

Use of DNA Markers in Breeding for Bunt Resistance at the National Agricultural Research and Development Institute Fundulea, Romania

M. CIUCA, F. ONCICA, M. IUORAS, M. ITTU and N.N. SAULESCU

Agricultural Research-Development Institute (ARDI) Fundulea, Romania, N. Titulescu
Street, No. 1, Calarasi, Fundulea, 912500, Romania, e-mail: matilda72001@yahoo.co.uk

Abstract: Breeding high-yielding bunt resistant wheat cultivars faces many difficulties related to: (1) The poor agronomic performance of most known sources of resistance. Our data show that, even after several cycles of crossing with high-yielding cultivars, the association of bunt resistance with some unfavourable traits remains difficult to break; (2) The incomplete expressivity of susceptibility alleles. Even under artificial inoculation, susceptible genotypes are not 100% diseased, and this creates difficulties in selection for resistance; (3) The need to pyramid genes giving complete resistance for achieving durability. Our work, aiming at the use of molecular markers for alleviating these problems, has been directed towards: (1) Use of already published marker for the gene *Bt10* to understand the resistance of breeding lines obtained by crossing with PI178383 (a *Bt9 + Bt10* carrier); (2) Work for identifying markers for the gene *Bt8*. A set of 23 randomly extracted lines from the cross F 676T2-1 (Yala 305/2* Aura//2* Rapid – *Bt-8* resistant) × Delabrad was phenotyped for bunt resistance, during 2 years. Resistance and susceptible lines have been analyzed with 56 primers. Preliminary results will be presented; (3) Preliminary identification of DNA polymorphism, possibly associated with bunt resistance genes *Bt11*, *Bt12*, *Bt5*, based on a comparison between resistance lines selected after several cycles of crossing and the respective susceptible parents. Results obtained so far will be used for a detailed analysis in specially built genetic population.

Keywords: wheat breeding; molecular markers; DNA polymorphism; *Bt* genes