

## Altitude and Forest Type Effects on Soils in the Jizera Mountains Region

LENKA PAVLŮ, LUBOŠ BORŮVKA, ANTONÍN NIKODEM, MARCELA ROHOŠKOVÁ  
and VÍT PENÍŽEK

*Department of Soil Science and Geology, Faculty of Agrobiological Sciences, Food and Natural Resources, Czech University of Life Sciences in Prague, Prague, Czech Republic*

**Abstract:** This paper is focused on the Jizera Mountains as a region strongly influenced by man in the past. The structure of the natural forest was changed. Species monocultures with similar tree ages were planted. High acidificants concentrations in atmosphere led to the decline of these monoculture forests in the top parts of the mountains and the high acidificants deposition damaged the soils in the whole region. The goals of this study are to describe the distribution of the soil properties in altitude transects, where temperature, precipitation, and vegetation gradients are recorded, and to compare the soil properties in spruce and beech forests. The soil samples were collected from soil pits in a surviving nature-close beech forest, in a production spruce forest, and also in the top dead forest area with a grass cover. Soil samples from sufficiently deep diagnostic horizons were taken for the study of chemical properties. The basic soil characteristics were determined by the commonly used methods (pH, effective cation exchange capacity – eCEC, and the contents of cations in the sorption complex,  $A_{400}/A_{600}$  as humus quality parameter, the contents of available Ca, Mg, K and P, pseudototal content of Ca and Mg, and two differently extracted Fe and Al forms contents). The soils of the Jizera Mts. are strongly acid with a low eCEC which is the result of the natural and anthropogenic acidification processes. Soil chemical properties of the most affected top mountainous parts are in some aspects more favourable than lower parts (binding of potentially toxic Al in organic matter, slightly higher pH), but in other aspects they are still endangered by the acidification symptoms (higher leaching of base cations, especially Mg). The soils of nature-close beech forests represent more favourable soil properties than those of planted spruce forests. Generally, it can be concluded that the natural systems have higher resilience, and that natural mechanisms are able to mitigate slightly the soil degradation.

**Keywords:** forest soils; altitude; forest type; beech; spruce; elevation transects

Soil acidification in the Czech Republic presents a serious problem in forest soils of mountainous areas. Soil acidification is a natural process conditioned by naturally acid parent rocks, high precipitations and slow organic matter decomposition producing various organic acids. This natural

system has been disturbed by human activities. Natural stable forest ecosystems were changed to spruce monoculture production forests with acid, hardly decomposed litter. This step led to natural acidification processes acceleration; soil pH decrease and base cation loss increase. A number

---

Supported by the Ministry of Agriculture of the Czech Republic, Project No. 1G57073, and by the Ministry of Education, Youth and Sports of the Czech Republic, Project No. MSM 6046070901.

of studies showed the differences in the soil characteristics between broadleaved and coniferous trees, especially between beech (*Fagus sylvatica*) and Norway spruce (*Picea abies*) (for example: CRONAN & GRIGAL 1995; ZIMKA & STACHURSKI 1996; MISSON *et al.* 2001; OULEHLE & HRUŠKA 2005). In 1970's and 1980's, a period of enormous coal mining, thermal power stations and heavy industry development, huge amounts of acidificants ( $\text{SO}_2$ ,  $\text{NO}_x$ , etc.) were emitted to the atmosphere. High levels of acidificants deposition increase the intensity of soil acidification and change some mechanisms of this process. The presence of mineral non-complexing acids ( $\text{H}_2\text{SO}_4$ ,  $\text{HNO}_3$ ) in soils leads to mobilisation of potentially toxic elements like Al or heavy metals (RICHTER 1986; HOVMAND & BILLE-HANSEN 1999; PIERZYNSKI *et al.* 2000; HRUŠKA & CIENCIALA 2003). Though sulphur emissions decreased dramatically in the last decades, the effect of S accumulation is long-termed; moreover, the nitrogen deposition is still fairly high (HRUŠKA *et al.* 2001; HRUŠKA & KRÁM 2003; UHLÍŘOVÁ *et al.* 2002).

This paper deals with the Jizera Mountains region located in the North of Bohemia. The region is one of the areas most affected by human activities in the Czech Republic (SUCHARA & SUCHAROVÁ 2002). High concentrations of acidificants in atmosphere, originating mainly from thermal power stations in the Czech Republic and in Poland, also called “Black triangle”, occurred in the past.

At present, the concentrations of acidificants in atmosphere are decreased. However, the forests still stay threatened by long-term changes of the soil conditions (lower base saturation, lower pH, and high contents of potentially toxic Al forms). In the Jizera Mountains, the damage of soil and forests led to the forest decline in the top parts of the mountains. This area was fast invaded by grass as a natural mechanism of ecosystem restoration. The soils under this grass cover have often better chemical properties than soils covered by forests due to the grass litter quality, its influence on N cycling, organic matter turnover and base cations leaching (ADAMS & EVANS 1989; KOIJMAN *et al.* 2000; FIALA *et al.* 2001).

This study is a continuation of the project dealing with the whole region of the Jizera Mts. Some results of the complex project showed effects of altitude and forest type on the soil properties distribution (BORŮVKA *et al.* 2005; MLÁDKOVÁ *et al.* 2005, 2006). The goals of this study are to focus on this relationship in greater detail, to describe the distribution of soil properties in altitude transect, and to compare the soil properties in spruce and beech forests.

## MATERIAL AND METHODS

### Area of interest and sampling

The Jizera Mountains region is plutonic area with uniform granite bedrock. Complex soil studies of this region are presented e.g. in MLÁDKOVÁ

Table 1. Description of sampling sites on Paličnick and Smědava transects

Pit	Paličnick			Smědava		
	soil type	forest	age	soil type	forest	age
1	Haplic Podzol	spruce	40–50	Haplic Podzol	spruce (+ pine)	10
2	Haplic Podzol	beech	80–120	Haplic Podzol	spruce	10
3	Haplic Podzol	spruce	10	Haplic Podzol	spruce	10
4	Haplic Podzol	beech	80	Colluvic Regosol	spruce	10
5	Haplic Podzol	spruce	30–40	Colluvic Regosol	spruce/beech	10/110
6	Haplic Podzol	beech	170	Colluvic Regosol	beech	110
7	Haplic Podzol	spruce	60	Haplic Podzol	beech	150
8	Entic Podzol	beech	80–120	Haplic Podzol	beech (+ birch)	150
9	Entic Podzol	spruce	90	Entic Podzol	beech	140
10	Cambisol	beech	120	Entic Podzol	beech	150




Figure 1. Map of sampling sites with colour emphasise of steepness (degree); scale change with distance, equidistance is 25 m (Paličnick – 972 m, Smědavská hora – 1084 m)

*et al.* (2004, 2005, 2006), BORŮVKA *et al.* (2005). This paper deals with two elevation transects in the north part of this region (Figure 1, Table 1). The first transect, Paličnick, represents the altitude range from 600 m to 950 m with western orientation. The soil samples were collected from ten soil pits. Five soil pits are placed in a different age spruce forest (*Picea abies*) and five in a more than 100 years old nature-close beech forest (*Fagus sylvatica*). The soil type changes with decreasing altitude from Haplic Podzols over Entic Podzols to Cambisols. The second transect, Smědava, represents the altitude range from 600 m to 1050 m with northern orientation. The soil samples were collected also from ten soil pits. The forest cover differs with decreasing altitude from the dead forest with young free-growing spruce and a high grass abundance to old beech forest. Haplic Podzols are the prevailing soil types. The middle steepest part of the slope contained Colluvic Regosols while that of the altitude of around 600 m Entic Podzols.

#### Basic soil characteristics

The soil samples were collected from sufficiently deep soil horizons: organic fermented (F), organic humified (H), mineral humic (A), mineral albic (eluvial) (E), mineral spodic humic-sesquioxidic, spodic sesquioxidic, spodic red, and cambic (B) soil horizons, and the substrate (C). In total, 104 samples

were collected. The samples were air dried in room temperature and sieved through 2 mm mesh.

Basic soil characteristics were determined. The values of  $\text{pH}_{\text{H}_2\text{O}}$ ,  $\text{pH}_{\text{CaCl}_2}$ , and  $\text{pH}_{\text{KCl}}$  were measured potentiometrically. Humus quality was assessed by the ratio of absorbances of pyrophosphate soil extract at the wavelengths of 400 nm and 600 nm ( $A_{400}/A_{600}$  – POSPÍŠIL 1981). Organic carbon content ( $C_{\text{org}}$ ) was determined oxidimetrically by a modified Tjurin method (POSPÍŠIL 1964); this analysis was done only on samples from the mineral horizons. Available nutrient contents were determined in Mehlich III extract ( $P_{\text{MIII}}$ ,  $K_{\text{MIII}}$ ,  $\text{Ca}_{\text{MIII}}$ ,  $\text{Mg}_{\text{MIII}}$ ) (ZBÍRAL 2002). Pseudo-total contents of Ca and Mg were measured after soil digestion with aqua regia ( $\text{Ca}_{\text{AR}}$ ,  $\text{Mg}_{\text{AR}}$ ) (ZBÍRAL 2002). Effective cation exchange capacity (eCEC) was determined by the Mehlich method with unbuffered 0.1M  $\text{BaCl}_2$  extraction solution (PODLEŠÁKOVÁ *et al.* 1992). The contents of exchangeable elements were measured together with eCEC ( $\text{Ca}_{\text{exc}}$ ,  $\text{Mg}_{\text{exc}}$ ,  $\text{K}_{\text{exc}}$ ,  $\text{Na}_{\text{exc}}$ ,  $\text{Mn}_{\text{exc}}$ ,  $\text{Al}_{\text{exc}}$ ). The contents of two different Al forms were determined according to DRÁBEK *et al.* (2003): potentially toxic Al forms were extracted with 0.5M KCl solution ( $\text{Al}_{\text{KCl}}$ ), and organically bound Al forms were extracted with 0.05M  $\text{Na}_4\text{P}_2\text{O}_7$  solution ( $\text{Al}_{\text{Na}_4\text{P}_2\text{O}_7}$ ). The contents of iron forms were measured in the same extracts as Al ( $\text{Fe}_{\text{KCl}}$ ,  $\text{Fe}_{\text{Na}_4\text{P}_2\text{O}_7}$ ). Aluminium and

iron concentrations in both extracts were determined by means of ICP-OES (inductively coupled plasma – optical emission spectrometry, VARIAN Vista Pro, VARIAN, Australia). Statgraphics Plus for Windows 4.0 (Manugistics 1997) was used to perform correlation analysis and *t*-tests.

## RESULTS AND DISCUSSION

Maximum and minimum values of the studied soil properties are shown separately for each transect in Tables 2–5. For a better illustration, Figures 2 and 3 show the distribution of the selected soil characteristics on both transects in the H horizon. This horizon was picked out because of its similarity in all soil pits from the pedogenesis point of view.

Acidity of the surface soil horizons ranges from strong to very strong (Table 2). Deeper horizons are less acid. Humus quality is rather low as it is typical for forest soils (Table 2). Table 3 shows the characteristics of the soil sorption complex. Soil sorption capacity decreases with the depth of the soil horizon, probably due to the decreasing of organic matter content. It is evident that the main part of the sorbed cations is composed of aluminium ions. The ratio of the base cations to

Al ions in the soil sorption complex is less than 1. However, the criteria of this ratio evaluation (where commonly used critical value is 1) exist only for the soil solution (CRONAN & GRIGAL 1995). The contents of the available elements decrease with depth (Table 4). While the pool of these elements is almost sufficient in the surface organic horizon, it is very low in the mineral horizons. The contents of  $P_{MIII}$  are often below the detection limit, and also  $Mg_{MIII}$  contents in the deepest mineral horizons are below this limit. The distribution of  $Mg_{AR}$  in the soil profile does not correspond to  $Mg_{MIII}$  distribution mentioned above. Pseudototal Mg content increases with depth and this distribution corresponds more to the high leaching in acid conditions. The distribution of  $Ca_{AR}$  corresponds probably to the distribution principles of the podzolization process. The lowest contents are found in eluvial E horizon. Al and Fe forms distribution in the soil profile is controlled by the podzolization process (Table 5). Their release from the E horizon and transport to the B horizon is the principal mechanism of this process. The relative assessment of this process intensity and the contents of the released potentially toxic Al forms could be a good indicator of the acidification level.

Table 2. Maximum and minimum values of studied basic soil characteristics on the Paličnick and Smědava transects separately for soil horizons

Horizon (samples)	pH <sub>H<sub>2</sub>O</sub>		pH <sub>KCl</sub>		pH <sub>CaCl<sub>2</sub></sub>		C <sub>org</sub> (%)		A <sub>400</sub> /A <sub>600</sub>	
	min	max	min	max	min	max	min	max	min	max
<b>Paličnick transect</b>										
F (10)	3.57	4.49	2.66	3.28	2.80	3.40			7.22	25.65
H (10)	3.09	4.02	2.83	3.42	2.90	3.50			7.71	17.00
A (5)	3.51	4.06	2.87	3.43	3.10	3.70	5.73	10.52	3.70	7.86
E (4)	3.69	3.87	3.04	3.25	3.25	3.45	4.01	7.86	4.88	6.04
B (19)	3.64	4.54	2.99	4.19	3.25	4.30	3.01	9.55	5.79	11.82
C (7)	4.45	4.63	4.05	4.25	4.10	4.45	0.98	4.48	3.57	12.19
<b>Smědava transect</b>										
F (8)	3.85	4.38	2.89	3.51	3.05	3.65			7.19	19.67
H (10)	3.85	4.15	3.02	3.64	3.15	3.80			6.20	10.50
A (5)	3.76	4.17	3.13	3.57	3.30	3.75	6.73	10.27	4.12	11.32
E (4)	3.95	4.33	3.33	3.61	3.55	3.85	1.67	7.54	4.68	8.42
B (8)	3.96	4.56	3.37	4.27	3.60	4.55	1.15	9.60	4.38	10.47
C (6)	4.44	4.59	3.89	4.35	4.15	4.55	1.10	5.14	3.73	10.53




Figure 2. Selected H horizon soil properties distribution on Paličník transect separately for beech (BK) and spruce (SM) forest (exchangeable cations contents (K<sub>ex</sub>, Mg<sub>ex</sub> and Ca<sub>ex</sub>) are presented in mmol/100 g, all other elements contents are presented in mg/kg).




Figure 3. Selected H horizon soil properties distribution on Smědava transect (exchangeable cations contents (K<sub>ex</sub>, Mg<sub>ex</sub> and Ca<sub>ex</sub>) are presented in mmol/100 g, all other elements contents are presented in mg/kg).



Table 3. Maximum and minimum values of soil sorption parameters on the Paličník and Smědava transects separately for soil horizons (eCEC and exchangeable cations contents are presented in mmol/100 g;  $Al_{ex}$  detection limit is 0.5 mmol/100 g)

Horizon (samples)	eCEC		$Ca_{ex}$		$Mg_{ex}$		$K_{ex}$		$Na_{ex}$		$Mn_{ex}$		$Al_{ex}$	
	min	max	min	max	min	max	min	max	min	max	min	max	min	max
<b>Paličník transect</b>														
F (7)	20.39	26.05	1.68	7.25	0.70	1.19	0.54	1.37	0.07	0.10	0.03	0.12	9.39	21.35
H (10)	10.13	27.03	0.37	1.28	0.41	0.86	0.34	0.80	0.06	0.09	0.01	0.05	3.76	18.38
A (5)	8.89	12.07	0.20	0.66	0.33	0.53	0.17	0.23	0.05	0.06	0.01	0.08	6.27	8.45
E (4)	7.21	10.45	0.35	0.62	0.15	0.49	0.10	0.22	0.05	0.08	0.01	0.01	4.56	9.20
B (19)	4.13	14.80	0.06	6.60	0.10	0.64	0.05	0.23	0.04	0.11	0.00	0.09	2.98	17.34
C (7)	1.64	4.41	0.09	0.43	0.23	0.53	0.04	0.08	0.03	0.16	0.01	0.03	<0.5	3.16
<b>Smědava transect</b>														
H (7)	10.06	24.00	0.37	1.84	0.68	1.47	0.22	0.81	0.04	0.10	0.01	0.30	6.07	24.44
A (3)	11.10	11.91	0.46	0.69	0.25	0.95	0.15	0.22	0.05	0.06	0.01	0.05	7.20	9.92
E (4)	2.87	9.82	0.28	0.45	0.21	1.25	0.05	0.11	0.04	0.06	0.01	0.03	0.24	8.65
B (8)	1.08	12.92	0.16	0.74	0.15	1.25	0.05	0.26	0.04	0.05	0.00	0.24	1.99	11.87
C (6)	0.88	5.81	0.16	0.48	0.11	0.69	0.05	0.07	0.04	0.05	0.01	0.05	0.80	5.85

#### Assessment of altitude effect

The altitude effect was assessed by correlation analysis. The results from the Paličník transect are shown in Table 6. The content of  $Al_{KCl}$  in the surface organic horizons decreases with increasing

altitude in the Paličník transect. It could indicate that a higher portion of Al is bound in organic matter in the higher parts of the transect. Negative correlations of  $Al_{KCl}$  contents in A and E horizons could be explained rather by a stronger podzolization process in the higher parts of the transect.

Table 4. Maximum and minimum values of available and pseudototal contents of studied elements on the Paličník and Smědava transects separately for soil horizons (all values are presented in mg/kg; detection limit is 10 mg/kg for available contents)

Horizon (samples)	$P_{MIII}$		$K_{MIII}$		$Mg_{MIII}$		$Ca_{MIII}$		$Mg_{AR}$		$Ca_{AR}$	
	min	max	min	max	min	max	min	max	min	max	min	max
<b>Paličník transect</b>												
F (10)	< 10	58	199	481	70	196	225	1630	524	1444	237	2058
H (10)	< 10	62	84	290	28	100	124	322	321	1871	201	525
A (5)	< 10	28	60	97	20	39	89	188	1136	2289	245	346
E (4)	< 10	<10	33	93	11	25	96	136	647	1567	129	330
B (19)	< 10	22	20	102	<10	58	38	147	1258	4031	167	544
C (7)	< 10	26	18	37	<10	50	55	123	3092	6303	441	1102
<b>Smědava transect</b>												
F (8)	< 10	25	164	677	47	186	129	906	809	1857	353	1380
H (10)	< 10	13	84	298	25	130	92	377	515	2318	255	615
A (5)	< 10	<10	46	82	19	29	72	153	904	3915	319	517
E (4)	< 10	<10	16	37	12	23	77	242	418	1450	248	316
B (8)	< 10	<10	16	94	<10	25	45	152	1516	4248	268	477
C (6)	< 10	23	19	28	<10	15	52	109	1246	5462	289	642

Table 5. Maximum and minimum values of studied Al and Fe forms contents on the Paličník and Smědava transects separately for soil horizons (all values are presented in mg/kg)

Horizon (samples)	Al <sub>KCl</sub>		Al <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>		Fe <sub>KCl</sub>		Fe <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>	
	min	max	min	max	min	max	min	max
<b>Paličník transect</b>								
F (10)	212.2	1 710.0	1 057.3	9 216.9	18.1	679.2	1 466.3	5 388.8
H (10)	310.2	1 633.4	1 885.8	13 229.7	18.3	453.7	2 498.4	8 132.3
A (5)	351.6	969.7	1 094.7	5 656.9	91.1	258.7	1 613.2	7 445.7
E (4)	307.6	737.4	1 074.7	1 677.8	62.2	173.0	347.0	3 559.1
B (19)	230.7	1 089.7	2 033.5	12 723.5	9.6	595.6	1 862.7	19 627.5
C (7)	140.5	281.3	2 188.5	7 006.7	8.2	19.6	300.6	3 825.4
<b>Smědava transect</b>								
F (8)	220.1	929.2	1 602.9	7 707.1	158.4	1 075.2	2 401.8	6 872.6
H (10)	24.8	1 318.1	3 344.4	8 565.0	31.8	1 266.6	4 764.9	8 490.6
A (5)	83.0	999.7	1 608.9	6 263.9	250.8	759.0	4 009.3	7 260.6
E (4)	69.1	887.7	100.2	1 646.6	43.5	417.8	183.6	1 553.2
B (8)	10.6	1 304.4	3 143.1	10 582.8	8.1	434.4	341.8	11 427.2
C (6)	9.4	486.4	1 654.9	7 350.0	6.0	371.8	132.8	5 034.1

The same process could explain the positive correlation of Fe<sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub> content with altitude in the B horizons. The results also showed negative correlation between Mg<sub>AR</sub> and altitude in the surface organic and mineral A and E horizons. Higher precipitation amounts in higher altitude and the consequent higher leaching could be responsible for this fact. However, no correlation was found in the case of Mg<sub>MIII</sub>, where sorption on organic matter could play an opposite role.

The results of correlation analysis for the Smědava transect are shown in Table 6. Mg<sub>AR</sub> content in the surface horizons decreases with increasing altitude in the Smědava transect as well as in the Paličník transect. The B horizons are also enriched with Fe<sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub> and, moreover, with Al<sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub> in the Smědava transect. Al<sub>KCl</sub> content decreasing with increasing altitude was not found in the surface horizons of this transect, but it was found in the A and E horizons.

The situation that correlation is in many cases rather weak can be caused in part by the fact that the relationship does not need necessarily to be linear. The search for a non-linear function describing the relationship between altitude

and soil properties will be underway in further research.

Generally, we can say that altitude has no effect by itself. Different altitudes represent changes in temperature, precipitation amount, vegetation cover, deposition rate, and so on. Soil chemical properties of the most affected top mountainous parts are in some aspects more favourable than those of lower parts (binding of potentially toxic Al in organic matter, slightly higher pH). Natural mechanisms as grass expansion in dead forests are able to mitigate slightly the soil degradation. In other aspects, this mountainous parts are still endangered by acidification symptoms (higher leaching of base cations, especially Mg).

#### Assessment of forest type effect

The Paličník transect to altitude 900 m was used for the assessment of the forest type effect on soil chemical properties. The soil pits placed above 900 m appertain to different vegetation zones by forest typology and for this reason they were removed from the data set evaluated. The results of the *t*-test performed are shown in Table 7. Beech

Table 6. Correlation coefficients of relationships of altitude versus studied soil properties on the Paličník and Smědava transects separately for soil horizons or groups of soil horizons; significant correlations at the significance level of 0.05 are in bold letters

Characteristic	All horizons	F + H	F	H	A+E	B	C
<b>Paličník transect</b>							
pH <sub>H<sub>2</sub>O</sub>	−0.048	0.111	0.386	−0.193	−0.002	−0.123	0.535
pH <sub>KCl</sub>	−0.111	−0.030	0.031	−0.090	−0.009	−0.216	0.395
pH <sub>CaCl<sub>2</sub></sub>	−0.128	−0.019	0.057	−0.095	−0.152	−0.267	−0.214
C <sub>org</sub>	0.267				−0.186	0.240	0.345
A <sub>400</sub> /A <sub>600</sub>	−0.166	−0.173	−0.213	−0.147	−0.530	−0.062	−0.163
Al <sub>KCl</sub>	−0.263	<b>−0.509</b>	−0.610	−0.396	<b>−0.675</b>	0.019	−0.237
Al <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>	0.171	0.201	0.080	0.301	−0.062	0.171	<b>0.907</b>
Fe <sub>KCl</sub>	−0.228	−0.367	−0.565	−0.133	−0.253	−0.252	0.632
Fe <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>	0.217	−0.217	−0.550	−0.083	−0.203	<b>0.548</b>	0.619
P <sub>MIII</sub>	0.226	0.342	0.555	0.212	0.596		
K <sub>M III</sub>	0.059	0.210	0.240	0.320	0.031	<b>0.513</b>	0.314
Mg <sub>MIII</sub>	−0.133	0.091	0.301	−0.251	−0.371	<b>−0.824</b>	−0.439
Ca <sub>MIII</sub>	0.119	0.254	0.377	0.169	0.307	0.242	−0.315
Mg <sub>AR</sub>	<b>−0.286</b>	<b>−0.606</b>	−0.562	<b>−0.672</b>	<b>−0.687</b>	−0.361	−0.359
Ca <sub>AR</sub>	0.021	0.238	0.360	0.129	−0.230	−0.196	0.091
eCEC	−0.054	−0.049	−0.264	0.166	<b>−0.717</b>	0.318	−0.112
Ca <sub>exc</sub>	0.131	0.206	0.598	0.573	0.458	0.231	0.482
Mg <sub>exc</sub>	−0.050	0.059	0.155	0.339	−0.247	0.364	−0.441
K <sub>exc</sub>	−0.081	−0.035	0.164	0.156	−0.067	0.445	0.093
Na <sub>exc</sub>	<b>0.283</b>	0.117	0.408	0.274	0.197	<b>0.552</b>	0.480
Mn <sub>exc</sub>	−0.107	−0.272	−0.184	−0.181	0.126	0.162	−0.021
Al <sub>exc</sub>	0.114	0.016	0.275	−0.041	−0.155	0.398	0.518
<b>Smědava transect</b>							
pH <sub>H<sub>2</sub>O</sub>	−0.145	−0.177	0.053	−0.504	0.239	−0.281	0.083
pH <sub>KCl</sub>	−0.177	0.105	0.469	−0.056	0.344	−0.459	−0.704
pH <sub>CaCl<sub>2</sub></sub>	−0.227	−0.020	0.317	−0.180	−0.004	−0.504	−0.579
C <sub>org</sub>	−0.040				−0.276	0.198	0.252
A <sub>400</sub> /A <sub>600</sub>	0.023	0.355	<b>0.748</b>	−0.406	−0.383	−0.625	0.007
Al <sub>KCl</sub>	−0.231	−0.196	0.292	−0.409	<b>−0.795</b>	0.268	−0.692
Al <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>	0.177	0.389	0.532	0.413	−0.016	<b>0.713</b>	0.234
Fe <sub>KCl</sub>	<b>0.385</b>	<b>0.475</b>	0.282	0.602	−0.021	0.589	0.833
Fe <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>	0.135	0.365	0.620	0.331	−0.196	<b>0.835</b>	0.020
P <sub>MIII</sub>	−0.115	−0.379	<b>−0.881</b>				
K <sub>M III</sub>	0.142	0.204	0.114	0.432	−0.138	0.212	0.150
Mg <sub>MIII</sub>	0.181	0.315	0.087	<b>0.676</b>	0.437	0.154	−0.182
Ca <sub>MIII</sub>	0.141	0.047	−0.268	0.630	0.508	<b>0.734</b>	<b>0.828</b>
Mg <sub>AR</sub>	<b>−0.449</b>	<b>−0.706</b>	−0.532	<b>−0.820</b>	−0.514	0.001	−0.667
Ca <sub>AR</sub>	0.068	−0.017	−0.340	0.623	0.165	0.508	−0.310
eCEC	0.256	0.386		0.252	−0.637	0.629	0.507
Ca <sub>exc</sub>	<b>0.418</b>	0.610		<b>0.870</b>	0.094	<b>0.867</b>	0.581
Mg <sub>exc</sub>	−0.244	0.042		−0.233	−0.544	−0.603	−0.299
K <sub>exc</sub>	0.330	0.617		<b>0.781</b>	−0.184	0.128	0.716
Na <sub>exc</sub>	<b>0.454</b>	<b>0.904</b>		<b>0.889</b>	−0.342	0.173	0.553
Mn <sub>exc</sub>	−0.063	−0.469		−0.474	−0.398	0.465	0.200
Al <sub>exc</sub>	−0.133	−0.492		−0.482	<b>−0.839</b>	0.402	−0.112



Table 7. *t*-values of the difference between beech and spruce forests on the Paličník transect; plus values present higher levels of concrete characteristic in beech forest, minus values in spruce forest; significant differences at the significance level of 0.05 are in bold letters

Characteristic	F	H	Bhs	C
pH <sub>H<sub>2</sub>O</sub>	1.272	0.155	1.116	0.499
pH <sub>KCl</sub>	<b>4.141</b>	<b>2.979</b>	3.505	–1.543
pH <sub>CaCl<sub>2</sub></sub>	<b>6.306</b>	<b>3.017</b>	4.243	
C <sub>org</sub>			–1.293	<b>–6.934</b>
A <sub>400</sub> /A <sub>600</sub>	–1.470	0.648	1.134	–3.092
Al <sub>KCl</sub>	–2.080	<b>–3.102</b>	–2.820	4.108
Al <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>	0.940	1.145	1.673	–2.026
Fe <sub>KCl</sub>	<b>–3.243</b>	–1.284	<b>–8.222</b>	–1.382
Fe <sub>Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub></sub>	–0.140	–0.190	–0.170	–0.737
P <sub>MIII</sub>	0.017	–1.327		
K <sub>MIII</sub>	0.255	–1.956	1.223	–0.680
Mg <sub>MIII</sub>	0.255	–1.511	–2.280	–1.482
Ca <sub>MIII</sub>	–1.749	–2.040	–0.057	–0.693
Mg <sub>AR</sub>	1.010	0.434	0.389	–0.478
Ca <sub>AR</sub>	1.202	1.180	–0.005	–1.213
eCEC	<b>–3.088</b>	–0.834	0.384	0.706
Ca <sub>exc</sub>	–0.092	–0.096	<b>4.561</b>	
Mg <sub>exc</sub>	0.478	–0.126	–2.600	–0.728
K <sub>exc</sub>	0.588	–1.161	1.400	–0.200
Mn <sub>exc</sub>	0.000	–0.195		0.447
Al <sub>exc</sub>	–0.670	–0.251	1.165	–0.292

forest soils had significantly higher pH (except pH<sub>H<sub>2</sub>O</sub>) in the surface organic horizons than spruce forest soils. Also, they contained less Al<sub>KCl</sub> and Fe<sub>KCl</sub> and they had lower eCEC in these horizons than spruce forest soils. This may be in connection with the lower content of organic matter in beech forest caused by an easier decomposition of leaves. Beech forest soils contained more Ca<sub>AR</sub> and Mg<sub>AR</sub> in the surface organic horizons, but this result was not significant.

The higher C<sub>org</sub> content in the C horizon of spruce forest soil could indicate an easier migration of the low quality organic matter through the soil profile to deeper soil horizons. The soils of natural close beech forests represent more favourable soil properties than the soils of planted spruce forests on the same stand.

Generally, it can be concluded that natural systems presented by beech forest have higher resilience to anthropogenic changes.

## References

- ADAMS W.A., EVANS G.M. (1989): Effects of lime applications to parts of an upland catchment on soil properties and the chemistry of drainage waters. *European Journal of Soil Science*, **40**: 585–597.
- BORŮVKA L., MLÁDKOVÁ L., DRÁBEK O. (2005): Factors controlling spatial distribution of soil acidification and Al forms in forest soils. *Journal of Inorganic Biochemistry*, **99**: 1796–1806.
- CRONAN CH.S., GRIGAL D.F. (1995): Use of calcium/aluminum ratios as indicators of stress in forest ecosystems. *Journal of Environmental Quality*, **24**: 209–226.

- DRÁBEK O., BORŮVKA L., MLÁDKOVÁ L., KOČÁREK M. (2003): Possible method of aluminium speciation in forest soils. *Journal of Inorganic Biochemistry*, **97**: 8–15.
- FIALA K., TŮMA I., HOLUB P., TESAŘOVÁ M., JANDÁK J., PÁVKOVÁ A. (2001): Importance of grass cover in reduction of negative processes in soil affected by air pollution. *Rostlinná Výroba*, **47**: 377–382.
- HOVMAND M.F., BILLE-HANSEN J. (1999): Atmospheric input to Danish spruce forests and effects on soil acidification and forest growth based on 12 years measurements. *Water, Air and Soil Pollution*, **116**: 75–88.
- HRUŠKA J., CIENCIALA E. (2003): Long-term Acidification and Nutritional Degradation of Forest Soils – Limiting Factors of Forestry Today. Ministry of Environment of the Czech Republic, Prague.
- HRUŠKA J., KRÁM P. (2003): Modelling long-term changes in stream water and soil chemistry in catchments with contrasting vulnerability to acidification (Lysina and Pluhuv Bor, Czech Republic). *Hydrology and Earth System Sciences*, **7**: 525–539.
- HRUŠKA J., CUDLÍN P., KRÁM P. (2001): Relationship between Norway spruce status and soil water base cations/aluminum ratios in the Czech Republic. *Water, Air and Soil Pollution*, **130**: 983–988.
- KOOIJMAN A.M., EMMER I.M., FANTA J., SEVINK J. (2000): Natural regeneration potential of the degraded Krkonose forests. *Land Degradation and Development*, **11**: 459–473.
- Manugistics (1997): Statgraphics Plus for Windows User Manual. Manugistics, Inc., Rockville.
- MISSEON L., PONETTE Q., ANDRÉ F. (2001): Regional scale effects of base cation fertilization on Norway spruce and European beech stands situated on acid brown soils: soil and foliar chemistry. *Annals of Forest Science*, **58**: 699–712.
- MLÁDKOVÁ L., BORŮVKA L., DRÁBEK O. (2004): Distribution of aluminium among its mobilizable forms in soils of Jizera Mountains region. *Plant Soil and Environment*, **50**: 346–351.
- MLÁDKOVÁ L., BORŮVKA L., DRÁBEK O. (2005): Soil properties and toxic aluminium forms in acid forest soils as influenced by the type of vegetation cover. *Soil Science and Plant Nutrition*, **51**: 741–744.
- MLÁDKOVÁ L., BORŮVKA L., DRÁBEK O., VAŠÁT R. (2006): Factors influencing distribution of different Al forms in the Jizera Mountains forest soils. *Journal of Forest Science*, **52** (Special Issue): 87–92.
- OULEHLE F., HRUŠKA J. (2005): Tree species (*Picea abies* and *Fagus sylvatica*) effects on soil water acidification and aluminium chemistry at sites subjected to long-term acidification in the Ore Mts., Czech Republic. *Journal of Inorganic Biochemistry*, **99**: 1822–1829.
- PIERZYNSKI G.M., SIMS J.T., VANCE G.F. (2000): *Soils and Environmental Quality*. 2<sup>nd</sup> Ed. CRC Press LLC, Boca Raton.
- PODLEŠÁKOVÁ E., NĚMEČEK J., SIROVÝ V., LHOTSKÝ J., MACUROVÁ H., IVANEK O., BUMERL M., HUDCOVÁ O., VOPLAKAL K., HÁLOVÁ G., BLAHOVEC F. (1992): *Analysis of Soils, Waters and Plants*. VÚMOP, Prague. (in Czech)
- POSPÍŠIL F. (1964): Fractionation of humus substances of several soil types in Czechoslovakia. *Rostlinná Výroba*, **10**: 567–580.
- POSPÍŠIL F. (1981): Group- and fractional composition of the humus of different soils. In: *Transactions 5<sup>th</sup> Int. Soil Science Conf. Vol. 1*. Research Institute for Soil Improvement, Prague, 135–138.
- RICHTER D.D. (1986): Sources of acidity in some forested udults. *Soil Science Society of America Journal*, **50**: 1584–1589.
- SUCHARA I., SUCHAROVÁ J. (2002): Distribution of sulphur and heavy metals in forest floor humus of the Czech Republic. *Water, Air and Soil Pollution*, **136**: 289–316.
- UHLÍŘOVÁ H., FADRHOŇSOVÁ V., BÍBA M., LOCHMAN V. (2002): Deposition and movement in forest ecosystems of selected substances with connection to the food chain. *Chemické Listy*, **96**: 598–606. (in Czech)
- ZBÍRAL J. (2002): *Analysis of Soils I, II, III – Unified Operating Procedures*. SKZÚZ, Brno. (in Czech)
- ZIMKA J.R., STACHURSKI A. (1996): Forest decline in Karkonosze Mts. (Poland). Part II. An analysis of acidity and chemistry of atmospheric precipitation, throughfall and forest streamwaters. *Polish Journal of Ecology*, **44**: 153–177.

Received for publication March 23, 2007

Accepted April 13, 2007

---

*Corresponding author:*

Ing. LENKA PAVLŮ, Ph.D., Česká zemědělská univerzita v Praze, Fakulta agrobiologie, potravinových a přírodních zdrojů, katedra pedologie a geologie, Kamýcká 129, 165 21 Praha 6-Suchbátov  
tel.: + 420 224 382 632, fax: + 420 234 381 836, e-mail: pavlu@af.czu.cz

---