

Selenium content in tissues and meat quality in rabbits fed selenium yeast

A. DOKOUPILOVÁ¹, M. MAROUNEK^{1,2}, V. SKŘIVANOVÁ¹, P. BŘEZINA³

¹Institute of Animal Science, Prague-Uhříněves, Czech Republic

²Institute of Animal Physiology and Genetics, Czech Academy of Sciences, Prague, Czech Republic

³Department of Food Technology, Thomas Bata University in Zlín, Zlín, Czech Republic

ABSTRACT: Ten weaned rabbits were fed a basal (control) diet containing 0.12 mg Se/kg. In 10 rabbits the basal diet was supplemented with Se-enriched yeast to increase the Se concentration to 0.50 mg/kg. Rabbits were slaughtered at the age of 11 weeks. Samples of meat, liver and hair of 4 rabbits from each group were taken and analyzed. No effect of Se on growth, feed conversion and dressing out percentage was observed. Loin and hindleg meat, liver and hair of control rabbits contained 93, 98, 521 and 267 µg Se/kg, respectively (average values). In treated rabbits the corresponding Se concentrations were 400, 389, 1 414 and 914 µg/kg. Supranutritional Se supply had no effect on the activity of glutathione peroxidase in meat, and oxidative stability of meat expressed as production of thiobarbituric acid-reactive substances in meat stored for 3 and 6 days. Thus, the enrichment of meat with Se is the main benefit of Se supplementation of rabbit diets.

Keywords: rabbits; selenium; meat

Fifty years ago, selenium (Se) was recognized as an important trace element, essential for human and animal health (Schwarz and Foltz, 1957). Se in the form of selenocysteine is the central structural component of various selenoproteins. Selenoproteins with known functions include five glutathione peroxidases (GSH-Px), two deiodinases, thioredoxin reductases and selenophosphate synthetase (Behne and Kyriakopoulos, 2001). The five GSH-Px detoxify H₂O₂ and fatty acid derived hydroperoxides, thus contributing to the antioxidant defence against reactive molecules and free radicals, and complement the effects of vitamin E. Se enters the food chain through plants. Selenomethionine (SeMet) and selenocysteine are the most common organic Se sources in foods. SeMet is incorporated into general proteins by the same codon as that to methionine, thus it is feasible to enrich the meat of animals with Se when excessive SeMet is given to animals.

Adequate intake of Se is needed to decrease the risk of several serious diseases in humans (Hartikainen, 2005). The recommended dietary intake of Se for adults is 55 µg/day. In pregnant and lactating women the estimated value for adequate supply is 60 and 70 µg/day, respectively (Surai, 2006). In many regions of the world including Central Europe the recommended Se intake is not achieved (Rayman, 2004). Consequently, it is desirable to increase the Se content in animal products. The consumption of rabbit meat is not high in the Czech Republic, nevertheless the enrichment of rabbit meat with Se may increase its attractiveness for consumers. Previous studies showed that the feeding of supplemental organic Se increased Se content in animal tissues (Molnár et al., 1998; Pavlata et al., 2001; Leng et al., 2003). In addition, a correlation exists between the Se concentration and the activity of GSH-Px in meat (Scholz et al., 1981; Daun et al., 2001). Therefore, the supplementation

of diets with Se may also increase the oxidative stability of meat. Information on the effects of dietary Se on Se deposition and meat quality in rabbits is scarce. A study of Erdélyi et al. (2000) was focused on the effect of supranutritional Se on Se concentration in blood and liver, and GSH-Px activity in tissue samples. Thus, the aim of this study was to investigate deposition of Se in meat, and the effect of supplemental Se on meat quality in rabbits fed a diet enriched with organic Se.

MATERIAL AND METHODS

Animals and diets

Ten Hyplus rabbits, weaned at 5 weeks of age, were fed a granulated diet containing 0.12 mg Se/kg. In ten rabbits the basal diet was supplemented with Se-yeast (Sel-Plex, Alltech) to increase the Se concentration to 0.50 mg/kg. Rabbits were housed individually in stainless mesh cages. The environmental temperature was about 16°C and the humidity ca 65%. The rabbits had *ad libitum* access to feed and water. Ingredients and chemical composition of diets are shown in Table 1. Consumption of feeds was recorded individually. Animals were weighed in one-week intervals. No rabbit died in the course of the trial. Rabbits were slaughtered at 11 weeks of

age. Hot carcass weight including head, liver, kidney and perirenal fat was measured 15–30 min after slaughter (Blasco and Ouhayoun, 1996). Dressing out percentage was calculated as the proportion of hot carcass weight from live weight.

Sampling and analyses

Loin and hindleg meat, liver and hair of four rabbits selected by random from each group were analyzed. Samples of liver and meat were taken 24 h *post mortem* and stored at –40°C. Meat samples for GSH-Px assay were stored at –70°C. Feed and meat dry matter (DM) was determined by oven drying at 105°C, and fat by extraction with petroleum ether in Soxtec 1045 apparatus (Tecator Comp., Sweden). Contents of feed protein and fibre were determined employing instruments Kjeltec Auto 1030 Analyser and Fibertec 2010 from the same firm; ash after burning at 550°C. To determine Se, feeds, tissues, and hair were mineralized using the microwave digestion technique in a closed system (Milestone Ethos TC, Italy), in the presence of HNO₃ and H₂O₂. Se in processed samples was measured by the atomic absorption spectrometry (Solaar M-6 instrument, TJA Solutions, U.K.). The analytical procedure was validated by the analysis of a certified reference material RM 8414 Bovine Muscle (NIST).

Table 1. Ingredients and determined chemical composition of the basal¹ rabbit diet

Ingredients (%)		Chemical composition (g/kg)	
Lucerne meal	30	dry matter	906
Wheat bran	26	crude protein	166
Barley	14.5	crude fibre	161
Oat	6	fat	41
Sugar beet pulp	4	ash	73
Sunflower meal (extracted)	13	Se	1.2×10 ⁻⁴
Soybean meal (extracted)	2		
Rapeseed oil	1.5		
Mineral supplement ²	2		
Supplement of vitamins and trace elements ³	1		

¹Experimental diet was supplemented with Se at 0.38 mg/kg

²limestone, dicalcium phosphate, and salt

³per 1 kg: vitamin A 1 200 000 IU, vitamin D₃ 20 000 IU, alpha-tocopherol 5 g, vitamin K₃ 200 mg, vitamin B₁ 300 mg, vitamin B₂ 700 mg, vitamin B₆ 400 mg, vitamin B₁₂ 2 mg, niacin amide 5 g; calcium pantothenate 2 g, biotin 20 mg, folic acid 170 mg, choline 60 g, DL-methionine 100 g, L-lysine HCL 20 g, sodium salinomycinate 2 250 mg, cobalt 100 mg, copper 2 g, iron 5 g, iodine 120 mg, manganese 4.7 g, zinc 5 g, selenium 15 mg, antioxidant 10 g

Table 2. Growth, feed intake, feed conversion and carcass yield in rabbits¹ fed a basal diet and diet supplemented with Se-yeast (mean values ± SD)

	Control	Se
Initial weight (g)	1 015 ± 19	1 016 ± 25
Final weight (g)	2 944 ± 319	2 801 ± 302
Weight gain (g)	1 929 ± 320	1 785 ± 305
Feed intake (kg)	6.21 ± 0.96	5.56 ± 0.68
Feed/gain (kg/kg)	3.22 ± 0.14	3.11 ± 0.29
Carcass yield (%)	60.1 ± 1.3	60.0 ± 2.0

¹10 rabbits/group

Table 3. Dry matter (DM), fat and Se concentration in loin and hindleg meat of rabbits¹ fed a basal diet and diet supplemented with Se-yeast (mean values ± SD)

	Loin		Hindleg	
	control	Se	control	Se
DM (g/kg)	250 ± 4	250 ± 4	267 ± 12	266 ± 11
Fat (g/kg)	10.6 ± 1.5	8.7 ± 2.5	44.8 ± 13.8	37.7 ± 11.5
Se (μg/kg)	93 ± 3	400 ± 26*	98 ± 13	389 ± 34*

¹meat of 4 rabbits per group was analyzed

*significant effect of Se supplementation ($P < 0.05$)

The activity of GSH-Px was measured with tert-butyl hydroperoxide as a substrate by a coupled assay, recording the oxidation of NADPH by the decrease in absorbance at 340 nm. The activity was expressed as μmol NADPH oxidized min/g meat or liver tissue (DeVore and Greene, 1982). Lipid oxidation in minced meat samples was measured by the thiobarbituric acid method of Piette and Raymond (1999) and results were expressed as thiobarbituric acid-reactive substances (TBARS) in mg of malondialdehyde per kg muscle.

The *t*-test was used to evaluate the effects of Se supplementation.

RESULTS

No effect of Se on growth, feed conversion and dressing out percentage was observed (Table 2). Loin and hindleg meat of rabbits fed the Se-supplemented diet contained four times more Se than meat of control rabbits. The content of DM and fat in meat was not significantly influenced (Table 3). Concentrations of Se in the liver and hair exceeded those in meat. Supplementation with Se significantly increased Se concentration in liver and hair by 171 and 242%, respectively.

Table 4. Concentration of Se in liver, hair, and activity of GSH-Px in loin meat of rabbits¹ fed a basal diet and diet supplemented with Se-yeast (mean values ± SD)

	Control	Se
Se in liver (μg/kg)	521 ± 58	1 414 ± 48*
Se in hair (μg/kg)	267 ± 33	914 ± 142*
GSH-Px in meat (U/g) ²	1.29 ± 0.48	1.49 ± 0.50

¹meat of 4 rabbits per group was analyzed

²expressed as μmol NADPH oxidized min/g meat

*significant effect of Se supplementation ($P < 0.05$)

Table 5. Production of thiobarbituric acid-reactive substances (TBARS, mg MDA/kg) in loin and hindleg meat of rabbits¹ fed a basal diet and diet supplemented with Se (mean values ± SD)

	Loin		Hindleg	
	control	Se	control	Se
Day 0	0.94 ± 0.04	0.80 ± 0.14	0.83 ± 0.20	0.95 ± 0.22
Day 3	1.13 ± 0.26	1.13 ± 0.25	1.14 ± 0.17	1.73 ± 0.40
Day 6	1.70 ± 0.46	1.59 ± 0.32	1.86 ± 0.44	2.28 ± 1.10

¹samples from 4 rabbits per group were analyzed

MDA, malondialdehyde

There was no treatment effect on GSH-Px activity in loin meat (Table 4). Dietary Se did not influence the formation of TBARS in meat stored for 3 and 6 days at 4°C (Table 5).

DISCUSSION

The nutritional requirement of rabbits for Se is low, only 0.08 mg per kg of feed (NRC, 1977). Thus, it is not surprising that the Se supplementation did not significantly influence the rate of growth, feed efficiency and carcass yield. Se was well deposited in all tissues examined. Rabbits did not differ in this respect from other animals. The supplement of Se increased Se concentration in meat of broiler chickens (Ševčíková et al., 2006), pigs (Goehring et al., 1984), calves (Pavlata et al., 2001), and in eggs (Skřivan et al., 2006). One hundred and forty g of meat of Se-fed rabbits would cover the recommended Se daily intake for adults (Rayman, 2004). High Se exposure could be detrimental. However, it should be mentioned that toxic Se doses are more than 10 times higher than the physiological requirement (Surai, 2006). There was no treatment effect on DM and fat content of meat.

Erdélyi et al. (2000) did not observe a relationship between the Se status and the GSH-Px activity in femoral muscles of rabbits. On the other hand, several authors reported a correlation between GSH-Px activity and Se content of tissues of poultry (Daun and Åkesson, 2004), pigs (Daun et al., 2001) and cattle (Scholz et al., 1981; Gatellier et al., 2004). This relationship was not observed in our experiment with rabbits. According to Lee et al. (1979) rabbit liver and kidney contain a sufficient level of non Se-dependent GSH-Px, whereas lungs, heart, spleen, erythrocytes and plasma have only the Se-dependent GSH-Px activity. This makes rabbits poorly responsive to dietary Se deficiency.

The increase in the dietary Se level from 0.12 to 0.50 mg/kg did not decrease oxidation of lipids expressed as production of TBARS. Therefore, in rabbits the supranutritional Se supply has a limited potential for increasing the oxidative stability of meat. It can be concluded that the enrichment of meat with Se is the main benefit of Se supplementation of rabbit diets.

REFERENCES

- Behne D., Kyriakopoulos A. (2001): Mammalian selenium-containing proteins. *Annu. Rev. Nutr.*, 21, 453–473.
- Blasco A., Ouhayoun J. (1996): Harmonization of criteria and terminology in rabbit meat research. Revised proposal. *World Rabbit Sci.*, 4, 93–99.
- Daun C., Åkesson B. (2004): Comparison of glutathione peroxidase activity, and of total and soluble selenium content in two muscles from chicken, turkey, duck, ostrich and lamb. *Food Chem.*, 85, 295–303.
- Daun C., Johansson M., Önning G., Åkesson B. (2001): Glutathione peroxidase activity, tissue and soluble selenium content in beef and pork in relation to meat ageing and pig RN phenotype. *Food Chem.*, 73, 313–319.
- DeVore V.R., Greene B.E. (1982): Glutathione peroxidase in post-rigor bovine semitendinosus muscle. *J. Food Sci.*, 47, 1406–1409.
- Erdélyi M., Virág G., Mézes M. (2000): Effect of supranutritional additive selenium supply on the tissue selenium concentration and the activity of glutathione peroxidase enzyme in rabbit. *World Rabbit Sci.*, 8, 183–189.
- Gatellier P., Mercier Y., Renerre M. (2004): Effect of diet finishing mode (pasture or mixed diet) on antioxidant status of Charolais bovine meat. *Meat Sci.*, 67, 385–394.
- Goehring T.B., Palmer I.S., Olson O.E., Libal G.W., Wahlstrom R.C. (1984): Effect of seleniferous grains and

- inorganic selenium on tissue and blood composition and growth performance of rats and swine. *J. Anim. Sci.*, 59, 725–732.
- Hartikainen H. (2005): Biochemistry of selenium and its impact on food chain quality and human health. *J. Trace Elem. Med. Biol.*, 18, 309–318.
- Lee Y.H., Layman D.K., Bell R.R. (1979): Selenium-dependent and non selenium-dependent glutathione peroxidase activity in rabbit tissue. *Nutr. Rep. Int.*, 20, 573–578.
- Leng L., Bobcek R., Kuricová S., Boldizárová K., Grešáková L., Ševčíková Z., Révajová V., Levkutová M., Levkut M. (2003): Comparative metabolic and immune responses of chickens fed diets containing inorganic selenium and Sel-PlexTM organic selenium. In: Proc. Alltech's 19th Ann. Symp., Nottingham University Press, Nottingham, UK, 131–145.
- Molnár J., Macpherson A., Molnár P. (1998): The effects of selenium supplementation in feeding of lambs. *Acta Alim. Hung.*, 27, 167–179.
- NRC (1977): Nutrient Requirements of Rabbits. National Academy of Sciences. National Research Council, Washington, USA.
- Pavlata L., Illek J., Pechová A. (2001): Blood and tissue selenium concentrations in calves treated with inorganic or organic selenium compounds – a comparison. *Acta Vet. Brno*, 70, 19–26.
- Piette G., Raymond Y. (1999): Vergleichende bewertung verschiedener methoden. Bestimmung der Ranzigkeit in Fleischprodukten. *Fleischwirtschaft*, 7, 69–73.
- Rayman M.P. (2004): The use of high-selenium yeast to raise selenium status: how does it measure up? *Brit. J. Nutr.*, 92, 557–573.
- Scholz R.W., Todhunter D.A., Cook L.S. (1981): Selenium content and glutathione peroxidase activity in tissues of young cattle fed supplemented whole milk diets. *Am. J. Vet. Res.*, 42, 1718–1723.
- Schwarz K., Foltz C.M. (1957): Selenium as an integral part of factor 3 against dietary necrotic liver degeneration. *J. Am. Chem. Soc.*, 79, 3292–3293.
- Skřivan M., Šimáně J., Dlouhá G., Doučha J. (2006): Effect of dietary sodium selenite, Se-enriched yeast, Se-enriched Chlorella on egg Se concentration, physical parameters of eggs and laying hen production. *Czech J. Anim. Sci.*, 51, 163–167.
- Surai P.F. (2006): Selenium in Nutrition and Health. Nottingham University Press, Nottingham, UK, 643–808.
- Ševčíková S., Skřivan M., Dlouhá G., Koucký M. (2006): The effect of selenium source on the performance and meat quality of broiler chickens. *Czech J. Anim. Sci.*, 51, 449–457.

Received: 2007-01-03

Accepted after corrections: 2007-04-06

Corresponding Author

Ing. Adéla Dokoupilová, Institute of Animal Science, Přátelství 815, 104 01 Prague 10-Uhříněves, Czech Republic
Tel. +420 267 009 714, fax +420 267 710 779, e-mail: dokoupilova.adela@vuzv.cz
