

Survival Characteristics of *E. coli* O157:H7, *S. typhimurium* and *S. aureus* during Kefir Fermentation

NURAL KARAGÖZLÜ¹, CEM KARAGÖZLÜ² and BÜLENT ERGÖNÜL¹

¹Food Engineering Department, Engineering Faculty, Celal Bayar University, Manisa, Turkey; ²Department of Dairy Technology, Faculty of Agriculture, Ege University, İzmir, Turkey

Abstract

KARAGÖZLÜ N., KARAGÖZLÜ C., ERGÖNÜL B. (2007): **Survival characteristics of *E. coli* O157:H7, *S. typhimurium* and *S. aureus* during kefir fermentation.** Czech J. Food Sci., 25: 202–207.

In this research, the growth and survival of *E. coli* O157:H7, *Salmonella typhimurium* and *Staphylococcus aureus* were investigated during kefir fermentation. Two different levels of inoculation of the strains were conducted; the levels of 10² CFU/ml (EC-1, SA-1 and S-1) and 10³ CFU/ml (EC-2, SA-2 and S-2). At 0, 2, 6, 12, and 24 hours of kefir fermentation at 23 ± 1°C, samples were taken and the counts of *E. coli* O157:H7, *S. typhimurium*, and *S. aureus* were determined. EC-1 grew from 2.29 ± 0.02 log CFU/ml to 4.13 ± 0.18 log CFU/ml whereas EC-2 grew from 3.22 ± 0.04 log CFU/ml to 6.78 ± 0.99 log CFU/ml. Both S-1 and S-2 viable populations grew during the fermentation period, where sample S-1 grew from 2.37 ± 0.20 log CFU/ml to 4.64 ± 0.67 log CFU/ml and sample S-2 grew from 3.52 ± 0.07 log CFU/ml to 5.60 ± 0.10 log CFU/ml. SA-2 strains grew from 3.06 log CFU/ml to 3.64 log CFU/ml, SA-1 strains grew from 2.28 log CFU/ml to 2.66 log CFU/ml. According to the findings, *E. coli* O157:H7, *S. typhimurium*, and *S. aureus* can survive in kefir during fermentation.

Keywords: kefir fermentation; *E. coli* O157:H7; *Salmonella typhimurium*; *Salmonella aureus*

Kefir is a fermented milk product produced by the coculture of certain lactic acid bacteria, acetic acid bacteria, and yeasts. Kefir grains are small, irregularly shaped, yellowish-white particles that resemble miniature cauliflowers.

Kefir is used for the production of a refreshing probiotic fermented milk beverage, wellknown to East Europeans, by inoculating milk with kefir grains (GÜZEL-SEYDIM *et al.* 2000; WSZOLEK *et al.* 2001). The acidic fermented milk is slightly carbonated and contains small amounts of alcohol. Kefir differs from the traditional fermented milks (yoghurt) in that it is made only from kefir grains, which are composed of bacteria and yeasts (MARSHALL & COLE 1985). Kefir beverage contains

live microflora from the grains which are removed by filtration after the fermentation process. Kefir grains are conglomerates of lactic acid bacteria (*Lactobacilli*, *Lactococci*, and *Leuconostoc*), acetic acid bacteria (*Acetobacter aceti*), and yeasts (*Saccharomyces cerevisiae*, *Candida kefir*, *Kluyveromyces marxianus*) held together by a matrix of different exopolysaccharides.

Probiotics have been shown to possess inhibitory activity against the growth of pathogenic bacteria such as *Escherichia coli*, *Listeria monocytogenes*, *Salmonella* spp., and others (HARIS *et al.* 1989; CHATEAU *et al.* 1993). This inhibition is supposedly due to the production of inhibitory compounds such as bacteriocins, hydrogen peroxide, or organic acids,

as well as competitive adhesion to epithelium. Kefir also contains microorganisms that help to maintain a well-balanced intestinal flora and is known as an antitumour factor (ZUBILLAGA *et al.* 2001).

Kefir is made by fermentation of pasteurised milk. In the case of an inadequate pasteurisation, pathogen microorganisms such as *S. typhimurium*, *E. coli* O157:H7 and *S. aureus* may grow and multiply in milk and kefir. Since milking is realised by hand in open areas, especially in the countryside under poor hygienic conditions, milk can be contaminated by several sources such as diseased animals, milk handlers, and contaminated equipment (OGWARO *et al.* 2002). It is also known that raw milk is one of the sources of *E. coli* O157:H7. This pathogen has been isolated from raw milk (McDONOUGH *et al.* 1991; REITISMA & HENNING 1996). Therefore, the attention has to be extended to the contamination of dairy products.

In this study, the growth and survival of *E. coli* O157:H7, *Salmonella typhimurium*, and *Staphylococcus aureus* strains during kefir fermentation were examined.

MATERIAL AND METHODS

Preparation of kefir samples. Kefir grains were obtained from Microbiology Laboratory, Department of Dairy Technology, Faculty of Agriculture, Ege University, Izmir, Turkey. Sterilised milk (3% v/v fat) was purchased from Pinar Milk Company. *Escherichia coli* O157:H7, *S. typhimurium* (NRLE E.4463), and *S. aureus* strains were provided by the Food Microbiology Laboratory, Department of Food Engineering, Ege University, Izmir.

Seven different kefir samples were prepared by inoculating kefir grains into pasteurised milk in the proportion of 5% (w/v). One sample was inoculated with 10^2 CFU/ml of an overnight culture of *E. coli* O157:H7 EC-1. The second sample was inoculated with 10^3 CFU/ml of *E. coli* O157:H7 EC-2. The third (S-1) and fourth (S-2) samples were inoculated with 10^2 CFU/ml and 10^3 CFU/ml of an overnight culture of *Salmonella typhimurium*, respectively. The fifth (SA-1) and sixth (SA-2) kefir samples were inoculated, respectively, with 10^2 CFU/ml and 10^3 CFU/ml of an overnight culture of *Staphylococcus aureus*. The last sample (C) was not inoculated and served as the control sample. Immediately after inoculation, the samples were incubated at $23 \pm 1^\circ\text{C}$ for 24 h in an incubator (Memmert, model 400). The samples were then

collected after 0, 2, 6, 12, and 24 h and microbiological analysis was performed.

pH measurement. pH of the control sample was measured at the time of sampling. pH-meter (WTW pH537) was used for the measurement.

Microbiological counts. For the enumeration of *E. coli* O157:H7, *Salmonella typhimurium* and *Staphylococcus aureus*, the samples were serially diluted with peptone water (0.1% w/v), and the viable populations were determined by plating 0.1 ml of the diluted samples on Sorbitol Macconkey Agar (Oxoid, CM813), Brilliant Green Agar (Oxoid, CM263), and Baird Parker Agar (Oxoid, CM275), respectively. The samples were then incubated at 37°C for 24–48 hours. At the end of the incubation period, the colonies were counted and their numbers were determined (BAIRD-PARKER 1962; AOAC 2001; LARA *et al.* 2003; TSEGAYE & ASHENAFI 2005).

Statistical analysis. The mean values and the standard deviation of the data obtained from duplicate experiments were determined. Data were compared by Duncan's multiple range method (SAS 1999).

RESULTS AND DISCUSSION

Acidification is an important measure to control the growth and survival of pathogens and spoilage microorganisms. However, various acidic foods like yoghurt have been implicated in the foodborne outbreaks caused by *E. coli* O157:H7 (GRIFFIN & TAUXE 1991; ULJAS & INGHAM 1999; McCLURE 2000; HSIN-YI & CHOU 2001; GÜLMEZ & GÜVEN 2003; ŞİMSEK *et al.* 2007; BACHROURI *et al.* 2006). Since *E. coli* O157:H7 can survive in the pH range between 4.5 and 9.0 (GLASS *et al.* 1992; REINDERS *et al.* 2001), kefir can act as a medium for its growth and survival. In recent years, acid adaptation and increased resistance of *E. coli* O157:H7 (CHENG & CHOU 1999; McINGVALE *et al.* 2000) and *Salmonella* spp. (GOVERD *et al.* 1979; FOSTER & HALL 1990; ZHAO *et al.* 1993) have been observed by several researchers.

E. coli O157:H7 is known as acid resistant (ZHAO *et al.* 1993; BUCHANAN & DOYLE 1997; GURAYA *et al.* 1998) and it can survive in fermented dairy products for long periods of time (DIENEEN *et al.* 1998; TSEGAYE & ASHENAFI 2005). ZHAO *et al.* (1993) observed the survival of *E. coli* O157:H7 for up to 31 days in refrigerated unpasteurised apple juice at pH 3.9.

As reported by TSIAl and INGHAM (1997) and MILLER and KASPAR (1994), *E. coli* can survive in acidic foods such as apple cider (pH 3.7), mustard (pH 3.1), and ketchup (pH 3.6). It is also reported that acid adaptation prolonged the survival of some pathogens like *E. coli* O157:H7 and *Salmonella* spp., and may have important implications in terms of food safety (ACKERS *et al.* 1998; HSIN-YI & CHOU 2001; SAMELIS & SOFOS 2003).

It is reported that *E. coli* O157:H7 could grow and survive at pH 4.5 in a medium adjusted with HCl (CONNOR & KOTROLA 1995). According to UYTENDAELE *et al.* (2001), *E. coli* O157:H7 strains of approximately 10^2 CFU/ml, can survive for 5 days in meat, with pH adjusted with 50% lactic acid and fermented at 22°C. The growth and survival of *E. coli* O157:H7 during fermentation of kefir is shown in Table 1. Both EC-1 and EC-2 samples show similar growth patterns during fermentation. EC-1 grew from 2.29 ± 0.02 log CFU/ml to 4.13 ± 0.18 log CFU/ml whereas EC-2 grew from 3.22 ± 0.04 log CFU/ml to 6.78 ± 0.99 log CFU/ml. Similarly, the populations of *E. coli* O157:H7 strains in both samples were doubled in log units during the same period. During fermentation, pH of kefir decreased from 6.7 to 4.6 (Figure 1). In spite of this decrease in pH, viable populations of *E. coli* O157:H7 strains were increased. As seen in Table 1, the populations of the *E. coli* O157:H7 strains in both EC-1 and EC-2 significantly increased during the fermentation period ($P < 0.05$). This shows that stationary-phase strains of *E. coli* O157:H7 are acid tolerant and can survive during acid fermentation. Stationary-phase cells of *E. coli* O157:H7 are more resistant to acid than log-phase cells. Similar findings of *E. coli* O157:H7 survival during fermentation of yoghurt were reported by OGWARO *et al.* (2002). TSEGAYE *et al.* (2004) found that *E. coli* O157:H7 strains can




Figure 1. Changes in pH during kefir fermentation (control sample)

survive and grow 1 log unit during the first 24 h of fermentation of green Datta slurry (a traditional Ethiopian fermented beverage). Also *E. coli* O157:H7 was reported to grow in TSB at pH levels as low as 2.0 (KODIE & DHOPLE 2001). MILLER and KASPAR (1994) and ROERING *et al.* (1999) reported that *E. coli* O157:H7 had survived for several days in apple juice at pH < 3.5. According to MOON *et al.* (2006), *E. coli* O157:H7 was detected in apple juice at pH 3.5 during 10 days.

Survival of *E. coli* O157:H7 was also reported in fermented sausages (at pH 4.5) for 35 days (UYTENDAELE *et al.* 2001), in whey cheeses for 30 days (at pH 5.8) (GOVARIS *et al.* 2001), in fermented sausages for 8 weeks at pH 4.4 (GLASS *et al.* 1992), and in skim milk for 35 days at pH 4.7–5.4 (GURAYA *et al.* 1998). MAREK *et al.* (2004) revealed that *E. coli* O157:H7 can survive in pasteurised and unpasteurised whey samples up to 21 days and indicated the potential risk of persistence of *E. coli* O157:H7 in whey in the event of contamination with this pathogen. Also LEKKAS *et al.* (2006) re-

Table 1. Survival of *E. coli* O157:H7, *Salmonella* spp. and *Staphylococcus aureus* during fermentation of kefir (log CFU/ml)

Time (h)	EC-1	EC-2	S-1	S-2	SA-1L	SA-2
0	2.29 ± 0.02^c	$3.22 \pm 0.04^{*c}$	2.37 ± 0.20^c	$3.52 \pm 0.07^{*b}$	2.28 ± 0.03^d	$3.06 \pm 0.05^{*,d}$
2	$2.87 \pm 0.60^{b,c}$	3.40 ± 0.14^c	2.50 ± 0.28^c	$4.53 \pm 1.34^{a,b}$	2.46 ± 0.03^c	3.25 ± 0.03^c
6	$3.38 \pm 0.08^{a,b}$	$4.05 \pm 0.11^{b,c}$	$3.36 \pm 0.51^{b,c}$	$4.91 \pm 0.82^{a,b}$	2.60 ± 0.02^b	3.44 ± 0.02^b
12	3.77 ± 0.4^a	4.64 ± 0.15^b	$4.13 \pm 0.03^{a,b}$	5.57 ± 0.09^a	$2.64 \pm 0.02^{a,b}$	3.52 ± 0.01^b
24	4.13 ± 0.18^a	6.78 ± 0.99^a	4.64 ± 0.67^a	5.60 ± 0.10^a	2.66 ± 0.02^a	3.64 ± 0.02^a
	$P < 0.05$					

*Data represent the mean values of two independent experiments and their standard deviations

ported the survival of *E. coli* O157:H7 in Galotyri cheese at pH 3.7. TSEGAYE and ASHENAFI (2005) investigated the survival characteristics of *E. coli* O157:H7 in Ergo and Ayib, traditional fermented dairy beverages. In the study, it was revealed that during Ayib processing *E. coli* O157:H7 in souring milk increased by 3 log units in 24 hours. Also in Ayib production *E. coli* O157:H7 could survive for 7 days at pH 3.8–3.9.

GURAYA *et al.* (1998), found that *E. coli* O157:H7 can survive after a period of a week in non-fat yoghurt. BACHROURI *et al.* (2006) studied the survival of *E. coli* O157:H7 in milk and during refrigeration of home-made yoghurt. They indicated that the results of their study were of high public health significance, since the survival of *E. coli* O157:H7 was long enough to represent a hazard for the consumer. *E. coli* O157:H7 survival was also reported in various tomato products (4.2–4.8 pH) and it was indicated that the survival was notably prolonged at refrigeration temperatures (ERIBO & ASHENAFI 2003). In another study supporting ours, ROZAND *et al.* (2005), indicated that *E. coli* O157:H7 survives the lactic cheese manufacturing process, and that the presence of low numbers of *E. coli* O157:H7 in milk destined for the production of raw milk lactic cheeses can constitute a threat to the consumer.

The growth and survival of *Salmonella typhimurium* during fermentation of kefir are given in Table 1. Both S-1 and S-2 viable populations grew during the fermentation period, where the sample S-1 grew from 2.37 ± 0.20 log CFU/ml to 4.64 ± 0.67 log CFU/ml and the sample S-2 grew from 3.52 ± 0.07 log CFU/ml to 5.60 ± 0.10 log CFU/ml. S-1 and S-2 *S. typhimurium* strains showed resistance to increasing acidity (final pH was 4.6) and grew approximately 2 log units. The fermentation time was found statistically significant with the population of *S. typhimurium* in both kefir samples ($P < 0.05$). Also according to the findings of ROERING *et al.* (1999), *S. typhimurium* strains of 10^7 CFU/ml can survive well in apple cider (pH 3.3–3.5) for 21 days at 4°C.

Staphylococcus aureus is considered to be the second most common pathogen causing outbreaks of food poisoning, outnumbered only by *Salmonella* spp. (LARA *et al.* 2003). *Staphylococcus aureus* was demonstrated in dairy products (MARTIN & MARSHALL 1995; IURLINA & FRITZ 2004). Enterotoxins involved in food poisoning are produced by approximately one-third of the coagulase positive

strains of *S. aureus* (PORTOCARRERO *et al.* 2002). Enterotoxin A and Enterotoxin D of *Staphylococcus* are associated with many foodborne diseases.

The growth and survival of *Staphylococcus aureus* strains in the samples SA-1 and SA-2 are given in Table 1. In both SA-1 and SA-2 samples, *Staphylococcus aureus* strains survived and showed similar growth patterns during fermentation. The SA-2 strains grew from 3.06 log CFU/ml to 3.64 log CFU/ml, while the SA-1 strains from 2.28 log CFU/ml to 2.66 log CFU/ml, respectively. Growth populations of the strains were found less than 1 log unit, but as concerns the statistical analysis, the fermentation period was found significant for the growth of *Staphylococcus aureus* strains ($P < 0.05$).

CONCLUSION

The results of microbiological analysis revealed that *Salmonella typhimurium*, *E. coli* O157:H7, and *S. aureus* can survive and multiply during fermentation of kefir. Especially *E. coli* O157:H7 and *Salmonella typhimurium* doubled in log units. Since these pathogens are resistant to acidic conditions, in the case of the contamination of milk used for kefir, they can presumably survive and cause foodborne illnesses. To prevent the foodborne diseases caused by contaminated kefir, stricter preventive measures should be taken reaching from the dairy farm to the manufacturing unit. For that, the diseased animals should not be used for milking and the equipment for kefir making should be cleaned and sanitised. Milk must be pasteurised or boiled before the production, and the storage of milk should be realised under refrigeration.

With respect to the food safety and hygiene management, pathogen reduction systems like HACCP can be adopted to manufacturing plants to assure microbiological safety.

References

- ACKERS M., MAHON B.E., LEAHY E., GOODE B., DAMROW T., HAYES P.S., BIBB W.F., RICE D.H., BARET T.J., HUTWAGNER L., GRIFFIN P.M., SLUTSKER L. (1998): An outbreak of *Escherichia coli* O157:H7 infections associated with leaf lettuce consumption. *The Journal of Infectious Diseases*, **177**: 1588–1593.
- AOAC (Association of Official Analytical Chemists) (2001): *Bacteriological Analytical Manual*. 5th Ed. AOAC, Washington.

- BACHROURI M., QUINTO E.J., MORA M.T. (2006): Kinetic parameters of *Escherichia coli* O157:H7 survival during fermentation of milk and refrigeration of home-made yoghurt. *International Dairy Journal*, **16**: 474–481.
- BAIRD-PARKER A.C. (1962): An improved diagnostic and selective medium for isolating coagulase positive staphylococci. *Journal of Applied Bacteriology*, **25**: 12–19.
- BUCHANAN R.L., DOYLE M.P. (1997): Foodborne disease significance of *Escherichia coli* O157:H7 and other enterohemorrhagic *Escherichia coli*. *Food Technology*, **51**: 61–76.
- CHATEAU N., CASTELLANOS I., DESCHAMPS A.M. (1993): Distribution of pathogen inhibition in the *Lactobacillus* isolates of a commercial probiotic consortium. *Journal of Applied Bacteriology*, **74**: 36–40.
- CHENG H.Y., CHOU C.C. (1999): Survival of acid-adapted *Escherichia coli* O157:H7 in some acid foods and subsequent environmental stress. [MS Thesis.] National Taiwan University, Taipei.
- CONNOR D.E., KOTROLA J.S. (1995): Growth and survival of *Escherichia coli* O157:H7 under acidic conditions. *Applied and Environmental Microbiology*, **61**: 382–385.
- DIENEEN S.S., TAKEUCHI K., SOUDAH J.H., BOOR K.J. (1998): Persistence of *Escherichia coli* O157:H7 in Dairy Fermentation Systems. *Journal of Food Protection*, **61**: 1602–1608.
- ERIBO B., ASHENAFI M. (2003): Behavior of *Escherichia coli* O157:H7 in tomato and processed tomato products. *Food Research International*, **36**: 823–830.
- FOSTER J.W., HALL H.K. (1990): Adaptive acidification tolerance response of *Salmonella typhimurium*. *Journal of Bacteriology*, **172**: 771–778.
- GLASS K.A., LOEFTEHOLZ J.M., FORD J.P., DOYLE M.P. (1992): Fate of *Escherichia coli* O157:H7 as affected by pH or sodium chloride and in fermented dry sausage. *Applied and Environmental Microbiology*, **58**: 2513–2516.
- GOVARIS A., KOIDIS P., PAPTAEODOROU K. (2001): The fate of *Escherichia coli* O157:H7 in Myzithra, Anthotyros and Manouri whey cheeses during storage at 2 and 12°C. *Food Microbiology*, **18**: 565–570.
- GOVERD K.A., BEECH F.W., HOBBS R.P., SHANNON R. (1979): The occurrence and survival of coliforms and salmonellas in apple juice and cider. *Journal of Applied Bacteriology*, **46**: 521–530.
- GRIFFIN P.M., TAUXE R.V. (1991): The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E. coli* and the associated hemolytic uremic syndrome. *Epidemiology Reviews*, **13**: 60–98.
- GÜLMEZ M., GÜVEN A. (2003): Survival of *E. coli* O157:H7, *Listeria monocytogenes* 4b and *Yersinia enterocolitica* O3 in different yogurt and kefir combinations as prefermentation containment. *Journal of Applied Microbiology*, **95**: 631–636.
- GURAYA R., FRANK J.F., HASSAN A.N. (1998): Effectiveness of salt, pH, and diacetyl as inhibitors of *Escherichia coli* O157:H7 in dairy foods stored at refrigerator temperatures. *Journal of Food Protection*, **61**: 1092–1098.
- GÜZEL-SEYDİM Z., SEYDİM A.C., GREENE A.K. (2000): Organic acid and volatile flavor components evolved during refrigerated storage of kefir. *Journal of Dairy Science*, **83**: 275–277.
- HARIS L.J., DAESCHEL M., STILES M.E., KLAENHAMMER T.R. (1989): Antimicrobial activity of lactic acid bacteria against *Listeria monocytogenes*. *Journal of Food Protection*, **52**: 348–387.
- HSIN-YI C., CHOU C.C. (2001): Acid adaptation and temperature effect on the survival of *E. coli* O157:H7 in acidic fruit juice and lactic fermented milk product. *International Journal of Food Microbiology*, **70**: 189–195.
- IURLINA M.O., FRITZ R. (2004): Microbiological quality of Port Salut Argentino cheese stored at two temperature treatments. *Lebensmittel-Wissenschaft und Technologie*, **37**: 739–748.
- KOODIE L., DHOPE A.M. (2001): Acid tolerance of *Escherichia coli* O157:H7 and its survival in apple juice. *Microbios*, **104**: 167–175.
- LARA J.A.F., SENIGALIA S.W.B., OLIVEIRA T.C.R., DUTRA I.S., PINTO F., SHIMOKOMAKI M. (2003): Evaluation of survival of *Staphylococcus aureus* and *Clostridium botulinum* in charqui meats. *Meat Science*, **65**: 609–613.
- LEKKAS C., KAKOURI A., PALEOLOGOS E., VOUTSINAS L.P., KONTOMINAS M.G., SAMELIS J. (2006): Survival of *Escherichia coli* O157:H7 in Galotyri cheese stored at 4 and 12°C. *Food Microbiology*, **23**: 268–276.
- MARTIN J.H., MARSHALL D.L. (1995): Characteristics and control of potential foodborne pathogens in cultured dairy foods. *Journal of the Mississippi Agricultural and Forestry Experiment Station*, **30**(3): 9–16.
- MAREK P., NAIR M.K.M., HOAGLAND T., VENKITA-RAYANAN K. (2004): Survival and growth characteristics of *Escherichia coli* O157:H7 in pasteurized and unpasteurized Cheddar cheese whey. *International Journal of Food Microbiology*, **94**: 1–7.
- MARSHALL V.M., COLE W.M. (1985): Methods for making kefir and fermented milks based on kefir. *Journal of Dairy Research*, **52**: 451–456.
- MCCLURE P. (2000): The impact of *E. coli* O157:H7 on the food industry. *World Journal of Microbiology and Biotechnology*, **16**: 749–755.

- MCDONOUGH S., HEER F., SHIRLEY L. (1991): Foodborne outbreak of gastroenteritis caused by *Escherichia coli* O157:H7 – North Dakota, 1990. Morbidity and Mortality Weekly Report, **40**: 265–267.
- MCINGVALE S.C., CHEEN X.Q., MCKILLIP J.L., DRAKE M.A. (2000): Survival of *Escherichia coli* O157:H7 in buttermilk as affected by contamination point and storage temperature. Journal of Food Protection, **63**: 441–444.
- MILLER L.G., KASPAR C.W. (1994): *E. coli* O157:H7 acid tolerance and survival in apple cider. Journal of Food Protection, **57**: 460–464.
- MOON K.D., DELAQUIS P., TOIVONEN P., STANICH K. (2006): Effect of vanillin on the fate of *Listeria monocytogenes* and *Escherichia coli* O157:H7 in a model apple juice medium and in apple juice. Food Microbiology, **23**: 169–174.
- OGWARO B.A., GIBSON H., WHITEHEAD M., HILL D.J. (2002): Survival of *Escherichia coli* O157:H7 in traditional African yoghurt fermentation. International Journal of Food Microbiology, **79**: 105–112.
- PORTOCARRERO S., NEWMAN M., MIKEL B. (2002): *Staphylococcus aureus* survival, staphylococcal enterotoxin production and shelf stability of country-cured hams manufactured under different processing procedures. Meat Science, **62**: 267–273.
- REINDERS R.D., BIIESTERVELD S., BIJKER P.G. (2001): Survival of *Escherichia coli* O157:H7 ATCC 43895 in a model apple juice medium with different concentrations of proline and caffeic acid. Applied and Environmental Microbiology, **67**: 2863–2866.
- REITISMA C.J., HENNING D.R. (1996): Survival of enterohemorrhagic *Escherichia coli* O157:H7 during the manufacture and curing of Cheddar cheese. Journal of Food Protection, **59**: 460–464.
- ROERING A.M., LUCHANSKY J.B., IHNOT A.M., ANSAY S.E., KASPAR C.W., INGHAM S.C. (1999): Comparative survival of *Salmonella typhimurium* DT 104, *Listeria monocytogenes*, and *Escherichia coli* O157:H7 in preservative-free apple cider and simulated gastric fluid. International Journal of Food Microbiology, **46**: 263–269.
- ROZAND C.V., CRUCHAUDET C.M., BAVAI C., MONTET M.P., BONIN V., DERNBURG A., RICHARD Y. (2005): Growth and survival of *Escherichia coli* O157:H7 during the manufacture and ripening of raw goat milk lactic cheeses. International Journal of Food Microbiology, **105**: 83–88.
- SAMELIS J., SOFOS J.N. (2003): Organic acids. In: ROLLER S. (ed.): Naturel Antimicrobials for the Minimal Processing of Foods. CRC Woodhead Publishing Ltd, Cambridge: 98–132.
- SAS (1999): SAS Ins. Inc., Cary, NC, USA. Proprietary Software Release 8.2 (TS2MO).
- ŞİMSEK B., SAĞDIÇ O., ÖZÇELİK S. (2007): Survival of *Escherichia coli* O157:H7 during the storage of ayran produced with different spices. Journal of Food Engineering, **78**: 676–680.
- TSEGAYE M., ASHENAFI M. (2005): Fate of *Escherichia coli* O157:H7 during the processing and storage of Ergo and Ayib, traditional Ethiopian dairy products. International Journal of Food Microbiology, **103**: 11–21.
- TSEGAYE M., EPRAHIM E., ASHENAFI M. (2004): Behaviour of *Escherichia coli* O157:H7 during fermentation of Datta and Awaze, traditional Ethiopian fermented condiments, and during product storage at ambient and refrigeration temperatures. Food Microbiology, **21**: 6, 223–232.
- TSIAI Y.W., INGHAM S.C. (1997): Survival of *Escherichia coli* O157:H7 and *Salmonella* spp. in acidic condiments. Journal of Food Protection, **60**: 751–755.
- ULJAS H.E., INGHAM S.C. (1999): Combinations of intervention treatments resulting in 5-log 10-unit reductions in numbers of *Escherichia coli* O157:H7 and *Salmonella typhimurium* DT104 organisms in apple cider. Applied and Environmental Microbiology, **65**: 1924–1929.
- UYTENDAELE M., TAVERNIERS I., DEBEVERE J. (2001): Effect of stress induced by suboptimal growth factors on survival of *Escherichia coli* O157:H7. International Journal of Food Microbiology, **66**: 31–37.
- WSZOLEK M., TAMIME A.Y., MUIR D.D., BARCLAY M.N.L. (2001): Properties of kefir made in Scotland and Poland using bovine, caprine and ovine milk with different starter cultures. Lebensmittel-Wissenschaft und Technologie, **34**: 251–261.
- ZHAO T., DOYLE M.P., BESSAR R.E. (1993): Fate of enterohemorrhagic *Escherichia coli* O157:H7 in apple cider with and without preservatives. Applied and Environmental Microbiology, **59**: 2526–2530.
- ZUBILLAGA M., WEILL R., POSTAIRE E., GOLDMAN C., CARO R., BOCCIO J. (2001): Effect of probiotics and functional foods and their use in different diseases. Nutrition Research, **21**: 569–579.

Received for publication January 10, 2007

Accepted after corrections March 9, 2007

Corresponding author:

Dr. BÜLENT ERGÖNÜL, Celal Bayar University, Engineering Faculty, Food Engineering Department, Muradiye, 45040 Manisa, Turkey
tel.: + 90 236 241 21 44, fax: + 90 236 241 21 43, e-mail: bulent.ergonul@bayar.edu.tr
