

The effect of weaning age on performance and nutrient digestibility of broiler rabbits

L. ZITA¹, E. TŮMOVÁ¹, V. SKŘIVANOVÁ², Z. LEDVINKA¹

¹Department of Animal Husbandry, Czech University of Life Sciences in Prague, Czech Republic

²Institute of Animal Science, Prague-Uhřetěves, Czech Republic

ABSTRACT: The effect of weaning age on growth, feed consumption, digestibility of nutrients, carcass yield, mortality and parasite incidence was studied in a balance experiment. Forty Hyplus[®] rabbits proceeding from a commercial rabbitry were assigned to four groups of 10 rabbits according to weaning age at 25, 28, 31 and 35 days. The experimental diet had 17.03% crude protein, 4.66% fat and 18.30% crude fibre. Feed and water were available *ad libitum*. At the end of the experiment, at 84 days of age, six rabbits per group were slaughtered. In the experiment, most results of performance were not significantly ($P \leq 0.05$) affected by weaning age. Only the rabbits weaned at 25 days of age had higher live weight ($P \leq 0.05$) at the age of 35 days (996, 986, 971 and 910 g, respectively), but at the end of the experiment we did not find any differences between groups. Higher digestibility of nutrients was recorded in the second collection period. In the first collection period the digestibility of nutrients was higher in rabbits weaned at 25 days of age in comparison with the other groups. No differences between groups in the digestibility of nutrients were observed in the second period. The age of weaning did not influence carcass characteristics. Dressing percentage was insignificantly higher in rabbits weaned at 25 and 35 days of age (53.11 and 53.07%, respectively) in comparison with rabbits weaned at 28 and 31 days of age (50.89 and 52.67%, respectively). There was no significant effect of weaning age on mortality. The incidence of *Eimeria* ssp. was the highest after weaning in rabbits at the age of 35 days.

Keywords: rabbit; early weaning; growth; carcass characteristics; nutrient digestibility; parasite incidence; mortality

Early weaning is one of the solutions allowing the rabbits to be offered a diet adapted to their needs from a very early age. Kits begin to consume limited amounts of solid feed at 18 to 20 days of age (Maertens and de Groote, 1990; Scapinello et al., 1999), then the microflora begins caecal colonization and fermentation develops soon afterwards (Gidenne, 1997), and some of the enzymatic digestive activities show important changes (Marounek et al., 1995). Early weaning reduces the incidence of digestive disorders and reduces pathogen transmission by limiting contacts between litters and does (Schlölaut, 1988).

The growth of rabbits could be affected by weaning age. Trocino et al. (2001) observed that early

weaned kits showed a lower live weight in comparison with rabbits weaned at 32 days of age. Similar results were reported by Gidenne and Fortun-Lamothe (2001). The negative effect of early weaning on live weight was also found by Ferguson et al. (1997), Gidenne and Fortun-Lamothe (2001, 2003, 2004) and Gallois et al. (2003, 2004). In contrast Petersen et al. (1992), Gidenne et al. (2004) and Tůmová et al. (2006a) did not prove any significant differences in live weight during the time of fattening.

Feed consumption influences the development of digestive tract. According to Gallois et al. (2003, 2004) slight solid feed consumption led to the earlier development of digestive tract. On the other

Supported by the Ministry of Education, Youth and Sports of the Czech Republic (Project No. MSM 6046070901) and by the Ministry of Agriculture of the Czech Republic (Project No. MZe 0002701403).

hand, the digestive tract development does not affect feed consumption of early or traditionally weaned rabbits (Xiccato et al., 2000; Tůmová et al., 2006a).

There is a lack of information on the digestibility of nutrients in early weaned rabbits. Nicodemus et al. (2004) and Tůmová et al. (2006a) found that the dry matter and crude protein digestibility was higher in early weaned rabbits and decreased with age (Tůmová et al., 2002, 2006a,b).

Mortality can also be affected by the age of weaning. McNitt and Moody (1992), Ferguson et al. (1997), Gidenne and Fortun-Lamothe (2001, 2004) reported higher mortality in early weaned rabbits. According to Piattoni et al. (1999), Trocino et al. (2001) and Xiccato et al. (2003) mortality was not affected by the age of weaning.

Little is known about the carcass performance of early weaned rabbits. Trocino et al. (2001) indicated the lower empty body weight in early weaned rabbits at 21 days of age, but at 56 days of age the empty body weight was similar in weaning groups. In our previous experiments (Tůmová et al., 2006c) the age of weaning did not affect dressing percentage.

The aim of the present study was to evaluate the effect of weaning age on growth, feed consumption,

digestibility of nutrients, carcass yield, mortality and parasite incidence in broiler rabbits.

MATERIAL AND METHODS

Animals, diet and design

In a balance experiment forty Hyplus® rabbits (♂ PS59 × ♀ PS19) were used. The rabbits were from a commercial rabbitry and were taken from multiparous does at the weaning age of 25, 28, 31 and 35 days and were divided into 4 groups of 10. The rabbits were housed in all-wire individual metabolic cages with urine and faeces separation. The floor density was 0.2 m² per rabbit. The temperature of 16°C and relative humidity about 65% were maintained during the whole period of fattening. A twelve-hour photoperiod was used. Feed and water were available *ad libitum*. During the experiment rabbits received a commercial pelleted feed mixture until 84 days of age. The diet composition is given in Table 1 and the analysed content of nutrients per 1 kg of diet is in Table 2. Before weaning, kits received a feed mixture for does without anticoncidual drugs.

Animals were individually weighed every week, feed consumption was measured daily. Mortality and morbidity were also recorded in groups over the whole period of fattening.

At the end of the experiment, at 84 days of age, six rabbits per group of the average live weight were slaughtered. The slaughter and carcass dissection

Table 1. Ingredients of experimental diet (g/kg)

Ingredients	g/kg
Soybean meal	30
Sunflower meal	170
Barley	80
Oats	90
Lucerne meal	300
Wheat bran	225
Sugar-beet pulp	60
Rapeseed oil	15
Vitamin-mineral supplement*	10
Ground limestone	10
Dicalcium phosphate	5
Feeding salt	5

*Composition of vitamin and mineral supplement in 1 kg: vitamin A 1 200 000 IU; vitamin D₃ 200 000 IU; alpha tocopherol 5 000 mg; vitamin K₃ 200 mg; vitamin B₁ 300 mg; vitamin B₂ 700 mg; vitamin B₆ 400 mg; vitamin B₁₂ 2 mg; niacin amide 5 000 mg; calcium pantothenate 2 000 mg; biotin 20 mg; folic acid 170 mg; choline 60 000 mg; DL-methionine 100 g; L-lysine HCL 20 g; sodium salinomycin 2 250 mg; cobalt 100 mg; copper 2 000 mg; iron 5 000 mg; iodine 120 mg; manganese 4 700 mg; zinc 5 000 mg; selenium mg; antioxidant 10 000 mg

Table 2. Analysed nutrient content of experimental diet (g/kg)

Composition	g/kg
Dry matter	905.0
Crude protein	170.3
Fat	46.6
Crude fibre	183.0
Ash	74.3
Nitrogen-free extract	430.8
Starch	152.2
ADF	223.6
NDF	354.8
Nutritive values:	
Digestible protein	120.0
Digestible energy (MJ)	9.9
Digestible protein/digestible energy (g/MJ)	12.1

Table 3. Performance traits and mortality of rabbits

Measurement	Group (days of age)				Signifi- cance	SEM
	25	28	31	35		
Live weight (g)						
at weaning (initial)	564	631	746	910		
at 35 days of age	996 ^a	986 ^{ab}	971 ^{ab}	910 ^b	*	14.177
at 84 days of age	3 210	3 246	3 226	3 238	NS	42.659
Average daily weight gain (g)						
35–84 days of age	45.59	46.14	46.02	47.55	NS	0.759
Feed consumption (35–84 days of age)						
per day/rabbit (g)	150.8	177.5	174.8	140.9	NS	6.704
per kg live weight (kg)	3.83	3.87	3.81	3.69	NS	0.041
Mortality (<i>n</i>)	2	0	0	3		

* $P \leq 0.05$; NS = not significant; SEM = standard error mean

^{a,b}means with common superscripts do not differ significantly, determined by Scheffe's test

were carried out in an experimental slaughterhouse. Rabbits were fasted overnight, and slaughtered the following morning by electric stunning and bleeding by jugular cut. The method of carcass measurements was harmonised with Blasco and Ouhayoun (1996).

The digestibility of nutrients was determined by a quantitative method, according to the modified method of Perez et al. (1995). There were two collection periods: the first from 42 to 49 days of age and the second from 56 to 63 days of age. Each collection period lasted for 7 days. During the collection period, the total faeces excretion was collected daily into plastic bags and stored at -18°C until analysis. Faeces were analysed for dry matter, crude protein, fat, crude fibre, ash and nitrogen-free extract.

Parasite incidence was determined a day after weaning (e.g. at 26, 29, 32 and 36 days of age) and at the age of 42, 49, 56 and 63 days.

Analytical methods

The feed and faeces were analysed using standard AOAC methods (1995). Protein and fat concentrations were determined employing the Kjeltex Auto A 1030 Analyser and Soxtec 1043 from Tecator AB (Sweden), respectively. Crude fibre was determined according to the procedure of Van Soest (1991), using Fibertec 2010.

Coefficients of digestibility of nutrients were determined according to the formula:

$$\frac{\{(\text{quantity of feed} \times \text{nutrients of feed}) - (\text{quantity of faeces} \times \text{nutrients of faeces})\}}{\text{quantity of feed} \times \text{nutrients of feed}} \times 100$$

Samples of excrements were examined for the presence of parasites according to the procedures of Bréza (1957) and Pavlásek (1991).

Statistical analyses

The statistical analysis of data on fattening characteristics and carcass yield was performed by one-way analysis of variance by the GLM procedure of SAS (SAS, 2003). The significance of differences between groups was tested by Scheffe's test on the level of significance $P \leq 0.05$. Data on digestibility were evaluated by two-way analysis of variance, treatment and balance interactions using the GLM procedure of SAS (SAS, 2003). Means marked with a different superscript letter within each column are significantly different.

RESULTS AND DISCUSSION

In the experiment, most results of performance (Table 3) were not significantly ($P \leq 0.05$) affected by weaning age. Only the rabbits weaned at 25 days of age had higher live weight ($P \leq 0.05$) at the age of 35 days, but at the end of the experiment we did not find any differences between groups. The results of our experiment are in accordance with findings of Xiccato et al. (2000, 2003), Gidenne and Fortun-Lamothe (2001), Pascual et al. (2001), Trocino et al. (2001), Gidenne et al. (2004) and Tůmová et al.

Table 4. Digestibility of nutrients (%)

Nutrient	Weaning age (days)	Balance (days of age)		Significance			SEM
		42–49	56–63	balance	group	balance × group	
Dry matter	25	58.94	59.79	< 0.0001	0.0085	0.0034	0.818
	28	58.31	60.91				
	31	49.21	60.81				
	35	49.20	60.67				
Crude protein	25	73.57	73.80	0.0171	0.1021	0.5419	0.509
	28	71.79	74.29				
	31	68.52	72.73				
	35	70.18	72.77				
Crude fibre	25	23.00	17.90	0.6710	0.7123	0.0837	0.934
	28	21.31	19.97				
	31	14.39	22.44				
	35	20.64	22.24				
Fat	25	88.89	90.15	< 0.0001	0.1836	0.0070	1.393
	28	88.32	91.66				
	31	75.61	94.14				
	35	74.36	94.22				
Ash	25	49.31	47.40	0.1435	0.8830	0.3488	0.894
	28	46.87	48.07				
	31	41.22	45.33				
	35	40.20	47.08				
Nitrogen-free extract	25	66.85	70.90	< 0.0001	0.0003	0.0006	0.894
	28	67.41	71.89				
	31	56.42	71.46				
	35	56.94	70.94				

SEM = standard error mean

(2006a). On the other hand, lower live weight of early weaned rabbits was recorded at 35 days of age by Xiccato et al. (2000), Gidenne and Fortun-Lamothe (2001), Trocino et al. (2001).

Feed consumption was insignificantly higher in rabbit weaned at 28 days of age (Table 3). The results correspond with Gallois et al. (2003, 2004), who demonstrated that the early weaned rabbits had the higher feed consumption in comparison with rabbits weaned at 35 days of age.

The results of digestibility of nutrients are presented in Table 4. The digestibility of dry matter and nitrogen-free extract was significantly affected by weaning age. In the 1st collection period the digestibility of dry matter and nitrogen-free extract was higher in rabbits weaned at 25 and 28 days of age compared with those weaned at 31 and 35 days of age. The higher digestibility of dry matter in early

weaned rabbits was also reported by Nicodemus et al. (2004) and Tůmová et al. (2006a). A significant difference was determined between collection periods in digestibility of dry matter, crude protein, fat and nitrogen-free extract whereas higher digestibility of nutrients was in the second collection period. This result was obtained also by Tůmová et al. (2002, 2006a,b), who reported that nutrient digestibility decreased with the age in all rabbits. There were significant interactions among groups and age in digestibility of dry matter, fat and nitrogen-free extract.

Carcass characteristics were not influenced by weaning age (Table 5). Dressing percentage was insignificantly higher in rabbits weaned at 25 and 35 days of age in comparison with rabbits weaned at 28 and 31 days of age. Tůmová et al. (2006c) also reported that the rabbits of different weaning

Table 5. Carcass analysis at 84 days of age

Characteristic	Group (days of age)				Significance	SEM
	25	28	31	35		
Live weight (g)	3 184.00	3 153.00	3 091.00	3 167.00	NS	33.760
Carcass weight (g)	1 689.00	1 657.00	1 626.17	1 676.83	NS	19.270
Dressing percentage (%)	53.11	50.89	52.67	53.07	NS	0.708
Fore part of the carcass (%)	51.72	51.59	51.77	51.01	NS	0.303
Hind part of the carcass (%)	48.27	45.20	48.28	49.01	NS	0.905
Loin (%)	16.90 ^b	16.76 ^b	17.02 ^{ab}	18.28 ^a	*	0.240
Hind legs (%)	31.39	31.67	31.27	30.73	NS	0.216
Thigh muscles (%)	24.17	24.15	23.18	23.23	NS	0.213
Kidney (%)	1.24	1.23	1.28	1.31	NS	0.020
Renal fat (%)	2.25	2.06	1.95	2.25	NS	0.139
Heart (%)	0.64 ^a	0.52 ^b	0.50 ^b	0.52 ^b	*	0.022
Liver (%)	5.86	5.97	5.89	6.21	NS	0.287
Lung (%)	0.74 ^b	0.72 ^b	0.98 ^a	0.77 ^{ab}	**	0.034
Skin from live weight (%)	15.60	15.95	16.80	15.77	NS	0.260

* $P \leq 0.05$, ** $P \leq 0.01$; NS = not significant; SEM = standard error mean

^{a,b} means with common superscripts do not differ significantly, determined by Scheffe's test

age had similar dressing percentage at the end of the experiment. Differences between groups in the proportion of single parts from the carcass were not significant except the loin. The proportion of loin from carcass decreased with weaning age. An insignificantly lower proportion of renal fat was in the group weaned at 31 days of age. The age of weaning had a significant ($P \leq 0.05$, 0.01) effect on the proportion of heart and lungs from the carcass. A higher proportion of heart was in rabbits weaned

at 25 days of age while in rabbits weaned at 31 days of age a higher proportion of lungs from the carcass was recorded.

There was no significant effect of weaning age on mortality (Table 3). This result coincides with findings of Piattoni et al. (1999), Trocino et al. (2001) and Xiccato et al. (2003), who did not report the effect of weaning age on mortality either.

During the experiment, oocysts of *Passalurus ambiguus*, *Eimeria magna*, *Eimeria perforans*, *Eimeria*

Table 6. Parasite incidence per rabbit (OPG – oocysts per gram of faeces)

Age at sampling (days of age)	Parasite	Group (days of age)			
		25	28	31	35
26	<i>Eimeria magna</i>	184.7	–	–	–
	<i>Eimeria perforans</i>	396	x	–	–
29	<i>Passalurus ambiguus</i>	xx	6.6	–	–
32		xx	xx	N	–
	<i>Eimeria exigua</i>	xx	xx	xx	176
36	<i>Eimeria media</i>	xx	xx	xx	58 088.9
	<i>Eimeria perforans</i>	xx	xx	xx	75 515.6
42	<i>Eimeria magna</i>	N	N	N	28.3
49		N	N	N	N
56	<i>Passalurus ambiguus</i>	346.5	488.4	N	N
63	<i>Passalurus ambiguus</i>	N	6.6	N	N

xx = parasite incidence not determined; N = negative

exigua and *Eimeria media* were found (Table 6). The incidence of *Eimeria* spp. was highest in rabbits weaned at the age of 35 days.

In the present study, the age of weaning did not have a significant effect on the majority of the parameters of performance. Rabbits weaned at 25 days of age had significantly higher live weight at the age of 35 days and then the live weight was not significantly affected till the end of fattening. It is possible to draw a conclusion that weaning age and collection period affected digestibility of dry matter and fat and that the digestibility of nutrients decreased with age. The age of weaning did not influence carcass characteristics (except the proportion of loin, heart and lungs from carcass) and mortality. The highest incidence of *Eimeria* spp. was in rabbits at the age of 35 days, probably being caused by a longer time of coexistence with does.

REFERENCES

- AOAC (1995): Official Methods of Analysis. Association of Official Analytical Chemists, 16th ed., Arlington, Virginia, USA.
- Bréza M. (1957): Several practical information and subjects about parasites diagnosis. *Helminológia*, 1, 57–63.
- Blasco A., Ouhayoun J. (1996): Harmonization of criteria and terminology in rabbit meat research. Revised proposal. *World Rabbit Sci.*, 4, 93–99.
- Ferguson F.A., Lukefahr S.D., McNitt J.I. (1997): A technical note on artificial milk feeding of rabbit kits weaned at 14 days. *World Rabbit Science*, 5, 65–70.
- Gallois M., Gidenne T., Fortun-Lamothe L. (2003): Digestive development in the young rabbit: impact of a weaning at 21 d. In: Proc. 3rd and 4th COST Action 848, Prague, Czech Republic, 15 pp.
- Gallois M., Gidenne T., Fortun-Lamothe L., Le Huerou-Luron I., Lalles J.P. (2004): Weaning age and development of the small intestinal mucosa in the young rabbit. In: Proc. 8th World Rabbit Congress, Puebla, Mexico, 1079–1085.
- Gidenne T. (1997): Caeco-colic digestion in the growing rabbit: impact of nutritional factors and related disturbance. *Livest. Prod. Sci.*, 51, 73–88.
- Gidenne T., Fortun-Lamothe L. (2001): Early weaning: effect on performances and health. In: Proc. 2nd Meetings of Workgroup 3rd and 4th COST Action 848, Gödöllő, Hungary, 34 pp.
- Gidenne T., Fortun-Lamothe L. (2003): Microbial activity in the early weaned rabbit. In: Proc. 3rd Meetings of Workgroup 3rd and 4th COST Action 848, Prague, Czech Republic, 14 pp.
- Gidenne T., Fortun-Lamothe L. (2004): Growth, health status and digestion of rabbits weaned at 23 or 32 days of age. In: Proc. 8th World Rabbit Congress, Puebla, Mexico, 846–852.
- Gidenne T., Lapanouse A., Fortun-Lamothe L. (2004): Feeding strategy for the early weaned rabbit: interest of a high energy and protein starter diet on growth and health status. In: Proc. 8th World Rabbit Congress, Puebla, Mexico, 853–860.
- Maertens L., de Groote G. (1990): Feed intake of rabbit kits before weaning and attempts to increase it. *J. Appl. Rabbit Res.*, 13, 151–158.
- McNitt J.I., Moody G.L. (1992): A method for weaning rabbit kits at 14 days. *J. Appl. Rabbit Res.*, 15, 661–665.
- Marounek M., Vovk S.J., Skřivanová V. (1995): Distribution of activity of hydrolytic enzymes in the digestive tract of rabbits. *Brit. J. Nutr.*, 73, 463–469.
- Nicodemus N., Pérez-Alba L., Carabano R., de Blas C., Badiola I., Pérez de Rozas A., García J. (2004): Effect of level of fibre and level of ground of fibre sources on digestion and ileal and caecal characterization of microbiota of early weaned rabbits. In: Proc. 8th World Rabbit Congress, Puebla, Mexico, 928–929.
- Pascual J.J., Cervera C., Fernández-Carmona J. (2001): *In vivo* evolution of the corporal condition on primiparous rabbit does. In: Proc. 2nd Meeting of Workgroup 3rd and 4th COST Action 848, Gödöllő, Hungary, 49 pp.
- Pavlašek I. (1991): Use of glycerine in the detection of the oocysts of *Cryptosporidium parvum* and *C. baileyi* in the excrements of mammals and birds. *Vet. Med.*, 36, 255–256.
- Perez J.M., Lebas F., Gidenne T., Maertens L., Xiccato G., ParigiBini R., Dalle Z.A., Cossu M.E., Carazzolo A., Villamide M.J., Carabano R., Fraga M.J., Ramos M.A., Cervera C., Blas E., Fernandez J., Falcao e Cunha L., Bengala Freire J. (1995): European reference method for *in vivo* determination of diet digestibility in rabbits. *World Rabbit Sci.*, 3, 41–43.
- Petersen J., Klausdeinken F.J., Gerken M. (1992): Influence of weaning age on development of live-weight and food consumption in young rabbits. *J. Appl. Rabbit Res.*, 15, 856–863.
- Piattoni F., Maertens L., Mazzoni D. (1999): Effect of weaning age and solid feed distribution before weaning on performances and caecal traits of young rabbits. In: Proc. 2nd Internat. Conference on Rabbit Production in Hot Climates. Adana 7–9 September, 1998. *Cah. Options Méditerran.*, 41, 85–92.

- SAS (2003): SAS Institute Inc., the SAS System for Windows. Release 9.1.3
- Scapinello C., Gidenne T., Fortun-Lamothe L. (1999): Digestive capacity of the rabbit during the post-weaning period according to the milk/solid feed intake pattern before weaning. *Reprod. Nutr. Dev.*, 39, 423–432.
- Schlolaut W. (1988): Present husbandry and management conditions and development trends in rabbit production. In: *Proc. 4th World Rabbit Congress*, Budapest, Hungary, 1, 93–112.
- Trocino A., Xiccato G., Sartori A., Queaque P.I. (2001): Effect of starter diet and weaning age on growth, caecal fermentation and body composition of young rabbits. In: *Proc. 2nd Meetings of Workgroup 3rd and 4th COST Action 848*, Gödöllő, Hungary, 52–53.
- Tůmová E., Skřivanová V., Skřivan M. (2002): Effect of restricted feeding time and quantitative restriction in growing rabbits. *Arch. Geflügelkd.*, 66, 1–9.
- Tůmová E., Zita L., Marounek M., Skřivanová V., Berladyn A. (2006a): The effect of weaning age on performance, nutrient digestibility and lipase activity in broiler rabbits. In: *Proc. 3rd Rabbit Congress of the Americas*, Maringá, Brasil, CD.
- Tůmová E., Zita L., Skřivanová V., Fučíková A., Skřivan M., Burešová M. (2006b): Digestibility of nutrients, organ development and blood picture in restricted and ad libitum fed broiler rabbits. *Arch. Geflügelkd.*, 71, 6–12.
- Tůmová E., Zita L., Štolc L. (2006c): Carcass quality in restricted and *ad libitum* fed rabbits. *Czech J. Anim. Sci.*, 51, 214–219.
- Van Soest P.J., Robertson J.B., Lewis B.A. (1991): Methods for dietary fibre, neutral detergent fibre, and non starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.*, 74, 3583–3597.
- Xiccato G., Trocino A., Sartori A., Queaque P.I. (2000): Early weaning of rabbits: effect of age and diet on weaning and post-weaning performance. In: *Proc. 7th World Rabbit Congress*, Valencia, Spain, C, 483–490.
- Xiccato G., Trocino A., Sartori A., Queaque P.I. (2003): Effect of weaning diet and weaning age on growth, body composition and caecal fermentation of young rabbits. *Anim. Sci.*, 77, 101–111.

Received: 2006–11–02

Accepted after corrections: 2007–07–02

Corresponding Author

Ing. Lukáš Zita, Ph.D., Department of Animal Husbandry, Czech University of Life Sciences Prague, Kamýcká 129, 165 21 Prague 6-Suchbát, Czech Republic
Tel. +420 224 383 053, e-mail: zita@af.czu.cz
