

Changes in fungal communities in organically fertilized soil

B. Cwalina-Ambroziak¹, T. Bowszys²

¹*Department Phytopathology and Entomology, University of Warmia and Mazury, Olsztyn, Poland*

²*Department of Agricultural Chemistry and Environmental Protection, University of Warmia and Mazury, Olsztyn, Poland*

ABSTRACT

The research project was carried out as a field experiment with application of the following crop rotation system: industrial potato, spring barley for fodder, winter rape and winter wheat, established in the random distribution of blocks in triplicate on gleyic luvisol formed of silty light loam. The aim of the research was to determine the influence of diversified organic fertilization based on composted wastewater sediments and farm manure on the community of soil fungi as compared to fields without fertilization and with NPK fertilization only. The fungi were cultured on the Martin medium and were counted and identified afterwards. As a result of three-year field experiments it was established that organic fertilization had a more determining effect on qualitative composition than numbers of soil fungi. Total number of fungal colony-forming units in the soil fertilized with Biohum at 10 t/ha and 5 t/ha was significantly higher than in soil with mineral NPK fertilization and without fertilization. Most frequently pathogens populated the soil in fields without fertilization and to a lesser extent the soil with mineral NPK fertilization. A positive influence of organic fertilizers on the fungal community structure was recorded. The number of pathogens was limited (to 1.2% in fields fertilized with farm manure) while the population of saprotrophic fungi possessing antagonistic properties increased.

Keywords: soil; organic fertilization; pathogenic fungi; saprotrophic fungi; water extract from compost

Composted wastewater sediments formed as a result of biological treatments can be used for fertilization of soil enriching it with organic matter and at the same time improving its structure. These fertilizers, due to their content of micro- and macroelements, represent a source of nutrients for plants. The increase of the microbiological diversity and activity of soil as a consequence of organic fertilization is widely documented in literature (Shannon et al. 2002, Larkin et al. 2006). It contributes to limiting some soil fungi development, including pathogens from genera *Pythium*, *Phytophthora* and *Fusarium* (Hoitink and Boehm 1999). At the same time the numbers of positive microflora possessing antagonistic influence in relation to these pathogens increase, which is then a very positive phenomenon. Soil micro-

organisms, including fungi, represent biological protection of plants against pathogenic factors (Mathur et al. 2006). The above changes in the structure of soil fungi community contribute to improving the health of cultivated crops. Also, the excretions from roots of the cultivated crops can influence the qualitative composition of the soil fungi community (Angus et al. 1994, Grayston et al. 1998).

MATERIAL AND METHODS

The experiment was established by the Chair of Agricultural Chemistry and Environment Protection in 2004 and is located at the Production-Experimental Enterprise in Bałcyny on gleyic lu-

This article was presented at the conference "Practical Solutions for Managing Optimum C and N Content in Agricultural Soils IV", 19–23 June 2007, Prague, Czech Republic.

visol formed of silty light loam (Polish Standard BN-78/9180-11). Prior to the experiment the soil content of P (38.2 mg/kg), K (105.8 mg/kg) and Mg (48.3 mg/kg) and pH (5.04) were determined. The experiment involved rotation system of four crops, namely industrial potato, fodder spring barley, winter rape and winter wheat, as well as an option of organic fertilization based on sewage sediments and composts. The phytopathological studies were carried out during the first three years of the experiment. The studies comprised the following plots: (I) control (no fertilization), (II) mineral NPK fertilization, (III) farm manure 10 t/ha, (IV) farm manure 5 t/ha, (V) Biohum 10 t/ha (municipal sewage sediment with addition of straw, composted, N – 1.07% dry matter, P – 0.75% dm, K – 0.13% dm, Mg – 0.19% dm), (VI) Biohum 5 t/ha, (VII) soil improvement medium 10 t/ha (municipal sewage sediment dried and granulated, N – 1.88% dm, P – 1.28% dm, K – 0.18% dm, Mg – 0.37% dm), (VIII) soil improvement medium 5 t/ha, (IX) municipal sewage sediment from the treatment plant in Tyrowo 10 t/ha (N – 4.66% dm, P – 2.99% dm, K – 0.25% dm, Mg – 0.78% dm), (X) municipal sewage sediment from the treatment plant in Tyrowo 5 t/ha. The farm manure and composted sewage sediments at 10 t/ha were applied once in the crop rotation system, in 2004 before planting potato (Jasia cultivar). Mineral fertilization for potato was 150 kg N (ammonium nitrate 34%), 65 kg P (superphosphate 40%) and 166 kg K/ha (potassium salt 60%). The mineral fertilization on the plots fertilized with NPK was applied in whole before sowing. On the plots with farm manure and sewage sediments the fertilization with N was balanced to 150 kg/ha, depending on their content of total nitrogen, and completed after the main crop with ammonia nitrate. In 2005, only mineral fertilization (90 kg N, 26 kg P and 100 kg K/ha – forms of fertilizers as above) was applied for spring barley (Justyna cultivar); after the harvest and before sowing of winter rape mineral fertilization was applied at 120 kg N, 42 kg P and 134 kg K/ha. Organic fertilization at 5 t/ha was applied only on plots IV, VI, VIII and X. Supplementary fertilization with N was applied on the latter plots to reach 120 kg/h, depending on the content of total nitrogen in composts.

To determine the composition of species and the quantity of fungi, soil samples were collected at a depth of 10 cm in the first week of August of each year from three points on each plot representing specific combinations. In the laboratory

the samples from each plot were combined; test samples of 10 g were weighted to 250 ml flasks and shaken for 20 min in 90 ml of sterile water obtaining the dilution of 10^{-1} . Higher dilutions (10^{-2} to 10^{-4}) were made from that suspension. The fungi was cultured on the Martin's nutrient medium at 22–23°C, and the fungal colonies developed after five days of incubation were converted to grams of dry matter of soil (Dhingra and Sinclair 1995). Next, the fungi were inoculated on agar slopes for later identification of species. At the same time the linear growth of seven potentially pathogenic fungal species, namely *Alternaria alternata*, *Botrytis cinerea*, *Colletotrichum coccodes* and genus *Fusarium*, was assessed on the PDA medium supplemented with water extracts from composts analyzed during the experiment. Each compost type in the form of 2 g of dry matter was flooded with 100 ml of sterile water and after 24 h the extracts were filtered through sterile linen. Petri dishes were filled with 2 ml of the filtrate, covered with 10 ml of cooled medium, and mixed by circular movements. On top of the solidified medium, 5 mm agar discs overgrown with 7-day cultures of fungi were placed. The control was provided by dishes with inoculum of pathogens on medium without the filtrates. The experiment was carried out in 4 repetitions (dishes), twice for each species. After 4 and 8 days the diameters of fungal colonies were measured (along two perpendicular lines); on that basis the growth inhibition coefficient, expressed by the formula:

$$I = [(\phi_k - \phi) / \phi_k] \times 100\%$$

where: ϕ_k and ϕ represent the diameter of the control culture and the culture on the medium with the extract from the composts, was computed

The results were processed statistically and subjected to variance analysis. The Duncan's test was used for comparison of the averages.

RESULTS AND DISCUSSION

The results obtained from the three-year experiment indicate that application of organic fertilization diversified the quantitative and qualitative composition of soil fungi communities. The highest total number of fungal colony-forming units was observed in the soil fertilized with Biohum at 10 t/ha (applied during the first year under cultivation of potato) and twice at 5 t/ha (under potato and before sowing of winter rape), and the

lowest total number of fungal colonies was found in the soil without fertilization and with mineral NPK fertilization (the differences were statistically significant – Figure 1). The literature (Sigler and Turco 2002, Spedding et al. 2004, Larkin and al. 2006) reports that mineral and organic fertilization is, next to the type of crop, rotation system and applied plant protection media, a cause of changes in the structure of soil microorganisms' communities.

Among the selected fungal colonies, 52 species and cultures forming no spores as well as yeasts-like fungi were identified (Table 1). Pathogens were represented by *Alternaria alternata*, *Aureobasidium pullulans*, *A. bollei*, *Botrytis cinerea*, *Colletotrichum coccodes*, *Sclerotinia sclerotiorum* and genus *Fusarium*. The largest number of pathogen colonies was obtained during the second vegetation season of the experiment (9.4%).

Changes in the structure of fungal community between individual setups of the experiment and between sampling dates were found (Figure 2a–j). The most numerous populations of pathogens were obtained from the plots without fertilization (14% of the total number of isolates in the setup – Figure 2a). Among them the following species were identified: *A. pullulans*, *S. sclerotiorum* and genus *Fusarium*. Their numbers increased 2- and 3-fold during the following two years of the experiment, as compared to year one. At the same time the lowest number of saprophytic fungi (23.7%) was obtained from the soil in that combination, as compared to the other combinations. Among them the species of genera *Gliocladium*, *Paecilomyces*, *Penicillium*, *Trichoderma* and order *Mucorales*

were identified. They had a particularly low share in the fungal community during the first vegetation season. As a consequence, it can be assumed that the weakened biological barrier in the form of small numbers of antagonistic fungi was the cause of a more intensive development of the above-indicated pathogens.

Less numerous shares of pathogens in soil fungi community during the three-year study was observed on the plots with mineral NPK fertilization – 10.5% (Figure 2b). During the second year of experiment a decrease in their number by a half in comparison to the beginning of the period was observed in the soil under spring barley, while during the third year their presence was not confirmed. At the same time the number of the antagonistic fungi was twofold higher compared to the soil without fertilization.

Organic fertilization had a positive influence on the structure of soil fungi communities. Its influence was more visible in changes in the qualitative composition of fungi than in their numbers. Hoitink and Boehm (1999) showed that in organically fertilized soil the growth of microorganisms is stimulated; they protect plants against pathogens of genera *Pythium* and *Phytophthora* due to their antibiotic and parasitic influence. In our experiment farm manure showed the best phytosanitary influence among the analyzed organic fertilizers. The share of pathogens in the soil fertilized with farm manure at 10 t/ha was 1.2% (average for three years of studies – Figure 2c). A higher share of these microorganisms (4.1%) was recorded in the soil fertilized by double application of farm manure at 5 t/ha (Figure 2d). In the soil with farm manure




Figure 1. The number of fungal colony-forming units (CFUs) per 1 g soil sample




Figure 2a–f. Fungi isolated from soil of different treatments of experiment (years 2004–2006)

applied once or in two doses a total reduction in number of plant pathogens was observed during the third year of cultivation. This was accompanied by an increase in the population of saprophytic fungi, particularly of genus *Trichoderma*, which could explain the inhibited process of pathogen development. The latter fungi are known for their

lignolytic and cellulolytic properties (Papavizas 1985).

Among the composts made of sewage sediments from the treatment plant applied in the experiment, the soil improvement medium had the best influence on the biological life of the soil. Presence of pathogens was found only during the first year




Figure 2g–j. Fungi isolated from soil of different treatments of experiment (years 2004–2006)

of cultivation (Figure 2g and h). The numbers of saprotrophic organisms in soil with double fertilization showed a decreasing trend over the study period while in the soil with single application of the fertilizer under potato, the largest number of these fungi was obtained during the second year of the study. Abbo-Sedera (2006) observed that under the influence of organic fertilization the number of microorganisms in the potato root zone increased. In the soil fertilized with composts of sewage sediments from the Tyrowo treatment plant the share of pathogens exceeding 11% was found (Figure 2i and j). The dynamics of changes in the fungal community of the soil during the period of three years of the study was more favorable in the case of a single application at 10 t/ha of the fertilizer; a decrease in number of pathogens from 20% during the first year of study to 6% during the following year and to the total reduction of population during the third vegetation season was observed coupled with simultaneous increase of

saprotrophes to ca. 70% in the total population of fungi. Double application at 5 t/ha resulted in a two-fold increase in the number of pathogens during the later two years of the experiment as compared to the first year, coupled with a decrease in the population of saprotrophes.

In the case of experimental setup with Biohum fertilization, the average share of pathogens of 2.8% and 4.8% during the three study years was found for the dose of 10 t/ha and twice 5 t/ha, respectively (Figure 2e and f); however, the changes occurring in the structure of fungal community were not satisfactory. During the two later vegetation seasons, in particular the third one when winter rape was cultivated, a decrease in saprophytes is recorded, which might be caused by the simultaneous increase in the number of pathogens. The modifying influence of plant roots excretions on soil microorganisms may be suggested; it would be in line with the literature (Funk-Jensen and Hockenhull 1984).

Table 1. Fungi isolated from organically fertilized soil during investigation period 2004–2006 (%)

| Fungi | Treatments | | | | | | | | | |
|---|------------|------|------|------|------|------|------|------|------|------|
| | I | II | III | IV | V | VI | VII | VIII | IX | X |
| <i>Acremoniella</i> spp. | 9.7 | | | | | | | | | |
| <i>Acremonium strictum</i> W. Gams | 2.2 | 2.2 | 1.2 | 21.1 | 19.3 | 4.7 | | 2.6 | 3.8 | 3.8 |
| <i>Alternaria alternata</i> (Fr.) Keissler | | | | | | | | | 2.6 | 1.3 |
| <i>Arthrinium sphaeospermum</i> Fuckel | 1.1 | | 1.2 | 1.0 | | 1.9 | 1.2 | | 3.8 | 1.3 |
| <i>Aspergillus</i> spp. | 1.1 | 1.1 | 4.7 | | | | | | | 1.3 |
| <i>Aureobasidium bolleyi</i> | | | | | | | | 3.8 | 1.3 | 1.3 |
| <i>Aureobasidium pullulans</i> (de Bary) Arnaud | 6.5 | 7.4 | | 3.2 | 0.9 | 3.8 | 2.4 | | | 3.9 |
| <i>Botrytis cinerea</i> Pers. | | | | | | | | | | 1.3 |
| <i>Colletotrichum coccodes</i> Hughes (Wallr.) | | 1.1 | | | 0.9 | | | | | |
| <i>Chaetomium globosum</i> Kunze ex Fr. | 1.1 | 11.6 | 1.2 | 13.4 | 0.9 | 1.0 | 2.4 | 7.7 | | 2.6 |
| <i>Cladosporium cladosporioides</i> de (Fres) Vries | | 1.1 | 1.2 | | | 3.8 | 1.2 | 3.8 | 6.4 | 1.3 |
| <i>Coniothyrium minitans</i> Sacc. | | 1.1 | 3.5 | 1.0 | | | | | 2.6 | |
| <i>Endothia</i> spp. | | | | | | 1.0 | | | | 1.3 |
| <i>Epicoccum</i> spp. | | | | | | | | | | 1.3 |
| <i>Fusarium culmorum</i> (W.G. Sm.) Sacc. | 1.1 | | | | | | | | 3.9 | |
| <i>Fusarium equiseti</i> (Corda) Sacc. | | | 1.2 | | | 1.0 | | 3.8 | 1.3 | 1.3 |
| <i>Fusarium oxysporum</i> Schlecht. | 3.2 | | | | 0.9 | | | | | |
| <i>Fusarium poae</i> | | | | | | | | | | 2.6 |
| <i>Gilmaniella humicola</i> Barron | 5.3 | | | | | | | 5.1 | | 3.9 |
| <i>Gliomastix murorum</i> (Corda) Quequen | 1.1 | 3.3 | 1.2 | 6.2 | 1.8 | 1.9 | 2.4 | 7.7 | 1.3 | 3.9 |
| <i>Gliocladium catenulatum</i> Gilman and Abbott | 1.1 | | 3.6 | | | 1.0 | | | 3.8 | |
| <i>Gliocladium penicillioides</i> Corda | | | | | 0.9 | | | | | |
| <i>Humicola brevis</i> Gilman and Abbott | 1.1 | | 1.2 | | | 4.7 | | 1.3 | | 1.3 |
| <i>Humicola fuscoatra</i> Traaen | | | | | | | 1.2 | | | 2.6 |
| <i>Humicola grisea</i> Traaen | 1.1 | 1.1 | | | | | 2.4 | 1.3 | 1.3 | 1.3 |
| <i>Monodictis glauca</i> (Cooke et Harkn.) Hughes | | | | | | | 1.2 | | | |
| <i>Monodictis laevis</i> (Wiltshire) Hughes | | | | | 1.8 | 1.0 | | | | |
| <i>Mortierella acuminata</i> Linn. | | | 8.2 | | 3.6 | | | | | |
| <i>Mortierella alpina</i> Peyronel | | | | 1.0 | 6.4 | 1.0 | | | 2.6 | 3.9 |
| <i>Mortierella isabelina</i> Qudemans | 5.3 | 1.1 | 1.2 | | 10.1 | | 1.2 | 2.6 | 1.3 | 2.6 |
| <i>Mortierella zonata</i> Linn. | | | | | 9.7 | 6.7 | 1.2 | | 6.4 | |
| <i>Mucor hiemalis</i> Wehmer | | 2.2 | 1.2 | 1.0 | 0.9 | 1.0 | | 3.8 | 1.3 | |
| <i>Paecilomyces lilacinus</i> (Thom.) Samson | | | | 4.1 | | | | | | |
| <i>Paecilomyces nivalens</i> (Thom.) Samson | 1.1 | 4.2 | | | | 7.6 | 4.7 | 1.3 | | 2.6 |
| <i>Paecilomyces roseum</i> (Thom.) Samson | | 3.3 | 1.2 | 3.2 | | 1.0 | | | | 1.3 |
| <i>Papulaspora sepedonioides</i> Preuss | | | | | | | | 2.6 | 1.3 | |
| <i>Penicillium</i> spp. | 6.5 | 11.6 | 4.7 | 4.1 | 2.7 | 11.4 | 7.0 | 10.3 | 20.5 | 16.9 |
| <i>Periconia macrospinosa</i> Lefebvre et Johnson | | | | | | | 1.2 | 2.6 | | |
| <i>Phoma eupyrena</i> Sacc. | 1.1 | 6.4 | | | 0.9 | 4.7 | 9.4 | 1.3 | 1.3 | 3.9 |
| <i>Rhizopus nigricans</i> Ehrenberg | | 2.2 | 1.2 | 1.0 | 0.9 | 18.1 | 7.0 | 7.7 | | 10.4 |
| <i>Sclerotinia sclerotiorum</i> (W.G. Sm.) Sacc. | 3.2 | 2.2 | | 1.0 | | | 2.4 | | 2.6 | |
| <i>Sporormia</i> spp. | | | | | | | | | 1.3 | |
| <i>Sporotrichum olivaceum</i> Fries | 1.1 | 4.2 | 1.2 | 2.1 | 0.9 | | 1.2 | 2.6 | 1.3 | 2.6 |
| <i>Taeniolella stilbospora</i> Corda (Hughes) | | 3.3 | | | | | | | | |
| <i>Trichocladium canadense</i> Hughes | | | | 1.0 | | | | | | |
| <i>Trichoderma aureoviride</i> Rifai | | | 3.5 | 3.2 | | 4.7 | | 3.8 | | |
| <i>Trichoderma hamatum</i> (Bon.) Bain | 5.3 | 11.5 | 9.4 | | | 1.9 | 8.2 | 2.6 | 3.8 | 6.5 |
| <i>Trichoderma harzianum</i> Rifai | 3.2 | | 3.5 | | 4.6 | 1.0 | 5.8 | | | |
| <i>Trichoderma koningii</i> Qudemans | | 5.4 | 14.1 | 1.0 | 1.8 | | 1.2 | | 1.3 | 1.3 |
| <i>Trichoderma polysporum</i> (Link: Pers) Rifai | 1.1 | 3.3 | 2.4 | 9.3 | 4.6 | | | | | |
| <i>Trichoderma viride</i> Pers. ex S.F. Gray | | 5.4 | 1.2 | 1.0 | 5.5 | | 2.4 | 3.8 | 1.3 | 5.2 |
| <i>Zygorhynchus</i> spp. | | | | | | | | | | 1.3 |
| Nosporulatis fungi | 5.3 | 1.1 | | 1.0 | 2.7 | 14.3 | 16.3 | 2.6 | 11.3 | 1.3 |
| Yeast-like fungi | 31.2 | 2.4 | 26.8 | 20.2 | 17.4 | 1.0 | 16.3 | 15.4 | 10.3 | 2.6 |
| Total | 186* | 190 | 170 | 194 | 218 | 210 | 172 | 156 | 156 | 154 |

I – control (no fertilization); II – mineral NPK fertilization; III – farm manure 10 t/ha; IV – farm manure 5 t/ha; V – Biohum 10 t/ha; VI – Biohum 5 t/ha; VII – soil improvement medium 10 t/ha; VIII – soil improvement medium 5 t/ha; IX – sewage sediments (treatment plant in Tyrowo) 10 t/ha; X – sewage sediments (treatment plant in Tyrowo) 5 t/ha; *number of isolates




Figure 3. Percentage of pathogen mycelium growth inhibition on PDA with water extracts from composts (a) after 4 days, (b) after 8 days

The authors show that the species of *Brassicaceae* family that produce substances of glucosinolates and isocyanates can inhibit the development of saprophytic species. However, Smith and Kirkegaard (2002) found out that species of *Trichoderma* genus are tolerant to the above compounds. Curl (1982) even stated that plant excretions can improve the phytosanitary condition of soils through a better development of fungi inhibiting the growth of pathogens. As a conclusion, farm manure and soil improving medium were characterized by the best agricultural suitability.

The aquatic extract from composted sewage sediments tested *in vitro* inhibited the growth of mycelia of the studied pathogenic fungi at both times of measurement. The stronger fungistatic influence was found after 8 days than after 4 days of culture. The strongest inhibition of the fungal growth was recorded in the case of Tyrowo compost; fungal mycelium growth inhibition coefficient as compared to the controls was 33.9% and 44.4% during consecutive measurements, respectively (Figure 3). The lowest biological activity was presented by compounds and microorganisms con-

tained in the extract of soil improving medium (22.1% and 11.8 % inhibition of mycelium growth). The *Fusarium poae* and *Colletotrichum coccodes* species were the most susceptible after 4 days of culture and *C. coccodes* and *F. equiseti* after 8 days of culture. *In vitro* studies confirmed the inhibitory influence of composts of sewage sediments on development of selected plant pathogens in soil although differences in effectiveness of individual fertilizers were found, which is understandable because of the differences in conditions. Many authors indicated in their studies the protective influence of extracts from various fertilizers against plant pathogens from genera *Phytophthora*, *Fusarium* and *Rhizoctonia solani* (Aryantha and Guest 2006).

REFERENCES

- Abbo-Sedera S.A. (2006): Evaluation of organic and chemical fertilizers effect on rhizosphere microflora and potatoes (*Solanum tuberosum* L.) yield in loamy sand soil. Arab. Univ. J. Agric. Sci., 14: 613–623.
- Angus J.F., Van Herwaarden A.F., Howe G.N. (1994): Productivity and break crop effects of winter-growing oilseeds. Aust. J. Exp. Agric., 31: 669–677.
- Aryantha I.N.P., Guest D.I. (2006): Mycoparasitic and antagonistic inhibition on *Phytophthora cinnamomi* rands by microbial agents isolated from manure compost. Plant Pathol. J., 5: 291–298.
- Curl A.E. (1982): The rhizosphere relation to pathogen behavior and root disease. Plant Dis., 66: 623–630.
- Dhingra O.D., Sinclair J.B. (1995): Basic Plant Pathology Methods. CRC Press, Boca Raton.
- Funck-Jensen D., Hockenhull J. (1984): Root exudation, rhizosphere microorganisms and disease control. Växtskyddsnotiser, 48: 49–54.
- Grayston S.J., Wang S., Campbell C.D., Edwards A.C. (1998): Selective influence of plant species on microbial diversity in the rhizosphere. Soil Biol. Biochem., 30: 369–378.
- Hoitink H.A.J., Boehm M.J. (1999): Biocontrol within the context of soil microbial communities: A substrate-dependent phenomenon. Annu. Rev. Phytopathol., 37: 427–446.
- Larkin R.P., Honeycutt C.W., Griffin T.S. (2006): Effect of swine and dairy manure amendments on microbial communities in three soils as influenced by environmental conditions. Biol. Fertil. Soils, 43: 51–61.
- Mathur K., Bansal R.K., Gurjar R.B.S. (2006): Organic management of Fusarium wilt of fenugreek (*Trigonella foenumgraecum* L.) – a seed spice. Indian J. Mycol. Plant Pathol., 36: 94–95.
- Papavizas G.G. (1985): *Trichoderma* and *Gliocladium*: Biology, ecology and potential for biocontrol. Annu. Rev. Phytopathol., 23: 23–54.
- Shannon D., Sen A.M., Johnson D.B. (2002): A comparative study of the microbiology of soils managed under organic and conventional regimes. Soil Use Manag., 18: 274–283.
- Sigler W.V., Turco R.F. (2002): The impact of chlorothalonil application on soil bacterial and fungal populations as assessed by denaturing gradient gel electrophoresis. Appl. Soil Ecol., 21: 107–118.
- Smith B.J., Kirkegaard J.A. (2002): *In vitro* inhibition of soil microorganisms by 2-phenylethylisothiocyanate. Plant Pathol., 51: 585–593.
- Spedding T.A., Hamel C., Mehugs G.R., Madramootoo C.A. (2004): Soil microbial dynamics in maize-growing soil under different tillage and residue management systems. Soil Biol. Biochem., 36: 499–512.

Received on December 12, 2007

Corresponding author:

Dr. hab. Bożena Cwalina-Ambroziak, University of Warmia and Mazury, Department Phytopathology and Entomology, Prawocheńskiego 17, 10-720 Olsztyn, Poland
phone: + 488 952 341 47, e-mail: bambr@uwm.edu.pl
