

Anthocyanins Content in Blueberries (*Vaccinium corymbosum* L.) in Relation to Freezing Duration

A. VOLLMANNOVÁ*, T. TÓTH, D. URMINSKÁ, Z. POLÁKOVÁ, M. TIMORACKÁ
and E. MARGITANOVÁ

Faculty of Biotechnology and Food Sciences, Slovak University of Agriculture in Nitra,
949 76 Nitra, Slovak Republic, *E-mail: alena.vollmannova@uniag.sk

Abstract: The changes of total anthocyanins (TA) content in fruit of blueberries in relation to freezing duration are the aim of this study. The samples of 21 chosen cultivars obtained from the research breed station were stored in polyethylene bags in freezing box at temperature -18°C . In ethanolic blueberry extracts TA contents were by spectrophotometric method during the time of 6 months at period of 3 months determined. Significant differences of TA content among the observed cultivars were confirmed. The value interval was 615.8–3206.3 mg/kg. The average TA content in 10 very early, resp. early cultivars was 1655.8 mg/kg, in 5 medium early cultivars 1515.6 mg/kg, in 6 medium late, resp. late cultivars 1431.2 mg/kg. During 6 months of blueberries store in the freezing box the average decrease 16.7% of TA content was observed. The average decrease of TA content in very early, resp. early cultivars was 19.6%, in medium early cultivars 16.5% and in medium late, resp. late cultivars 14%.

Keywords: cultivars; blueberries; anthocyanin; freezing

INTRODUCTION

The positive influence of phytochemicals present in fruit and vegetable is accounted for their regular consumption. Small fruits as raspberries, blackberries, blueberries, blackcurrants are rich in phytochemicals such as anthocyanins which are glycosidic-linked flavonoids responsible for their red, violet, purple and blue colours. Berries are edible superstars that may protect against heart disease, cancers and ageing (BEATTIE *et al.* 2005). In many works a positive correlation between the intake of fruits and vegetables and prevention of diseases like atherosclerosis, cancer, diabetes, arthritis and also ageing is documented (KAUR & KAPOOR 2001). Bioactive extracts of blueberries are rich in anthocyanins and proanthocyanins with a significant antioxidant activity (SMITH *et al.* 2000) which is influenced by various factors (temperature, pH, oxidation) and is about 3-fold higher than either strawberries or raspberries (KALT *et al.* 1999). Total phenolics and total ant-

hocyanin concentrations in wild crop are higher than in cultivated blueberries (GIOVANELLI & BURATTI 2009). The changes of total anthocyanins (TA) content in fruit of blueberries in relation to freezing duration are the aim of this study.

MATERIAL AND METHODS

The samples of 10 very early, resp. early (VE, E), 5 medium early (ME) and 6 medium late, resp. late (ML, L) blueberry cultivars were obtained from the research breed station in Kriva on Orava. Locality of blueberry cultivation is characterised by altitude 700 m, average yearly temperature 6°C and precipitation 800 till 900 mm. From manually collected blueberries 100 g samples were weighted and stored in PE bags in freezing box at temperature -18°C . From blueberries samples 50 g were homogenised and extracted by 100 ml 80% ethanol during 12 hours. In obtained extracts total anthocyanins contents by modified method of LAPORNIK *et al.*

(2005) were spectrophotometrically (Shimadzu UV/VIS – 1240) determined. Into two tubes 1 cm³ of extract was pipetted and 1 cm³ 0.01% HCl in 80% ethanol was added. Then 10 cm³ 14% HCl into the first tube and 10 cm³ McIlvain agents (pH 3.5) into another tube were added. Absorbance was measured at 520 nm against blank sample. The obtained results were statistically evaluated by ANOVA single factor analyse ($\alpha = 0.01$).

RESULTS AND DISCUSSION

Significant differences of total anthocyanin content among the observed cultivars in interval 615.8 mg/kg (Puru) – 3206.3 mg/kg (Northland) were confirmed (Table 1).

The statistically significant differences in TA content among blueberry cultivars were confirmed also by ANOVA single factor analyse ($\alpha = 0.01$). In VE, E cultivars P -value = 4.12E-38, in ME cultivars P -value = 2.53E-15 and in ML, L cultivars P -value = 5.99E-20.

The average TA content in 10 VE, E cultivars was 1942.3 mg/kg, in 5 ME cultivars 1515.5 mg/kg, in 6 ML, L cultivars 1431.1 mg/kg. During first 3 months of blueberries store in the freezing box the average decrease (12.6%) of TA content was observed. At the same time significant differences among the observed cultivars were confirmed. The average decrease value of TA content in VE, E cultivars was 16.6%, in ME cultivars 7.5% and in ML, L cultivars 4.3%. The average decrease value of TA content after 6 months of blueberry storage at – 18°C in relationship to the first measurement in VE, E cultivars was 19.5%, in ME cultivars 16.5% and in ML, L cultivars 14%.

Our results correspond with results of GAO and MAZZA (2006). They determined simple and acylated anthocyanins and other phenolics in 10 cultivars and hybrids of lowbush (*Vaccinium corymbosum*) and one cultivar of highbush (*V. angustifolium*) blueberries. TA content in blueberries ranged from 110 to 260 mg/100 g of fresh berries. Similar results (12.7–197.34 mg/100 g f.m.) are presented also by SELAPAN *et al.* (2002).

Table 1. Total anthocyanins content (mg/kg) in blueberry samples during time storage at temperature –18°C

| Group of cultivars | Cultivar | I. (0 months) | II. (3 months) | III. (6 months) |
|------------------------------------|---------------|---------------|----------------|-----------------|
| Very early, resp. early (VE, E) | Goldtraube 23 | 1971.57 | 1679.83 | 1033.04 |
| | Earlyblue | 1860.88 | 1202.64 | 1660.78 |
| | Sunrise | 2013.92 | 1461.46 | 1531.16 |
| | Puru | 615.78 | 566.7 | 575.08 |
| | Duke | 2152.18 | 1809.73 | 1934.3 |
| | Nui | 1724.56 | 1797.3 | 1642.68 |
| | Reka | 900.47 | 1023.54 | 1140.05 |
| | Spartan | 2073.89 | 1476.69 | 1596.44 |
| Medium late, resp. late (ML, L) | Goldtraube | 2903.9 | 2399.34 | 2324.3 |
| | Northland | 3206.29 | 2144 | 2177.54 |
| | Pemberton | 2145.2 | 1599.79 | 1419.8 |
| | Blueray | 1335.1 | 1281.48 | 1052.16 |
| Medium late, resp. late (ML, L) | Bluejay | 1398.48 | 1217.55 | 1485.86 |
| | Patriot | 1882.17 | 1745.82 | 1547.9 |
| | Bluecrop | 816.75 | 917.96 | 821.66 |
| | Nelson | 1852.74 | 1810.47 | 1579.1 |
| Medium late, resp. late (ML, L) | Berkley | 1581.8 | 1416.17 | 1087.3 |
| | Brigitte | 1116.26 | 1090.4 | 1030.18 |
| | Persey | 1603.21 | 1382.46 | 1264.44 |
| | Herbert | 1183.13 | 1028.37 | 1211.2 |
| | Darrow | 1249.01 | 1454.01 | 1212.54 |

CHO *et al.* (2005) determined in blueberries higher TA contents (1435.2–8227.3 mg/kg). Changes of antioxidant activity of blueberries in relation to their TA, polyphenols and flavonols content and their cultivar dependence are confirmed also by VELIOGLU *et al.* (1998). Our results are in agreement also with data published by HÄKKINEN *et al.* (1999). They confirmed, that concentrations of phytochemicals in berries are influenced by many factors including environmental conditions, degree of ripeness, cultivar, cultivation site, processing and storage of the fruit. After SCIBISZ and MITEK (2007) the long-term frozen storage of blueberries did not induce significant changes in TA content. It is most probable that the anthocyanin in frozen fruits becomes more easily extractable. An increase in TA content in raspberry during freezing has been reported by ANCOS *et al.* (2000). GONZÁLEZ *et al.* (2003) documented anthocyanins losses in frozen blackberry stored for 12 months at -24°C . Other investigations have demonstrated dramatic losses of anthocyanins during frozen storage. CHAOVANALIKIT and WROLSTAD (2004) reported 88% anthocyanins degradation in sweet cherries after 6 months of storage at -23°C . Our results confirmed 16.7% average decrease of TA content after 6 months of blueberries storage at -18°C .

Acknowledgements: The work was supported by the Scientific Grant Agency (VEGA) of the Ministry of Education of the Slovak Republic and the Slovak Academy of Sciences (Grants KEGA 3/5081/07 and VEGA 1/0030/09).

References

- ANCOS B., GONZALEZ E.M., CANO M.P. (2000): Ellagic acid, vitamin C, and total phenolic contents and radical scavenging capacity affected by freezing and frozen storage in raspberry fruit. *Journal of Agricultural and Food Chemistry*, **48**: 4565–4570.
- BEATTIE J., CROSIER A., DUTHIE G.D. (2005): Potential health benefits of berries. *Current Nutrition & Food Science*, **1**: 71–86.
- CHAOVANALIKIT A., WROLSTAD R.E. (2004): Anthocyanin and polyphenolic composition of fresh and processed cherries. *Journal of Food Science*, **69**: 73–83.
- CHO M.J., HOWARD L.R., PRIOR R.L., CLARK J.R. (2005): Flavonol glycosides and antioxidant capacity of various blackberry and blueberry genotypes determined by high-performance liquid chromatography/mass spectrometry. *Journal of the Science of Food and Agriculture*, **85**: 2149–2158.
- GAO L., MAZZA G. (2006): Quantitation and distribution of simple and acylated anthocyanins and other phenolics in blueberries. *Journal of Food Science*, **59**: 1057–1059.
- GIOVANELLI G., BURATTI S. (2009): Comparison of polyphenolic composition and antioxidant activity of wild Italian blueberries and some cultivated varieties. *Food Chemistry*, **112**: 903–908.
- GONZÁLEZ E., DE BEGONA A., CANO M. (2003): Relation between bio active compounds and free radical – scavenging capacity in berry fruits during frozen storage. *Journal of the Science of Food and Agriculture*, **83**: 722–726.
- HÄKKINEN S., HEINONEN M., KÄRENLAMP I., MYKKÄNEN H., RUUSKANEN J., TÖRRÖ R. (1999): Screening of selected flavonoids and phenolic acids in 19 berries. *Food Research International*, **32**: 345–353.
- KALT W., FORNEY C.F., MARTIN A., PRIOR R.L. (1999): Antioxidant capacity, vitamin C, phenolics, and anthocyanins after fresh storage of small fruits. *Journal of Agricultural and Food Chemistry*, **47**: 4638–4644.
- KAUR C., KAPOOR H.C. (2001): Antioxidants in fruits and vegetables – the millennium's health. *International Journal of Food Science & Technology*, **36**: 703–725.
- LAPORNIK B., PROŠEK M., WONDRA A.G. (2005): Comparison of extracts prepared from plant by-products using different solvents and extraction time. *Journal of Food Engineering*, **71**: 214–222.
- SCIBISZ I., MITEK M. (2007): The changes of antioxidant properties in highbush blueberries (*Vaccinium corymbosum* L.) during freezing and long-term frozen storage. *Acta Scientiarum Polonorum Technologia Alimentaria*, **6**: 75–82.
- SELLAPAN S., AKOH C.C., KREWER G. (2002): Phenolic compounds and antioxidant capacity of Georgia-grown blueberries and blackberries. *Journal of Agricultural and Food Chemistry*, **50**: 2432–2438.
- SMITH M.A.L., MARLEY K.A., SEIGLER D., SINGLETARY K.W., MELINE B. (2000): Bioactive properties of wild blueberry fruits. *Journal of Food Science*, **65**: 352–356.
- VELIOGLU Y.S., MAZZA L., GAO L., OOMAH B.D. (1998): Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *Journal of Agricultural and Food Chemistry*, **46**: 4113–4117.