

Importance of *Enterococcus* spp. for Forming a Biofilm

L. NECIDOVÁ^{1*}, B. JANŠTOVÁ¹, S. KARPÍŠKOVÁ², Š. CUPÁKOVÁ¹, M. DUŠKOVÁ¹
and R. KARPÍŠKOVÁ¹

¹Department of Milk Hygiene and Technology, University of Veterinary and Pharmaceutical Sciences Brno, 612 42 Brno, Czech Republic; ²Czech Collection of Microorganisms, Faculty of Science Masaryk University, 602 00 Brno, Czech Republic, *E-mail: necidoval@vfu.cz

Abstract: The aim of this study was to monitor the capability of *Enterococcus faecalis* and *Enterococcus fecium* to form biofilms. Enterococci isolates originated from individual milk, bulk milk samples and environmental swabs obtained at farm level, dairy plant level including semi and final dairy products. Biofilm formation potential was determined by growing the tested strains in glass tubes containing BHI medium. The capability of forming biofilms was detected in 28% of *Enterococcus* spp. strains. Higher number of biofilm forming strains of the *Enterococcus faecium* (33%) than *Enterococcus faecalis* (28%) has been registered. Isolates obtained at plant level were forming biofilms more often than isolates from farm level and in final products (cheese and curd cheese), no isolate has been seen to be able to form biofilm.

Keywords: biofilm; *Enterococcus faecalis*; *Enterococcus fecium*; dairy products; bacteria

INTRODUCTION

Permanently wet surfaces of the food industry facilities often support the adherence and colonisation of bacteria with subsequent formation of a biofilm. Bacteria in biofilm structures are more resistant not only to detergents but also against drying than planktonic ones (BOLTON *et al.* 1988). Biofilm formation in food industry represents a serious issue (MCLANDSBOROUGH *et al.* 2006).

Nowadays, enterococci take up a significant position among the bacteria causing nosocomial infections. One of the main factors of enterococci virulence is the biofilm formation (TENDOLKAR *et al.* 2006). Enterococci are a part of common intestinal microflora in mammals and therefore they are often present in raw material such as raw milk and meat.

The aim of this study was to assess the capability of forming biofilm in two enterococci species isolated from the samples at farm and dairy plant level.

MATERIAL AND METHODS

Enterococci isolates. Enterococci strains originating from the strain collection of Department of Hygiene and Milk Hygiene and Technology (University of Veterinary and Pharmaceutical Sciences, Brno) isolated from dairy chain in 2005–2007 and stored at –75°C were used in this study.

Genus and species identification of enterococci. Genus specific identification was carried out using the PCR method based on the detection of genus specific *tuf*-gene (CUPÁKOVÁ *et al.* 2005). *Enterococcus faecalis* and *Enterococcus fecium* identification was carried out using the PCR method based on the detection of genus specific sections of the *sodA* gene encoding the enzyme manganese-dependent superoxide dismutase (JACKSON *et al.* 2004).

Detection of biofilm formation. For the assessment of the capability of biofilm formation in enterococci the phenotypic method was used. Suspension of tested strain was incubated in the glass tubes containing Brain Heart Infusion Broth (BHI

broth) aerobically at the temperature of 35°C for the period of 2 days. Then the supernatant was discarded, the glass tube has been stained by 0.1% safranin solution, washed with distilled water three times and dried. In the case of biofilm formation, a grainy red structure on the test tube bottom was found. All strains were tested in triplicates. The test tube finding was compared with the biofilm formation in reference strains (*Staphylococcus epidermidis* CCM 4418, CCM 7221).

RESULTS AND DISCUSSION

In total, 324 strains (303 strains of *Enterococcus faecalis* and 21 strains of *Enterococcus faecium*) were tested.

Enterococcus spp. biofilm formation capability was proven in 92 strains (28%), 232 strains (72%) was assessed to be biofilm negative.

From this study results, higher capability of biofilm formation can be seen in strains of the *Enterococcus faecium* species (Table 1). Monitoring of biofilm formation in *E. faecalis* and *E. faecium*

was carried out by numerous authors who present variable results. Majority of the studies are focusing of the characteristics of clinical isolates from various sources. The conclusions of some of the studies indicate that *Enterococcus faecalis* forms biofilm more often than *Enterococcus faecium* (PRAKASH 2005; DI ROSA *et al.* 2006). The results of our study do not confirm this findings, however, the biofilm forming capability will most probably depend on the strains origin and the conditions of their cultivation.

Table 2 shows the distribution of biofilm positive enterococci according to the locality (farm and dairy plant). Isolates from farm level show a higher number of biofilm forming strains. The biofilm forming strains originated mainly from the bulk tank milk samples. Enterococci with both positive and negative biofilm formation were detected in swabs at the dairy plant level (production equipment, milk and cream samples prior to pasteurisation). In final products – cheese and curd cheese only strains incapable of biofilm formation were found. One of the reasons why enterococci isolated from the final products did not form biofilm

Table 1. Biofilm formation capability in the *Enterococcus faecalis* and *Enterococcus faecium* species

	Number of biofilm positive strains	Number of biofilm negative strains
<i>Enterococcus faecalis</i>	85 (28%)	218 (72%)
<i>Enterococcus faecium</i>	7 (33%)	14 (67%)

Table 2. *Enterococcus* spp. with a detected biofilm forming capability sorted by the sampling locations

Sample locality	Sample type	Number of biofilm positive strains			Enterococci in total
		<i>Enterococcus faecalis</i>	<i>Enterococcus faecium</i>		
Farm 1	milk	9	0	9	19.5%
	swab	7	0	7	
Farm 2	milk	7	0	7	44.0%
	swab	4	0	4	
Farm 3	milk	11	0	11	34.7%
	swab	6	0	6	
Farm 4	milk	5	1	6	55.0%
	swab	3	2	5	
Dairy plant	swab	22	4	26	24.3%
	semi-products	6	0	6	
	final products	4	0	4	

could be due to the presence of sodium chloride in cheese (up to 4%) and a higher acidity of curd cheese (up to 70 SH). Several authors state that biofilm formation is impacted by the nutrient content in the cultivation medium, glucose, blood serum, iron, CO₂ and pH (PILLAI *et al.* 2004). In *E. faecalis* as well as *E. faecium* strains isolated from three samples of butter and from one sample of whipped buttermilk, biofilm formation was confirmed.

CONCLUSION

A preventive measure against biofilm formation is the use of suitable production technologies. One of the repressive measure is efficient surface cleaning and sanitation of all the surfaces that come in touch with the raw materials.

Acknowledgements: This study was supported by Project No. 6215712402 "Veterinary Aspects of Food Safety and Quality" of the Ministry of Education, Youth and Sports of the Czech Republic, and by Project No. QF4048 of the Ministry of Agriculture of the Czech Republic.

References

- BOLTON K.J., DODD C.E.R., MEAD G.C., WAITES W.M. (1988): Chlorine Resistance of strains of *Staphylococcus aureus* isolated from poultry-processing plants. Letters in Applied Microbiology, **6**: 31–34.
- CUPÁKOVÁ Š., POSPÍŠILOVÁ M., KOLÁČKOVÁ I., KARPÍŠKOVÁ R. (2005): Genus-specific identification of enterococci by PCR method. Acta Veterinaria Brno, **74**: 633–637.
- DI ROSA R., CRETU R., VENDITTI M., D'AMELIO R., ARCIOLA C.R., MONTANARO L., BALDASSARRI L. (2006): Relationship between biofilm formation, the enterococcal surface protein (Esp) and gelatinase in clinical isolates of *Enterococcus faecalis* and *Enterococcus faecium*. FEMS Microbiology Letters, **256**: 145–150.
- JACKSON CH., FEDORKA-CRAY P.J., BARRETT J.B. (2004): Use of a genus- and species-specific multiplex PCR for identification of enterococci. Journal of Clinical Microbiology, **42**: 3558–3565.
- MCLANDSBOROUGH L., RODRIGUEZ A., PEREZ-CONESA D., WEISS J. (2006): Biofilms: At the interface between biophysics and microbiology. Food Biophysics, **1**: 94–114.
- PILLAI S.K., SAKOULAS G., ELIOPOULOS G.M., MOELLERLING R.C. JR., MURRAY B.E., INOUZE R.T. (2004): Effect of glucose on *fsr*-mediated biofilm formation in *Enterococcus faecalis*. Journal of Infect Diseases, **190**: 967–970.
- PRAKASH V.P. (2005): Clinical prevalence, identification and molecular characterization of enterococci. [Ph D. Thesis.] Pondicherry University, India. In: MUHAMED J.A., HUANG D.B. (eds): Biofilm formation by enterococci. Journal of Medical Microbiology, **56**: 1581–1588.
- TENDOLKAR P.M., BAGHDAYAN A.S., SHANKAR N. (2006): Putative surface proteins encoded within a novel transferable locus confer a high-biofilm phenotype to *Enterococcus faecalis*. Journal of Bacteriology, **188**: 2063–2072.