

Possibility to predict the yield of potatoes grown under two crop production systems on the basis of selected morphological and physiological plant indicators

KRYSTYNA ZARZYŃSKA*, MILENA PIETRASZKO

Plant Breeding and Acclimatization Institute, Potato Agronomy Department, Jadwisin, Serock

*Corresponding author: k.zarzyńska@ihar.edu.pl

ABSTRACT

Zarzyńska K., Pietraszko M. (2017): Possibility to predict the yield of potatoes grown under two crop production systems on the basis of selected morphological and physiological plant indicators. *Plant Soil Environ.*, 63: 165–170.

In the study conducted in the years 2014–2016 at the Institute of Plant Breeding and Acclimatization in Poland, the effect was tested of such indicators of plant productivity as: leaf area index (LAI), leaf greenery index (SPAD) and chlorophyll *a* fluorescence on the yield of potato tubers grown in two production systems, i.e. organic and conventional. It was found that in the organic production system the values of all tested indicators were significantly lower than in the conventional system and the decrease in chlorophyll content in the leaves, as well as the decrease in chlorophyll *a* fluorescence over time followed faster in the organic than in conventional system. The lower surface of the leaves, smaller greenery index and lower activity of photosynthesis contributed to a significant reduction in tuber yield in the organic system. The tuber yield in this system was about 30% lower than in the conventional one. The positive correlations between the value of the tested indicators and yield of tubers was obtained. The highest correlation was between the LAI index, the smallest between the performance index of photosystem II. In the conventional system the correlation coefficients were slightly higher than in organic, which indicates that in this system the possibility of predicting the final yield is simpler.

Keywords: *Solanum tuberosum* L.; tuberous crop; organic plantation; photosynthetic activity; fertilizer

Commonly, potato is grown under the conventional system, but more often the integrated and organic systems are met. Global trends are going in the direction of environmentally friendly systems. However, in these systems, especially in the organic ones, there are certain restrictions that limit the level of yields. There are limits on the use of pesticides and in some soils deficit of nutrients appears as a result of non-mineral fertilizers. These limitations affect plant growth and, consequently, the tuber yield and its structure. Plants growing on the organic plantation are characterized by lower aboveground weight and smaller leaf greenness (Zarzyńska and Szutkowska 2012, Zarzyńska and Pietraszko 2015). It can be assumed that their photosynthetic activity is also lower compared to the conventional system.

Potato is one of the most efficient plants in converting solar energy into human food, both quantitatively and qualitatively. According to Milthrope and Moorby (1997) the rate of tuber mass increase is determined by the time of tuberization and the state of photosynthetically active leaves. Photosynthesis is one of the basic physiological processes of plants with internal and external conditions. Any reduction of the intensity of this process causes a decrease in the amount and quality of crop plants. One of the indicators for early prediction of potato yielding can be the measurement of chlorophyll fluorescence. It is an informative tool for studying the effects of different environmental stresses on photosynthesis (Strasser et al. 2000). In the research work, the chlorophyll fluorescence techniques are used for

doi: 10.17221/101/2017-PSE

Table 1. Agronomic inputs in organic and conventional systems

| Crop production practice | Organic system | Conventional system |
|--------------------------------|--|--|
| Fertilization | manure – 28 t/ha + catch crop | 4–5 t ploughed rye straw + 1 kg mineral nitrogen per 100 kg straw + catch crop. (100 kg N/ha; 53 kg P/ha; 150 kg K/ha) |
| Weed control | only mechanical tillage | mechanical tillage + herbicides 2–3 times per season |
| Colorado potato beetle control | biological insecticide (<i>Bacillus thuringiensis</i>) 2–3 times per season | chemical insecticides 2–3 times per season |
| Late blight control | 2–3 copper fungicides | 3–5 chemical fungicides |

assessment of potato genotypes tolerant and sensitive to environmental stresses (Jefferies 1992, Mauromicale et al. 2006, Ierna 2007).

The aim of the study was to evaluate differences in the selected indicators of some morphological-physiological traits of potato plants grown in two production systems, i.e. organic and conventional, and to assess which of these indicators is the most correlated with the yield and whether it is possible to predict the final yield on this basis.

MATERIAL AND METHODS

A study was conducted in 2014–2016 at the Plant Breeding and Acclimatization Institute, in central Poland on a light loamy sand. Potatoes were grown in the two systems, i.e. organic and conventional. In the organic system the following crop rotation was used: potatoes → field peas → oat → rye with serradella undersown → buckwheat + white mustard as catch crops.

In the conventional system the crop rotation was: potatoes → spring wheat → winter wheat → lupines. The two systems also differed in fertilization, weed control and insect control practices. Late blight control consisted of 2 to 3 foliar applications of copper fungicides in the organic system, and

3 to 5 applications of chemical fungicides in the conventional system (Table 1).

Weather conditions during three years of the trial were varied. In the years 2014 and 2016, both the amount and distribution of rainfall and air temperature were favourable for the growth and development of potato plants, while in 2015 there was an shortage of rainfall, especially in June and August, so the irrigation making up the water shortages in both systems was done. The meteorological data for 3 years of investigations are presented in Table 2.

Four Polish potato table cultivars were chosen for this study based on maturity class and resistance to late blight (Table 3). All cultivars were planted at the same time i.e. about the 20th April. Plot size was 84 m² in three replications. Plants were grown with the spacing of 75 × 33.3 cm.

At the end of June (for the early cultivars) and 2 weeks later (for the mid-early ones) total above-ground biomass, and leaf area index were assessed. The assimilation area of leaves was measured by using the Li-3100 area meter (Lincoln, USA). The LAI was calculated as a ratio of assimilation area of one plant to the area of ground occupied per one plant. The measurements were conducted on four plants from each of the three replicates. In addition, six measurements of chlorophyll content and chlo-

Table 2. Total monthly rainfall (R, mm) and mean monthly temperatures (T, °C) during the vegetative growth period in the years 2014–2016 for Jadwisin

| Year | May | | June | | July | | August | | September | |
|------|------|------|------|------|-------|------|--------|------|-----------|------|
| | R | T | R | T | R | T | R | T | R | T |
| 2014 | 41.3 | 14 | 69.8 | 15.8 | 23.5 | 21.4 | 79.2 | 18.3 | 11.1 | 14.8 |
| 2015 | 39.5 | 12.9 | 15.4 | 17.5 | 62.6 | 19.7 | 8.6 | 22.5 | 36.6 | 15.1 |
| 2016 | 92.2 | 15.3 | 85.4 | 18.7 | 103.6 | 19.6 | 61.4 | 18.4 | 9.5 | 15.7 |

Table 3. Characteristics of four potato cultivars evaluated in organic and conventional production systems during 2014–2016

| Cultivar | Maturity | Resistance to <i>Phytophthora infestans</i> |
|-----------|-----------|---|
| Ignacy | early | 3 |
| Michalina | early | 3 |
| Malaga | mid early | 3.5 |
| Oberon | mid early | 3.5 |

1 – no resistance; 9 – full resistance

rophyll *a* fluorescence were made at 10-day intervals starting from about 45 days after planting (DAP). Evaluation of total chlorophyll content was based on a leaf greenness index (SPAD) measured by SPAD 502- Soil-Plant Analyses Development (Minolta, Japan) chlorophyll meter, measured on apex leaf.

Chlorophyll *a* fluorescence measurements were performed on the plants with a Pocket PEA (plant efficiency analyser, Hansatech, UK). The youngest fully expanded leaves were kept in darkness for at least 20 min in specially provided clips. Determined parameters were: F_v/F_m – the ratio of variable to maximal chlorophyll fluorescence (the photochemical efficiency of photosystem II) and PI – the performance index of photosystem II (Strasser et al. 2000). Measurements were made on weekdays from 7:30–9:00 a.m. with the saturation irradiance up to 3500 $\mu\text{mol}/\text{m}^2/\text{s}$.

Final harvest was performed after full maturity of plants, between August 5 and 12.

RESULTS

The significance of the studied factors differentiation. The analysis of variance showed the

Table 4. Significance of the tested parameters

| Tested parameter | Crop production system | Cultivar | Year |
|---|------------------------|----------|------|
| Leaf area index | ** | ns | ns |
| Leaf greenness index | ** | * | ns |
| Ratio of variable to maximal chlorophyll fluorescence | ns | ns | ns |
| Performance index | ** | * | ns |
| Total yield | ** | ns | ns |

** $\alpha \leq 0.01$; * $\alpha \leq 0.05$; ns – non significant

importance of the diversity of all studied factors depending on the production system except of the ratio F_v/F_m . Significant varietal differences were observed only in the leaf greenness index and the performance index of photosystem II. Years of research did not significantly differentiate any of the studied parameters (Table 4). There was a significant interaction of the years of research and production system.

Diversity of morphological and physiological plant parameters depending on the tested factors. In the conventional system, the values of all studied parameters were higher than in the organic system. The only exception was the rate of F_v/F_m that did not differ in both systems. The biggest differences were related to LAI index; in the conventional system, its value was almost 40% higher than in the organic. Much less variation was observed in the case of leaf greenness index and the performance index of photosystem II) parameters. The average value of SPAD was by 11.5 and PI by 14.1% higher in the conventional system. This diversity contributed to the tuber yield. The yield obtained in the conventional system was by 30.5% higher than in the organic one (Table 5).

Table 5. Differences of plant morphological and physiological parameters depending on the crop production system

| Tested parameter/production system | Conventional | Organic | Decrease in relation to conventional system (%) |
|---|--------------|---------|---|
| Leaf area index | 2.97 | 1.81 | 39.0 |
| Leaf greenness index | 42.3 | 37.4 | 11.5 |
| Ratio of variable to maximal chlorophyll fluorescence | 0.79 | 0.79 | 0 |
| Performance index | 3.82 | 3.28 | 14.1 |
| Yield (t/ha) | 51.4 | 35.7 | 30.5 |

doi: 10.17221/101/2017-PSE

Table 6. Cultivar differences of plant morphological and physiological parameters

| Cultivar | Leaf greenness index | Performance index |
|-----------|----------------------|-------------------|
| Ignacy | 39.7 ^b | 3.54 ^b |
| Michalina | 38.8 ^b | 3.68 ^b |
| Malaga | 37.1 ^a | 3.05 ^a |
| Oberon | 43.9 ^c | 3.94 ^c |

^{a,b,c}Mean values indicated by the same letters are not statistically significant at the 0.05 level by the Tukey's test

The cultivars significantly differed in the amount of the two indicators, i.e. SPAD and PI.

The biggest value of leaf greenness was recorded in cv. Oberon, the lowest in cv. Malaga. The highest rate of PI was also noted for cv. Oberon and the lowest for cv. Malaga (Table 6).

Although the significant differences of years did not prove concerning in tuber yield, significant differences were found in yielding of potatoes grown in both production systems in subsequent years. In 2015, the highest yield was obtained in the conventional system but the lowest in the organic one.

Changes of SPAD index values in conventional and organic systems over time. Performed every 10 days, the measurement of leaf greenness and the chlorophyll *a* fluorescence showed a variation of these parameters depending on the production system. At the beginning of the plant development, the chlorophyll content in the leaves of both systems was similar. Differentiation followed with the development of plants. The highest content of chlorophyll was observed in both systems at the

beginning of the measurements. In the organic system the chlorophyll content was generally lower and declined more rapidly over time than in the conventional system (Figure 1).

Similar situation is related to the chlorophyll *a* fluorescence. Photosynthetic activity of leaves determined by PI index was the highest in the initial period of plant growth and decreased as they developed. In the organic system, the chlorophyll *a* fluorescence was lower than in the conventional system and falling faster than in the conventional system. In the early stages of plant growth the photosynthesis intensity in both production systems was at a similar level (Figure 2).

The relationships between the value of morphological-physiological parameters and tuber yield. The calculated correlation coefficients between the value of the studied parameters and yield of tubers obtained in both production systems showed the highest relationship between LAI, then between SPAD index and the lowest between PI ratio and tuber yield. The correlation coefficients were, respectively, as follow: in the conventional system: 0.702, 0.570 and 0.381; in the organic system, they were slightly lower and amounted to: 0.687, 0.520 and 0.323 (Table 7).

DISCUSSION

Prohibition or major restrictions in the use of chemical and fertilizer inputs in the organic production system resulted in weaker plant growth, and consequently reduced tuber yield and smaller tuber size. Development of the aboveground plant




Figure 1. Effect of production system (conventional and organic) on leaf chlorophyll content (SPAD index) from 45 to 95 days after planting (DAP). Values are means of 4 cultivars and three years




Figure 2. Effect of production system (conventional and organic) on the performance index of photosystem II from 45 to 95 days after planting (DAP). Values are means of 4 cultivars and three years

Table 7. Correlation coefficients between the chosen morphological-physiological parameters and tuber yield. Values are means of 4 cultivars and three years

| Production system | Leaf area index | | Leaf greenness index | | Performance index | |
|-------------------|-----------------|-----------------------|----------------------|-----------------------|-------------------|-----------------------|
| | <i>P</i> | <i>r</i> ² | <i>P</i> | <i>r</i> ² | <i>P</i> | <i>r</i> ² |
| Conventional | 0.006 | 0.702 | 0.022 | 0.570 | 0.032 | 0.381 |
| Organic | 0.007 | 0.687 | 0.028 | 0.520 | 0.030 | 0.323 |

Statistically significant difference at *P*-level of significance $P < 0.05$

parts was largely determined by supply and availability of nitrogen in the first weeks after emergence (Van der Zaag 1992, Marscher 1995, Vos 1995, Goffart et al. 2008) and the time of flowering, when the tuberization process begins. For maximum tuberization, it is necessary to achieve a certain level of development of the aboveground plant parts, i.e. LAI ranging from 2.5 to 3 (Marschner 1995, Van Delen 2001). In our study, the average LAI in organic production system was only 1.8 and in the conventional system around 3. This difference in plant development was reflected in tuber yield that was 30% lower in organic system, although it should be emphasized that the resulting yields were as high as in organic system. In this study, it was found that at the beginning of the growing season the leaf chlorophyll content was similar in both production systems. Leaf chlorophyll content is closely related to nitrogen status of the plants (Tremblay 2004, Goffart et al. 2008, Ramirez et al. 2014). A decline in leaf chlorophyll content in the organic production system occurred starting about 50 DAP. This could have been related to a decline in available nitrogen from organic fertilizer sources. In conventional system, it started later. Cultivars generally differed in chlorophyll leaf content. With many years of research conducted in our institute, it is clear that for most cultivars, SPAD readings gave 43 result in optimum yield (Trawczyński 2012). Chlorophyll fluorescence is a measure of the efficiency of photosynthetic apparatus. The maximum efficiency of photosynthesis occurs when the surface of the lamina has already reached the minimum size, then decreases with the increase of the lamina surface (Pula et al. 1999, Michalek and Sawicka 2005); they found that the operation of photosystem PS II is the most sensitive indicator of various stressors on plants. Also Rykaczewska and Mańkowski (2015) stated that chlorophyll fluorescence is a very sensitive probe of the physiological status of leaves, which can provide a very

rapid assessment of plant performance in a wide range of situations. Our study also confirmed the highest activity of the photosynthetic apparatus in the early stages of plant development in both production systems. In the organic system, the activity of the photosynthesis was smaller and just as in the case of chlorophyll content it fell faster than in the conventional system. The obtained relationship between the studied indicators and yield of tubers showed the highest correlation between the leaf area index and the final yield. This indicator seems to be the key, because good development of plants proves their good nourishment, and thus life processes running properly. The slightly higher correlation in the conventional system provides for greater stability of the plant growth in this production system. In the organic system, plants are exposed to a stronger adverse factors. Weather factors play a huge role in this case. In the years of unfavourable weather conditions, more significant losses in yield can be expected in the organic than conventional production systems (Zarzyńska and Pietraszko 2015, Zarzyńska and Goliszewki 2015).

To summarize, it is possible to predict the tuber yield based on physiological-morphological plant parameters; in the conventional system, it is simpler to eliminate the effects of factors independent of farmer.

REFERENCES

- Goffart J.P., Olivier M., Frankinet M. (2008): Potato crop nitrogen status assessment to improve N fertilization management and efficiency: Past-present-future. *Potato Research*, 51: 355–383.
- Ierna A. (2007): Characterization of potato genotypes by chlorophyll fluorescence during plant aging in Mediterranean environment. *Photosynthetica*, 45: 568–575.
- Jefferies R.A. (1992): Effects of drought on chlorophyll fluorescence in potato (*Solanum tuberosum* L.). I. Plant water status

doi: 10.17221/101/2017-PSE

- and the kinetics of chlorophyll fluorescence. *Potato Research*, 35: 25–34.
- Marscher H. (1995): *Mineral Nutrition for Higher Plants*. 2nd Ed. London, Academic Press.
- Mauromicale G., Ierna A., Marchese M. (2006): Chlorophyll fluorescence and chlorophyll content in field-grown potato as affected by nitrogen supply, genotype, and plant age. *Photosynthetica*, 44: 76–82.
- Michałek W., Sawicka B. (2005): Chlorophyll content and photosynthetic activity of medium-late varieties of potato in the conditions of cropland in east-central Poland. *Acta Agrophysica*, 6: 183–195.
- Milthrope F.L., Moorby J. (1997): *Introduction to Physiology Plant Yielding*. Warszawa, Agriculture and Forest Publishing.
- Pula J., Skrzypek E., Łabza T., Dubert T. (1999): Chlorophyll fluorescence as one of the indicators of potato yielding. Potatoes for trade and processing – Agronomic and storage conditions guaranteeing a quality. *Radzików*, 23–25: 110–122.
- Ramírez D.A., Yactayo W., Gutiérrez R., Mares V., De Mendiburu F., Posadas A., Quiroz R. (2014): Chlorophyll concentration in leaves is an indicator of potato tuber yield in water-shortage conditions. *Scientia Horticulturae*, 168: 202–209.
- Rykaczewska K., Mańkowski D. (2015): The effect of physiological age of potato plants on chosen chlorophyll fluorescence parameters. *Plant, Soil and Environment*, 61: 462–467.
- StatSoft, Inc. (2014): *Statistica* (data analysis software system), version 12. Available at www.statsoft.com
- Strasser R.J., Srivastava A., Tsimilli-Michael M. (2000): The fluorescence transient as a tool to characterize and screen photosynthetic samples. In: Yunus M., Pathre U., Mohanty P. (eds.): *Probing Photosynthesis: Mechanisms, Regulation and Adaptation*. London, Taylor and Francis, 445–483.
- Trawczyński C. (2012): Preparation of field and fertilization of potato. In: Chotkowski J. (ed.): *Production and Market of Potato*, Warsaw, 182–197. (In Polish)
- Tremblay N. (2004): Determining nitrogen requirements from crop characteristics. *Recent Research Development in Agronomy Horticulture*, 1: 157–182.
- Van Delen A. (2001): Yield and growth components of potato and wheat under organic nitrogen management. *Agronomy Journal*, 93: 1370–1385.
- Van der Zaag D.E. (1992): *Potatoes and their cultivation in the Netherlands*. Hague, Netherlands Potato Consultative Institute.
- Vos J. (1995): Nitrogen and the growth of potato crops. In: Haverkort A.J., MacKerron D.K.L. (eds): *Potato Ecology and Modeling of Crops under Conditions Limiting Growth*. Dordrecht, Kluwer, 115–128.
- Zarzyńska K., Szutkowska M. (2012): Development differences, yield and late blight development (*Phytophthora infestans*) infection of potato plants grown under organic and conventional systems. *Journal of Agriculture Science and Technology A*, 3/4: 281–290.
- Zarzyńska K., Goliszewski W. (2015): Cultivar – Environmental determinants of potato yielding under organic system. *Journal of Research and Applications in Agricultural Engineering*, 60: 135–139.
- Zarzyńska K., Pietraszko M. (2015): Influence of climatic conditions on development and yield of potato plants growing under organic and conventional systems in Poland. *American Journal of Potato Research*, 92: 511–517.

Received on February 20, 2017

Accepted on April 20, 2017

Published online on April 25, 2017