

Effect of moisture content on some physical properties of *Canarium schweinfurthii* Engl. fruits

J.C. EHIEM, V.I.O. NDIRIKA, U.N. ONWUKA

Department of Agricultural and Bio-Resources Engineering, Michael Okpara University of Agriculture, Umudike, Umuahia, Nigeria

Abstract

EHIEM J.C., NDIRIKA V.I.O., ONWUKA U.N. (2016): **Effect of moisture content on some physical properties of *Canarium schweinfurthii* Engl. fruits.** Res. Agr. Eng., 62: 162–169.

The effect of moisture content on some physical properties of three varieties of *Canarium schweinfurthii* Engl. fruits (small, large and long) was studied at 40.91%, 34.92%, 23.44%, 18.5% and 11.03% moisture content on wet basis in order to solve problems associated with designing and development of processing and handling equipment for these fruits. The physical parameters investigated were major, intermediate, minor diameters, geometric mean diameter, unit mass, volume, sphericity, density, bulk density, roundness, aspect ratio, porosity, surface and specific surface area. The results obtained showed that the physical parameters decreased linearly with a decrease in moisture content. Major diameter and true density of all the fruit varieties were not affected significantly ($P < 0.05$) by moisture content. *Canarium schweinfurthii* Engl. fruits are not round but spherical and oblong, hence, they can rather roll than slide. Among the studied varieties, small *Canarium schweinfurthii* Engl. is less bulky, has the highest specific surface area and is more porous.

Keywords: sphericity; area; density; porosity; variety

Canarium schweinfurthii Engl. tree is one of the economic tree crops that belong to the family of *Burseraceae* and popularly is called “ube mgba” in Igbo and “elemi” in Hausa land. The English name is African bush candle. It is grown widely in the equatorial forest regions of East, West and Central Africa (ORWA et al. 2009). In Nigeria, it is mostly grown in the southeast part of the country. The tree produces edible fruits that contain 20.43% crude protein, 23% crude fat, 0.75% crude fibre, 20.10% carbohydrate, 11.8% cellulose and 3.25% ash (ONIMAWO, ADUKWU 2003). The fruits are eaten fresh or boiled. EDOU et al. (2012) reported that the fruit pulp and kernel contain about 30% to 50% oil, which are used industrially for the manufacturing of shampoo and waxes and, pharmaceutically to

produce drugs for treatment of wounds and microbial infections. OLAWALE (2012) also reported that the extraction of oils from elemi pulp and kernel is not being carried out at commercial level at present, despite the availability of the fruit in large quantity in Nigeria and elsewhere in Sub-Saharan Africa. This situation would improve if data needed for the design and operation of the oils’ extraction plants were available. In handling and processing agricultural products, some fundamental information about their characteristics is essentially needed. This information can be obtained through the knowledge of engineering properties of the products, which constitutes essential engineering data in designing modern technology for their production, handling, processing, storage, preservation, quality

Supported by the Tertiary Education Trust Fund through Michael Okpara University of Agriculture, Umudike, Abia State, Nigeria.

evaluation, distribution and marketing and utilization. Engineering properties of agricultural product is profitably used for planting, harvesting, drying and storage. It improves working efficiency of processing equipment, reduces losses and waste of constructional materials. It also saves time and money, offer ways in which products can be utilized effectively and helps to maintain quality even in adverse storage and handling conditions. Literature reveals that several studies have been carried out on engineering properties of different agricultural products; millet (*Pennisetum glaucum* L.) (NDIRIKA, OYELEKE 2006), *Lablab purpureus* L. (SIMONYAN et al. 2009), *Jatropha curcas* L. fruit, nut and kernel (SIRISOMBOON et al. (2007), *Jathropa curcas* L. seed (KABUTEY et al. 2011). Despite all these studies and numerous economic values of *Canarium schweinfurthii* fruits, no work has been published on the post-harvest physical properties of *Canarium schweinfurthii* Engl. fruits. The aim of this study therefore was to investigate the effect of moisture content on some physical properties of *Canarium schweinfurthii* Engl. fruits, like principal dimensions, unit mass, volume, sphericity, density, bulk density, roundness, aspect ratio, porosity, surface and specific surface area which will be useful in designing and developing processing and handling equipment for this fruits.

MATERIAL AND METHODS

Three varieties of *Canarium schweinfurthii* Engl. fruits (small, large and long size (CSHT_S, CSHT_L and CSHT_{LG}, respectively)) used for this investigation were sourced from Ebonyi (6°15'N 8°05'E) state of Nigeria. They were cleaned and initial moisture content of the fruits was attained by keeping the fruits in refrigerator at 0°C for 24 hours. Other moisture levels were attained by exposing the fruits to environmental temperature and measurement taken at two-days interval. Moisture content was determined as:

$$MC_{wb} = \frac{w_1 - w_2}{w_1} \times 100 \quad (1)$$

where:

MC_{wb} – moisture content wet basis; w_1 , w_2 – initial weight and final weight, respectively

For each of the moisture contents obtained, the principal dimensions (major (a), intermediate (b) and

minor diameters (c)) of 50 samples were measured using the digital vernier caliper (JIS7502; Mitutoyo, Aurora, USA) of 0.01mm accuracy while the weight of individual fruit was measured using the electronic weighing balance (FEJ-300; Fuzhou Furi Electronic, Ltd., Fuzhou, China) with sensitivity of 0.01g. Geometric mean diameter (GMD), sphericity, arithmetic mean diameter (AMD) and aspect ratio were calculated as shown in Eqs (2–5), NDIRIKA and OYELEKE (2006); EHIEM and SIMONYAN (2012):

$$GMD = \sqrt[3]{abc} \quad (2)$$

$$Sphericity = GMD/a \quad (3)$$

$$AMD = (a + b + c)/3 \quad (4)$$

$$Aspect\ ratio = b/a \quad (5)$$

Roundness was obtained by tracing each fruit in its rest position on a transparent paper using a sharp pencil. The radius of the sharpest corner and the circumscribed circle were measured and roundness estimated as:

$$R = A_p/A_c \quad (6)$$

where:

R – roundness; A_p – area of the sharpest corner (mm²); A_c – area of the circumscribed circle (mm²)

Water displacement method was used to estimate the volume of the fruits. The density (r) of each fruit was calculated using the Eq. (7):

$$r = mass/volume \text{ (g/cm}^3\text{)} \quad (7)$$

Bulk density was taken as the ratio of mass of samples to the volume of container it occupies. It was determined by filling to the brim a 2,580 cm³ container with the sample and level off the excess samples with a flat object. The whole sample was weighed and bulk density (ρ_b) and porosity calculated with Eq. (8) and Eq. (9), EKE et al. (2007):

$$\rho_b = M_p/V_c \quad (8)$$

where:

M_p – mass of sample (g); V_c – volume of the container (cm³)

$$Porosity = 1 - \frac{\rho_b}{\rho} \times 100 \quad (9)$$

Surface area (mm²) and specific surface area (mm²/cm³) of the fruits were estimated with formu-

doi: 10.17221/11/2015-RAE

as shown in Eq. (10) and Eq. (11), SIRISOMBOON et al. (2007):

$$A_s = \pi D^2 \quad (10)$$

$$A_{ss} = \frac{A_s \rho b}{mass} \quad (11)$$

where:

A_s – surface area (mm²); A_{ss} – specific surface area (mm²/cm³); D^2 – geometric mean diameter

The data obtained from the experiment were analysed using Excel and Genstat (Genstat discovery Ed. 3; United Kingdom.) statistical packages.

RESULTS AND DISCUSSION

Principal dimension and size distributions

Major, intermediate and minor diameters of *Canarium schweinfurthii* Engl. fruits for all the varieties evaluated decreased as moisture content decreased from 40.91% to 11.03% w.b. (Table 1). Similar trend was reported by NDIRIKA and OYELEKE (2006), SIMONYAN et al. (2009), RAZAVI et al. (2009), FATHOLHZADEH et al. (2008) for millet (*Pennisetum glaucum* L.), *Lablab purpureus*, canola seeds and barberry fruits respectively. Major, intermedi-

Table 1. Measured parameters and coefficient of variation of *Canarium schweinfurthii* Engl. fruits at different moisture content

Samples	Moisture content (% w.b.)	Major diameter (a, mm)	Intermediate diameter (b, mm)	Minor diameter (c, mm)	Mass (g)	Volume (cm ³)	Density (g/cm ³)
CSHT _s	40.91	26.06 ± 1.09 (4.18)	16.69 ± 0.59 (3.52)	16.17 ± 0.27 (1.98)	4.87 ± 0.26 (5.71)	3.69 ± 0.52 (2.94)	1.36 ± 0.37 (2.71)
	34.92	25.08 ± 1.37 (5.47)	15.89 ± 0.56 (3.56)	15.36 ± 1.54 (1.05)	3.99 ± 0.38 (4.42)	3.22 ± 0.46 (1.49)	1.23 ± 0.32 (2.58)
	23.44	24.20 ± 2.08 (8.62)	15.05 ± 0.78 (5.24)	14.57 ± 1.54 (1.05)	3.46 ± 0.39 (2.5)	2.8 ± 0.51 (1.81)	1.23 ± 0.35 (2.82)
	18.5	23.87 ± 2.24 (9.39)	14.45 ± 0.84 (5.8)	13.94 ± 1.59 (1.14)	3.18 ± 0.38 (1.2)	2.54 ± 0.49 (1.95)	1.26 ± 0.42 (3.36)
	11.03	23.05 ± 2.73 (1.18)	13.99 ± 0.96 (1.20)	13.48 ± 1.62 (1.2)	2.74 ± 0.48 (1.75)	2.31 ± 0.52 (2.26)	1.19 ± 0.42 (3.54)
CSHT _L	40.91	35.14 ± 0.95 (2.59)	22.44 ± 0.62 (1.72)	22.03 ± 0.61 (1.71)	11.37 ± 0.82 (2.84)	9.11 ± 0.65 (5.01)	1.25 ± 0.04 (0.14)
	34.92	34.65 ± 0.89 (2.58)	21.74 ± 0.62 (2.84)	21.35 ± 0.59 (2.75)	10.33 ± 0.78 (3.48)	8.43 ± 0.62 (2.38)	1.23 ± 0.04 (3.62)
	23.44	34.42 ± 0.89 (2.59)	21.05 ± 0.51 (2.44)	20.76 ± 0.55 (2.67)	9.18 ± 0.77 (3.85)	7.89 ± 0.55 (7.01)	1.16 ± 0.05 (4.43)
	18.5	34.09 ± 0.82 (2.33)	20.25 ± 0.57 (2.76)	20.41 ± 0.57 (2.81)	8.51 ± 0.67 (4.71)	7.58 ± 0.55 (6.85)	1.12 ± 0.059 (5.24)
	11.03	34.09 ± 0.80 (2.35)	20.25 ± 0.53 (2.63)	19.92 ± 0.52 (2.61)	8.04 ± 0.62 (3.74)	7.21 ± 0.49 (6.74)	1.12 ± 0.053 (4.73)
CSHT _{LG}	40.91	42.86 ± 0.91 (2.13)	20.05 ± 0.66 (3.27)	19.56 ± 0.65 (3.33)	9.68 ± 0.66 (4.81)	7.89 ± 0.59 (7.45)	1.226 ± 0.03 (2.39)
	34.92	42.59 ± 0.88 (2.09)	18.54 ± 0.63 (3.41)	18.06 ± 0.64 (3.56)	8.95 ± 0.49 (2.43)	7.13 ± 0.54 (7.55)	1.16 ± 0.17 (14.26)
	23.44	42.36 ± 0.93 (2.09)	18.44 ± 0.51 (2.77)	17.89 ± 0.53 (2.97)	7.09 ± 1.23 (1.73)	7.15 ± 1.07 (2.91)	1.144 ± 1.15 (1)
	18.5	42.02 ± 0.97 (2.30)	17.39 ± 2.38 (1.36)	17.18 ± 0.98 (3.74)	6.68 ± 1.25 (1.87)	6.59 ± 1.04 (2.82)	1.10 ± 1.80 (2.78)
	11.03	41.71 ± 1.36 (3.27)	16.90 ± 2.43 (1.44)	16.58 ± 1.13 (3.82)	6.22 ± 1.34 (2.16)	6.14 ± 1.07 (2.50)	1.06 ± 3.28 (2.27)

CSHT_s – *Canarium schweinfurthii* Engl. small; CSHT_L – *Canarium schweinfurthii* Engl. long; CSHT_{LG} – *Canarium schweinfurthii* Engl. large; values in brackets are the coefficient of variation

Table 2. ANOVA summary of physical parameters of *Canarium schweinfurthii* fruit studied at moisture content range of 40.91–11.03%

Source of variation	Df	<i>a</i>	<i>b</i>	<i>c</i>	Mass (g)	Sphericity	Vol.	Density	S. area	Roundness	<i>P</i>	
		(mm)									0.05	0.01
Samples	2	790.50**	976.50**	1,084.9**	31.52**	648.74**	1,678.39**	1.35 ^{ns}	920.72**	20.62°	4.46	8.65
Moisture content	4	2.62 ^{ns}	7.14°	68.02**	314.29**	13.15°	57.53	0.29 ^{ns}	38.69**	5.23°	3.84	7.01
Interaction	8	5.56°	43.40**	12.22°	68.79**	36.12**	137.26	8.10°	65.53**	8.10°	2.266	3.17
Error	8											

*significant; **highly significant; ns – not significant; Df – degree of freedom; *a* – major diameter; *b* – intermediate diameter; *c* – minor diameter; S. area – surface area; 50 samples; 5% level of significance

ate and minor diameters of CSHT_S, CSHT_L, and CSHT_{LG} fruits decreased by 11.5%, 3.22%, and 2.68%; 16.18%, 9.76% and 15.771%; 16.63%, 9.58% and 15.24%, respectively, as moisture reduced from 40.91 to 11.03% w.b. All the principal dimensions had low coefficient of variation, which indicates evenly distributed values about their means. Grains of chickpea (KONAK et al. 2002) and Samaru sorghum 17 (SIMONYAN et al. 2007) also behaved the same way at 16.52% d.b., 6.5% d.b., 16.5–8.89% d.b., and 13.5% w.b., respectively. The ANOVA shown in Table 2 also revealed that moisture content had no significant (5%) effect on the major diameter of all the varieties studied. Regression equations of the principal dimensions in Table 3 showed high values of coefficient of determination (R^2) indicating good relationship between the parameters evaluated.

Mass and volume

The mean values of mass and volume are summarized in Table 1. Mass and volume decreased by 43.74% and 29.29%; 36.34% and 17.87%; 27.56% and 14.51% for CSHT_S, CSHT_L and CSHT_{LG} respectively as moisture content decreased from 40.91 to 11.03% w.b. FATHOLLAHZADEH et al. (2008), AKINOSO and RAJI (2010) observed 70.77% and 57.82%; 9.71% and 8.89% decrease in mass and volume of barberry fruits and tenera, respectively as moisture content decreased from 89.19–12.64% and 11–5% w.b. Low coefficient of variation of mass and volume showed a uniform dispersion of their values about their means. Both physical properties are highly influenced by moisture content and variety at 5% and 1% levels of significance as shown in

Table 3. Equations representing relationship between physical parameters and moisture content

Parameters	CSHT _S		CSHT _L		CSHT _{LG}	
	regression equation	R^2	regression equation	R^2	regression equation	R^2
<i>c</i>	$0.062h^2 - 1.095h + 27.05$	0.987	$0.055h^2 - 0.592h + 35.66$	0.993	$0.130h^2 - 0.865h + 43.07$	0.906
<i>b</i>	$0.065h^2 - 1.074h + 17.72$	0.998	$0.044h^2 - 0.781h + 22.75$	0.996	$-1.89\ln(h) + 20.01$	0.989
<i>c</i>	$0.062h^2 - 1.053h + 17.17$	0.999	$-0.517h + 22.44$	0.986	$-1.73\ln(h) + 19.50$	0.975
Mass	$0.091h^2 - 1.068h + 5.823$	0.997	$0.115h^2 - 1.541h + 12.84$	0.997	$0.142h^2 - 1.771h + 11.47$	0.960
Sphericity	$0.000h^2 - 0.011h + 0.745$	0.907	$0.745e - 0.01h$	0.987	$0.001h^2 - 0.021h + 0.620$	0.973
Roundness	$0.009h^2 - 0.106h + 0.536$	0.977	$-0.01\ln(h) + 0.505$	0.984	$0.007h^2 - 0.090h + 0.582$	0.954
Volume	$0.045h + 1.75$	0.907	$0.019h + 6.924$	0.986	$0.069h + 5.468$	0.879
Density	$0.009h^2 - 0.095h + 1.437$	0.958	$0.022h + 0.795$	0.993	$0.001h^2 - 0.057h + 1.991$	0.986
Surfaces area	$0.001h^2 + 0.044h + 7.785$	0.991	$0.001h^2 + 0.039h + 17.51$	0.987	$0.002h^2 - 0.019h + 16.22$	0.844
Bulk density	$-0.016h^2 + 2.494h - 1.054$	0.997	$-0.005h^2 + 0.638h + 47.02$	0.983	$0.006h^2 + 0.409h + 41.72$	0.987
Porosity	$0.023h^2 - 2.406h + 104.0$	0.985	$0.007h^2 - 0.485h + 56.33$	0.925	$0.018h^2 - 1.298h + 65.61$	0.942

CSHT_S – *Canarium schweinfurthii* Engl. small; CSHT_L – *Canarium schweinfurthii* Engl. long; CSHT_{LG} – *Canarium schweinfurthii* Engl. large; *h* – moisture content; *e* – mean exponential expression; *a* – major diameter; *b* – intermediate diameter; *c* – minor diameter

doi: 10.17221/11/2015-RAE

Table 4. Mass and volume were linearly regressed with moisture content with high values of R^2 . The regression equations are shown in Table 3.

Sphericity, roundness and aspect ratio

The influence of moisture content on aspect ratio, sphericity and roundness of different varieties of *Canarium schweinfurthii* Engl. fruits are also presented in Table 4 and Figs 1 and 2. The values of the above physical parameters decreased with reduc-

tion in moisture content. Aspect ratio, sphericity and roundness of CSHT_S, CSHT_L and CSHT_{LG} were reduced by 5.27%, 3.80% and 45.10%; 4.65%, 6.98% and 4.62%; and, 9.54%, 13.30% and 35.80%, respectively as moisture content decreased from 40.91% to 11.03% w.b. FATHOLLAHZADEH et al. (2008), RAZAVI et al. (2009) and BERYDEH and MANGOPE (2002) also reported 40.43%, 2.29%, 18.5% and 0.475% reduction in sphericity of barberry fruits, canola seeds (SLM spp.) and pigeon pea as moisture content reduced from 89.23 to 11.64% w.b., 21 to 6% and at 25%, respectively. At 40.91% w.b., the

Table 4. Calculated parameters and coefficient of variation of *Canarium schweinfurthii* Engl. fruits at different moisture content

Samples	Moisture content (% w.b.)	GMD	AMD	SQMD	EQD	Surface area (mm ²)	Specific surface area (cm ² /cm ³)	Aspect ratio
CSHT _S	40.91	19.13 ± 1.09 (5.09)	19.64 ± 0.92 (4.67)	19.37 ± 0.98 (5.05)	19.38 ± 0.99 (5.10)	11.54 ± 1.18 (1.09)	1.74 ± 0.18 (1.09)	64.09 ± 2.14 (3.34)
	34.92	18.26 ± 1.06 (3.81)	18.78 ± 0.63 (3.33)	18.49 ± 0.96 (3.18)	18.51 ± 0.83 (4.16)	1,051 ± 1.10 (1.04)	1.77 ± 0.18 (1.04)	63.52 ± 2.97 (4.68)
	23.44	17.41 ± 1.25 (4.17)	17.94 ± 0.80 (4.47)	17.65 ± 1.18 (4.39)	17.67 ± 1.00 (4.68)	9.57 ± 1.25 (1.31)	1.31 ± 0.17 (1.31)	62.49 ± 4.27 (4.83)
	18.5	16.84 ± 1.34 (4.92)	17.42 ± 0.87 (4.98)	17.10 ± 1.26 (4.36)	17.12 ± 1.08 (4.28)	8.96 ± 1.28 (1.42)	1.18 ± 0.17 (1.42)	60.86 ± 4.32 (0.09)
	11.03	16.27 ± 1.47 (3.04)	16.84 ± 1.41 (3.62)	16.53 ± 1.43 (3.65)	16.55 ± 1.26 (3.62)	8.38 ± 1.38 (1.64)	0.74 ± 0.122 (1.64)	60.71 ± 4.88 (4.73)
CSHT _L	40.91	25.90 ± 0.644 (1.61)	26.54 ± 0.66 (1.62)	26.19 ± 0.65 (1.61)	26.21 ± 0.65 (1.61)	21.09 ± 1.04 (0.05)	1.117 ± 0.059 (0.05)	63.87 ± 1.51 (0.036)
	34.92	25.23 ± 0.63 (2.48)	25.91 ± 0.63 (2.44)	24.54 ± 0.63 (2.47)	25.56 ± 0.63 (2.46)	20.03 ± 9.9 (4.94)	1.21 ± 0.06 (4.94)	62.76 ± 1.37 (2.21)
	23.44	24.69 ± 0.57 (2.33)	25.41 ± 0.59 (2.31)	25.01 ± 0.58 (2.31)	25.04 ± 2.32 (2.32)	19.16 ± 0.89 (4.67)	1.24 ± 0.057 (4.67)	61.17 ± 1.29 (2.12)
	18.5	24.36 ± 0.55 (2.25)	25.11 ± 0.55 (2.18)	24.69 ± 0.55 (2.21)	24.72 ± 0.55 (2.21)	18.65 ± 0.84 (4.54)	1.22 ± 0.056 (4.54)	60.62 ± 1.60 (2.63)
	11.03	23.95 ± 0.54 (2.24)	24.75 ± 0.54 (2.17)	24.31 ± 0.54 (2.20)	24.33 ± 0.54 (2.19)	18.03 ± 0.808 (4.48)	1.20 ± 0.05 (4.48)	59.41 ± 1.36 (2.29)
CSHT _{LG}	40.91	25.61 ± 0.61 (2.37)	27.49 ± 0.57 (2.08)	26.39 ± 0.59 (2.24)	26.49 ± 0.58 (2.21)	20.62 ± 0.99 (4.32)	1.46 ± 0.0707 (4.32)	46.78 ± 1.67 (3.57)
	34.92	23.86 ± 57 (2.41)	25.73 ± 0.53 (2.07)	24.63 ± 0.56 (2.25)	24.74 ± 0.55 (2.23)	17.88 ± 0.89 (5)	1.419 ± 0.079 (5)	45.23 ± 1.43 (3.16)
	23.44	23.78 ± 1.87 (7.89)	26.08 ± 0.95 (3.66)	24.75 ± 1.25 (5.01)	24.87 ± 1.35 (5.40)	17.88 ± 2.18 (1.22)	1.36 ± 0.17 (1.22)	43.47 ± 1.43 (3.29)
	18.5	23.12 ± 2.07 (8.95)	25.55 ± 1.007 (3.94)	24.14 ± 1.33 (5.51)	24.27 ± 1.46 (5.99)	16.92 ± 2.24 1.27	1.29 ± 0.17 1.27	41.36 ± 5.72 (3.38)
	11.03	22.53 ± 2.29 (1.14)	25.07 ± 1.10 (4.4)	23.61 ± 1.42 (6)	23.74 ± 1.58 (6.6)	16.11 ± 2.36 (1.69)	1.23 ± 0.18 (1.69)	40.56 ± 5.94 (3.02)

CSHT_S – *Canarium schweinfurthii* Engl. small; CSHT_L – *Canarium schweinfurthii* Engl. long; CSHT_{LG} – *Canarium schweinfurthii* Engl. large; GMD – geometric mean diameter; AMD – arithmetic mean diameter; SMD – square mean diameter; EQD – equivalent mean diameter




Fig. 1. Influence of moisture content on (a) sphericity and (b) roundness of *Canarium schweinfurthii* Engl. varieties

values of aspect ratio and sphericity of all the varieties were high except the aspect ratio of CSHT_{LG}, showing that at higher moisture content, the fruits are spherical and oblong; hence, they can rather roll than slide. Moisture content affected ($P < 0.05$) the sphericity and roundness of CSHT_{LG}. It could be because the structure of CSHT_{LG} fruit has more pore spaces than other varieties, which resulted in high shrinkage effect during moisture loss. Roundness had low values for all the varieties studied, showing that *Canarium schweinfurthii* Engl. fruits are not round, therefore, the shape of screen opening for separation purposes must not be round. Regression equations are shown in Table 3.

True density, bulk density and porosity

The results of true density, bulk density and porosity as influenced by varied moisture content are




Fig. 2. Influence of moisture content on (a) bulk density and (b) porosity of *Canarium schweinfurthii* Engl. varieties

shown in Table 1 and Figs 3 and 4 respectively. From the table, it was observed that true density of all the varieties are higher than that of water indicating that the fruits cannot float in water. Hence, separation from other dense materials cannot take place in water. True density of CSHT_S decreased with decreased moisture content and had a polynomial relationship with moisture content. SIMONYAN et al. (2009) reported the same behaviour for *Lablab purpureus*, indicating that agricultural materials are not homogeneously formed. Bulk density and porosity also decreased as moisture content decreased. CSHT_S had the least (24.31%) and highest (79.65%) bulk density and porosity, respectively. This means that CSHT_S require lesser space per unit mass and will encourage easier aeration and water vapour diffusion during drying than other varieties. Similar results were observed for apricot kernel, cape fruits, two varieties of corn (Sc 704 and 370) and cowpea (IAR 339-1) by FATHOLLAHZADEH et al.

doi: 10.17221/11/2015-RAE

(2008), SESSIZ et al. (2007), SEIFI and ALIMARDANI (2010) and DAVIES and ZIBOKERE (2011), respectively. True density and bulk density of individual varieties studied had a positive linear correlation against moisture content showing a cordial relationship between the parameters. A similar positive correlation was reported on sunflower seed by BHISE et al. (2013). Negative correlations were also observed by BAUMLER et al. (2006), COSKUN et al. (2005) and DURSUN and DURSUN (2006) for safflower seed, sweet corn seed and caper seeds respectively. The ANOVA in Table 4 revealed that moisture content and variety had no significant effect on true density at 1% and 5% levels of significance.

Surface area and specific surface area

In the range of moisture content (40.91–11.03% w.b.) evaluated, surface area and specific surface area of CSHT_S, CSHT_L and CSHT_{LG} decreased from 11.54–8.38 mm² and 1.74–0.74 mm²/cm³; 21.09–18.03 mm² and 1.117–1.20 mm²/cm³; 20.62–16.11 mm² and 1.46–1.23 mm²/cm³ respectively (Table 2). The table also revealed that CSHT_L and CSHT_S had the highest surface area (21.09 ± 1.04) and specific surface area (1.74 ± 0.18), meaning that heat and mass transfer in CSHT_L and CSHT_S fruits are faster than in CSHT_{LG}. Hence, the prediction of their drying rate can easily be carried out. The coefficient of variation of both surface areas is low showing uniform dispersion about their means. Similar result was reported for *Jatropha curcas* fruits, nuts and kernel and, REDSPAR and DELBARSTIVAL apples by SIRISOMBOON et al. (2007) and KHEIRALIPOUR et al. (2008) respectively. Moisture content, variety and their interactions affected surface area of *Canarium schweinfurthii* Engl. fruits significantly at 1% and 5% levels of significance (Table 4).

CONCLUSION

The study of some physical properties of *Canarium schweinfurthii* Engl. fruits as influenced by moisture content revealed the following:

- All the physical parameters evaluated varied linearly from variety to variety with a decrease in moisture content. The varieties of *Canarium*

schweinfurthii Engl. fruits differ significantly (1% and 5%) from each other both in shape and size

- Moisture content does not significantly (1% and 5%) affect major diameter and true density of *Canarium schweinfurthii* Engl. fruits. The fruits cannot be separated by float method since they are denser than water. CSHT_S variety is less bulky, has the highest specific surface area and is more porous than other varieties.
- *Canarium schweinfurthii* Engl. fruits of the three varieties are not round but spherical and oblong, hence they can rather roll than slide.

References

- Akinoso R., Raji A.O. (2011): Physical properties of fruit, nut and kernel of oil palm. *International Agrophysics*, 25: 85–88.
- Baumler E., Cuniberti A., Nolasco S.M., Riccobene I.C. (2006): Moisture dependent physical and compression properties of safflower seed. *Journal of Food Engineering*, 72: 134–140.
- Baryeh E.A., Mangope B.K. (2002): Some physical properties of Qp-38 variety pigeon pea. *Journal of Food Engineering*, 56: 59–65.
- Bhise S., Kaur A., Manikantan M.R. (2013): Moisture dependant physical properties of sunflower seed (PSH 569). *International Journal of Engineering and Science*, 2: 23–27.
- Coskun M. B., Yalçın I., Özarslan C. (2005): physical properties of sweet corn seed (*Zea mays saccharata* Sturt.). *Journal of Food Engineering*, 74: 523–528.
- Davies R.M., Zibokere D.S. (2011). Effect of moisture content on some physical and mechanical properties of three varieties of cowpea (*Vigna unguiculata* (L.) walp). *Agricultural Engineering International: CIGR Journal*, 13.
- Dursun E., Dursun I. (2005): Some physical properties of caper seed. *Biosystems Engineering*, 92: 237–245.
- Edou E.P., Abdoul-Latif F.M., Obame L.C., Mewono L., Agnani H. (2012): Volatile constituents of *Canarium schweinfurthii* Engl. essential oil from Gabon. *International Journal of AgriScience*, 2: 200–203.
- Ehiem J.C., Simonyan K.J. (2012): Physical properties of wild mango (*Irvingia* spp.) fruits and nuts. *International Agrophysics*, 26: 95–98.
- Eke C.N.U., Asoegwu S.N., Nwandikom G.I. (2007): The Physical properties of Jackbean seed (*Canavalia ensiformis*). *Agricultural Engineering International: The CIGR E-Journal Manuscript FP 07 014 Vol. 4*.
- Fathollahzadeh H., Mobli H., Jafari A., Rajabipour A., Ahmadi H., Borghai A.M. (2008): Effect of moisture content on some physical properties of barberry. *American-Eurasian*

- Journal of Agriculture and Environmental Science, 3: 789–794.
- Kabutey A., Herák D., Sedláček A. (2011): Behaviour of different moisture contents of *Jatropha curcas* L. seeds under compressive loading. Research Agricultural Engineering, 57: 72–77.
- Kheiralipour K., Tabatabaefar A., Mobli H., Rafiee S., Sahra-roo A., Rajabipour A., Jafari A. (2008): Some physical properties of apple. Pakistan Journal of Nutrition, 7: 667–672.
- Konak M., Carman K., Aydin C. (2002): Physical properties of chick pea seeds. Biosystem Engineering, 182: 73–78.
- Masoumi A.A., Rajabipour A., Tabil L.G., Akrami A.A. (2006): Physical attributes of garlic (*Allium sativum* L.). Journal of Agricultural Science. Technology, 8: 15–23.
- Naderiboldaji M., Khadivikhub, A., Tabatabaefar A., Varnamkhasti M.G., Zamani Z. (2008): Some physical properties of sweet cherry (*Prunus avium* L.) fruits. American-Eurasian Journal of Agricultural & Environmental Science, 3: 513–520.
- Ndirika V.I.O., Oyeleke O.O. (2006): Determination of some selected physical properties and their relationship with moisture content for millet (*PennisetumGlaucum* L.). Applied Engineering in Agriculture, ASABE, 22: 2.
- Olawale A.S. (2012): Solid-liquid extraction of oils of African elemi's fruit. Agricultural Engineering International: CIGR E-Journal, 14: 159–160.
- Onimawo I.A., Adukwu B.I. (2003): Proximate composition and selected physicochemical properties of the fruits of *Canarium schweifurthii* Linn. African Journal of Science and Technology, 4: 1–4.
- Orwa C., Mutua A., Kindt R., Jamnadass R., Simons A. (2009). Agroforestry database: a tree reference and selection guide version 4.0. Available at <http://www.worldagroforestry.org/af/treedb/>
- Razavi S.M.A., Yeganehzad S., Sadeghi A. (2009): Moisture dependent physical properties of canola seeds. Journal Agricultural Science Technology, 11: 309–322.
- Seifi M.R., Alimardani R. (2010): Comparison of moisture dependent physical and mechanical properties of two varieties of corn (Sc 704 and Dc 370). Australian Journal of Agricultural Engineering, 1: 170–178.
- Sessiz A., Esgici R., Kizil S. (2007): Moisture dependent physical properties of caper (*Capparispp.*) fruit. Journal Food Engineering, 79: 1426–1431.
- Simonyan K.J., Yiljep Y.D., Oyatoyan O.B., Bawa G.S. (2009): Effect of moisture on some physical properties of *Lablab purpureus* (L.) sweet seeds. Agricultural Engineering International: CIGR Journal Manuscript 1279. Vol. XI.
- Sirisomboon P., Potncheloampom P., Romphophek T. (2007): Physical properties of greensoya bean: Citrus for sorting. Journal of Food Engineering, 79: 18–22.

Received for publication February 3, 2015

Accepted after corrections May 11, 2015

Corresponding author:

Engr. EHIEM JAMES CHINAKA, University of Agriculture, College of Engineering and Engineering Technology, Department of Agricultural and Bio-Resources Engineering, Umudike, P.M.B. 7267, Umuahia, Abia State, Nigeria; e-mail: chinaka71@yahoo.com
