

The use of inflammatory markers as a prognostic aid for traumatic reticuloperitonitis in water buffalo (*Bubalus bubalis*)

M. EL-ASHKER¹, M. SALAMA¹, A. RIZK¹, M. EL-BOSHY^{1,2}

¹Faculty of Veterinary Medicine, Mansoura University, Mansoura, Egypt

²Faculty of Applied Medical Science, Umm Al-Qura University, Makkah, Kingdom of Saudi Arabia

ABSTRACT: The present study was conducted to evaluate the prognostic significance of selected inflammatory markers for prediction of clinical outcomes of traumatic reticuloperitonitis (TRP) in water buffalo (*Bubalus bubalis*). Acute local TRP was initially diagnosed in 32 buffalo by clinical examination and confirmed by ultrasonography (USG), laparo-rumenotomy and/or necropsy findings in non-surviving cases. Ten clinically healthy buffalo were randomly selected and served as controls. Blood was drawn from all examined buffalo to measure the respective levels of tumor necrosis factor alpha (TNF- α), interleukin (IL)-1 β , IL-6, IL-10, interferon gamma (INF)- γ , serum amyloid A (SAA), haptoglobin (Hp), fibrinogen (Fb), C-reactive protein (CRP) and serum sialic acid (SSA). Clinically, the heart rates, but neither respiratory rate nor rectal temperature, were significantly higher in non-survivors compared with survivors ($P < 0.05$). In addition, the non-surviving buffalo were more likely to have anorexia and weakness compared with survivors. However, rumen stasis, recurrent ruminal tympany, lacrimation, lordosis, bruxism, and decreased milk production were commonly observed in all diseased animals. Biochemically, TNF- α , IL-1 β , IL-6, IL-10, SAA, Hp, Fb, CRP, and SSA levels were significantly higher in diseased buffalo compared with controls, and were higher in non-survivors than survivors ($P < 0.05$). The data herein indicate an ongoing cascade of systemic inflammatory responses in buffalo with TRP with concomitant compensatory anti-inflammatory reactions and the overall degree of cytokine network disruption may be an important prognostic indicator. Medical strategies to modulate inflammation must take into account the complex of cytokine biology in buffalo with TRP.

Keywords: cytokine profile; prognosis; traumatic reticulo-peritonitis; water buffalo

Traumatic reticuloperitonitis (TRP) is one of the most frequently occurring digestive diseases of bovines, which has drawn the attention of animal health professionals over the past years (Mousavi et al. 2007). It is caused by a combination of a lapse in good management and the non-selective eating habits of large ruminants. The buffalo is considered to be at higher risk of TRP than cattle (Misk et al. 2001). It has been suggested that TRP should be considered in any bovine animal with indigestion (Radostits et al. 2007); however, the variability of the clinical settings represents a significant challenge. The overall clinical symptoms of TRP are anorexia, recurrent tympany, fever, tachypnoea, and lordosis with abducted elbows (Saleh et al. 2008; Abdelaal et al. 2009). However, the observed signs are dependent upon the site of reticular perforation and

lesions caused by the foreign body, as visceral and muscular contractions promote migration within the body. Due to the complexity of TRP and possible coexistence with other syndromes, diagnosis is often difficult (Jafarzadeh et al. 2004). In practice, an initial diagnosis of TRP is mainly based upon clinical signs and laboratory findings, while it is confirmed by radiography and USG (Saleh et al. 2008; Abdelaal et al. 2009; Aref and Abdel-Hakiem 2013).

Systemic inflammation is a normal response to altered homeostasis and has an important role in several pathophysiological processes, such as infection or trauma. It is characterised by the endocrine release of different cytokines normally confined to paracrine regulation of a local inflammatory response (Koj 1997). These cytokines stimulate the release of acute phase proteins (APPs) from the

liver (Yoshioka et al. 2002). The initiation of the acute phase response starts when the animal is subjected to different internal or external stimuli (Johnson 1997). APPs have been widely used as a diagnostic tool for several human diseases, and recently in different animal species. In bovine species, the most commonly studied APPs include Hp, SAA, CRP, Fb, ceruloplasmin, alpha 1-antitrypsin and alpha 1-acid glycoprotein (Morimatsu et al. 1989; Jafarzadeh et al. 2004; Orro et al. 2011; El-Deeb and Iacob 2012). Although APPs constitute an element of the non-specific innate immune response which aims to restore homeostasis, their serum levels could reflect the degree of tissue damage in the diseased animal (Murata et al. 2004). Until recently, many clinicolaboratory advancements have been conducted for the diagnosis of TRP in water buffalo (Saleh et al. 2008; Abdelaal et al. 2009; Abouelnasr et al. 2012; Aref and Abdel-Hakim 2013; Abouelnasr et al. 2013; El-Ashker et al. 2013). However, the prognostic significance of immunological markers in buffalo with TRP has not been previously addressed. Therefore, this study aimed at evaluating the prognostic significance of selected inflammatory markers to predict clinical outcome in buffalo with TRP.

MATERIAL AND METHODS

Animal population and clinical examination. The study was conducted on thirty-two water buffalo cows (*Bubalus bubalis*). Twenty-four were within six weeks *post partum* and eight animals were in their final gestation period; their ages ranged from five to seven years. Ten clinically healthy peri-parturient buffalo cows, at three to five years of age, were randomly selected and served as controls. The clinical examination of all animals was carried out at the Veterinary Teaching Hospital, Faculty of Veterinary Medicine, Mansoura University, Egypt during 2012 and 2013. All examined buffalo were subjected to thorough clinical examination, in which all cardinal signs were recorded. Pain tests were also performed, including abdominal palpation caudal to the xiphoid, pinching of the withers, by placing a rod underneath the abdomen to elicit a grunt, and by using a foreign body detector that was applied over the ventral aspect of the abdomen to detect ferromagnetic foreign bodies. The diseased buffalo exhibited clinical signs of disturbed appetite, recurrent ruminal tympany, decreased defecation and milk yield, and mild pyrexia or eu-

thermia. Acute local TRP was initially diagnosed on the basis of clinical examination, and confirmed by USG, laparo-rumenotomy, and/or necropsy of non-surviving cases. Cases of chronic local TRP or diffuse peritonitis were not diagnosed during this investigation. The selection criteria were applied to the examined buffalo, excluding those with concomitant ailments and thereby removing potential confounding variables ($n = 6$). Once the diagnosis of acute local TRP had been made, medical and surgical management were subsequently offered to the owners. Conservative medical management included the immobilisation of the diseased animal, oral administration of a magnet, and intramuscular injection of broad spectrum antibacterial agents (penicillin and streptomycin twice daily at a dose of 22 000 IU/kg b.w. and 25 mg/kg b.w., respectively), concurrently with *i.v.* injection of flunixin meglumine at a dose of 2.2mg/kg b.w. once daily. Rehydrating *i.v.* fluids were also administered on an individual basis. In cases where an improvement in clinical signs was not observed within three days of the prescribed treatment, a laparo-rumenotomy was performed. Follow-up information was obtained via clinical examination, owner communication, and referring veterinarians with respect to the health status of animals following intervention. According to the clinical outcome, the diseased buffalo were allocated into two groups, survivors ($n = 12$), and non-survivors ($n = 20$).

Blood sampling and measurements. Blood was collected from all investigated buffalo by jugular vein puncture. The collected samples were added to heparinised tubes and tubes without anti-coagulant to yield plasma and serum, respectively. Blood plasma and serum samples were separated by centrifugation at 3000 g for 10 min and kept frozen at -20°C until required. Plasma samples were used to determine SAA, Hp, Fb and CRP by using commercially available bovine kits, according to Orro et al. (2011) (Kamiya Biomedical Company, USA) and an automated microplate reader (Bio TEC, ELX800G, USA) according to the manufacturer's instructions. Bovine cytokine kits were used for estimation of serum TNF- α , IL-6 (Kamiya Biomedical Company, USA), IL-1 β (Thermo Scientific Pierce Protein Biology Products-Rockford, Illinois), IL-10 (Genorise Scientific Inc., Philadelphia, USA), and IF- γ (BioSource Europe S.A. Rue de l'Industrie, Belgium). The cross-reactivity between the aforementioned buffalo and bovine cytokines has previously been documented by Mingala et al. (2007)

and Suzuki et al. (2011). Cytokine levels were determined according to the manufacturers' instructions; the plates were read at 450 nm and a correction wavelength of 550 nm was measured on a computerised automated microplate reader (Bio TEC, ELX800G, USA). Values were expressed in picograms per millilitre (pg/ml). Samples were run in duplicate for all of the examined cytokines and APPs. SSA was measured in the serum samples according to the method described by Shamberger (1984) using Ehrlich's reagent (made by adding 0.7 g of *p*-dimethylaminobenzaldehyde (Sigma Aldrich) to 150 ml of concentrated HCl and 100 ml of distilled water). The absorbance of samples was measured at 525 nm using a Jenway 7305 UV-VIS spectrophotometer (Jenway Scientific Equipments, UK). Data were expressed as mmol/l.

USG examination. USG was performed on all examined animals in standing position using 3.5 and 5.0 MHz transducers (Chison Medical imaging Co. Ltd., Wuxi, China 214142) according to Braun (2009) and Abouelnasr et al. (2013). The reticulum was examined by placing the transducer on both sides of the sternum and thorax up to the level of the elbow. The lung area was examined with the transducer held parallel to the 3rd to 11th intercostal space. For echocardiography, the 3rd, 4th and 5th intercostals were examined on both sides of the

thorax after moving the forelimbs cranially to help improve the contact between the probe and the intercostal space. Within the cardiac region, the heart, major blood vessels, valves, and the mediastinal region were scanned.

Laparo-rumenotomy. A left flank laparo-rumenotomy was performed in three cases in standing position under linear infiltrating anaesthesia using Lidocaine 2% according to Ducharme and Fubini (2004).

Gross post mortem examination (PM). Upon the approval of the Faculty Ethical Committee, four buffalo were subjected to complete PM examination and the findings were recorded.

Statistical analysis. Data were statistically analysed using the SPSS statistical software program (SPSS, version 15, USA). Significant differences ($P < 0.05$) among groups were compared using one-way ANOVA and the Duncan test. Means and standard deviation for each variable were calculated.

RESULTS

An overview of the clinical and laboratory findings in the diseased buffalo as well as controls is provided in Table 1 and in Figures 1–3. Clinically, the heart rates, but neither respiratory rate nor rectal

Table 1. Clinical findings in 32 buffalo with TRP compared to 10 controls

Variables	Control ($n = 10$)	Survivors ($n = 12$)	Non survivors ($n = 20$)
Heart rate (bpm)	63.0 \pm 8.07 ^a	87.66 \pm 6.18 ^b	88.80 \pm 6.79 ^b
Respiratory rate (breath/min)	12.66 \pm 1.96 ^a	13.00 \pm 2.28 ^a	14.33 \pm 1.63 ^a
Rectal temperature (°C)	37.85 \pm 0.17 ^a	38.80 \pm 1.13 ^a	38.90 \pm 0.67 ^a
Weakness and poor performance	absent ($n = 10$)	absent ($n = 7$) present ($n = 5$)	absent ($n = 9$) present ($n = 11$)
Abdominal distention	absent ($n = 10$)	absent ($n = 11$) present ($n = 1$)	absent ($n = 18$) present ($n = 2$)
Appetite	normal ($n = 10$)	inappetence ($n = 4$) anorexia ($n = 8$)	inappetence ($n = 4$) anorexia ($n = 16$)
Ruminal motility	normal ($n = 10$)	normal ($n = 0$) hypomotile ($n = 4$), ruminal stasis ($n = 8$)	normal ($n = 0$) hypomotile ($n = 5$), ruminal stasis ($n = 15$)
Recurrent tympany	absent ($n = 10$)	absent ($n = 2$) present ($n = 10$)	absent ($n = 1$) present ($n = 19$)
Lacrimation	absent ($n = 10$)	absent ($n = 4$) present ($n = 8$)	absent ($n = 6$) present ($n = 14$)
Lordosis/ bruxism	absent ($n = 10$)	absent ($n = 4$) present ($n = 8$)	absent ($n = 8$) present ($n = 12$)
Changes in milk yield	absent ($n = 10$)	sharp declined ($n = 12$)	sharp declined ($n = 12$) [*]

^{a, b}variables with different superscripts in the same row are significantly different at $P < 0.05$; ^{*}the remaining eight patients were dry cows in late gestation



Figure 1. Ultrasonogram of echogenic deposits on the reticulum of a buffalo with traumatic reticuloperitonitis viewed from the left ventral thorax. 1 = ventral abdominal wall, 2 = musculophrenic vein, 3 = echogenic deposits, 4 = intraperitoneal fluid, 5 = reticulum, Cr = cranial, Cd = caudal, D = diaphragm

temperature, were significantly higher in non-survivors compared to survivors (Table 1). In addition, the non-surviving buffalo were more likely to develop anorexia and weakness compared with survivors. However, rumen stasis, recurrent ruminal tympany, lacrimation, lordosis, bruxism and decreased milk production were commonly observed in both survivors and non-survivors. Biochemically, TNF- α , IL-1 β , IL-6, IL-10, SAA, Hp, Fb, CRP, and SSA levels were significantly higher in non-survivors compared with survivors as well as controls; on the other hand, serum IF- γ showed no difference between diseased buffalo and controls (Figure 2 and 3).

USG findings

In clinically healthy buffalo, the reticulum appeared as a semi-circular shape with a smooth

contour. However, in buffalo with TRP, the USG examinations showed fibrin deposits interspersed with fluid pockets between the reticulum, the dorsal ruminal sac and the diaphragm (Figure 1). Reticular abscesses were also observed as echogenic capsules with a hypoechogenic centre. None of the foreign bodies could be visualised. The amplitude and frequency of reticular contraction were decreased. With extensive adhesions, the frequency of contractions was decreased to one or zero contractions every two minutes.

Exploratory laparo-rumenotomy findings

Variable numbers and sizes of metallic foreign bodies (nails and wires) were removed from the ventral sac of the rumen and reticulum. Rumen contents were not fully macerated (impacted rumen). The animals were postoperatively treated with parenteral antibiotics, anti-histaminics, and anti-inflammatory drugs for five days. Complete recovery was achieved by 12 days postoperatively.

Necropsy findings

PM examination of slaughtered buffalo revealed the existence of foreign bodies (including pieces of wire and nails) that had perforated the reticular wall, were embedded in the diaphragm and formed peri-reticular abscess.

DISCUSSION

While significant progress has been made over the past years in the application of APPs as biomarkers in several animal diseases, the prognostic significance of inflammatory markers in buffalo




Figure 2. Means \pm SD of the selected acute phase cytokines in surviving and non-surviving buffalo with TRP compared to controls. Bars labelled with different letters are statistically significant ($P < 0.05$)




Figure 3. Means \pm SD of (A) Serum amyloid A ($\mu\text{g/ml}$), (B) fibrinogen (g/l), (C) haptoglobin (g/l), (D) CRP (mg/l), and (E) SSA (mmol/l) levels in surviving and non-surviving buffalo with TRP compared to controls. Bars labelled with different letters are statistically significant ($P < 0.05$)

with TRP has not yet been fully addressed. Our hypothesis was that inflammatory markers can help predict clinical outcome in buffalo with TRP. TRP in buffalo has become increasingly common and is particularly challenging to manage as its clinical presentation is often vague. In this study, a tentative diagnosis of TRP in diseased buffalo was initially based on clinical signs and laboratory findings, and later confirmed by USG evaluation, exploratory laparo-rumenotomy and/or necropsy in non-surviving cases. The classical clinical findings of TRP in diseased buffalo were consistent with previous reports (Saleh et al. 2008; Abdelaal et al. 2009; El-Ashker et al. 2013). Relative to cattle, indicators of abdominal pain including stiff gait, lordosis, and painful urination/defecation (Radostits et al. 2007) were less common in diseased buffalo, whereas lacrimation was very prominent. Thus, early diagnosis of TRP is more difficult in buffalo

compared to cattle. These findings are consistent with previous reports (Saleh et al. 2008; Abdelaal et al. 2009). Although clinical examination and USG were essential for diagnosis of TRP, they contribute little to discriminating non-survivors from survivors. Nonetheless, USG provides clear information with respect to the location and extent of the insult(s). Our findings in this regard were in part consistent with other reports (Abdelaal et al. 2009; Braun 2009; Abouelnasr et al. 2012).

Measurements of APPs in buffalo and cattle have been performed using standard commercial kits (El-Deeb and Iacob, 2012), and (Orro et al. 2011), which are similar to our methodology. On the other hand, the cross reactivity of some inflammatory cytokines (IL-1 α , IL-1 β , IL-6 and TNF- α) between buffalo and cattle has been previously described by Mingala et al. (2007). These authors reported the cloning, sequencing and phylogenetic analy-

sis of inflammatory cytokine (IL-1 α , IL-1 β , IL-6 and TNF- α) genes from swamp buffalo and two bubaline breeds, and the Bulgarian Murrah buffalo. The multiple sequence comparison showed a high homology between the bubaline breeds, which ranged from 99.3% to 100.0% similarity, whereas the comparison with cattle ranged from 98.6% to 99.0%. In addition, Suzuki et al. (2011), documented a cross reactivity between IL-2, IFN, IL-4 and IL-10 between African buffalo and cattle, and reported that the nucleotide sequence homology of IL-2, IFN- γ and IL-4 was more than 98%, which resulted in identical polypeptides. Also, the IL-10 gene of African buffalo and cattle exhibited 95% homology in nucleotide sequence, corresponding to thirteen amino acid residue substitutions. In this study, buffalo with TRP had higher values of all measured acute phase cytokines as well as inflammatory markers than controls, with higher levels in non-survivors compared with survivors. This intensive inflammatory reaction is followed by a compensatory anti-inflammatory response, during which the anti-inflammatory cytokines, including IL-10 are liberated into the bloodstream. The inhibitory effect of IL-10 on IL-1 β , IL-6, TNF- α , and IFN- γ supports a role for this cytokine in not only the regulation of T-cell responses, but also in acute inflammatory responses (Fiorentino et al. 1991). It is now recognised that pro-inflammatory cytokines are required for the initiation of an effective inflammatory response to infection and/or tissue injury. These pro-inflammatory cytokines are able to induce each other in a signalling cascade, thereby resulting in synergistic potentiating of pathological effects. Our findings were in part similar to those reported by Horadagoda et al. (2002) in buffalo following intravenous inoculation of *Pasteurella multocida* serotype B:2 endotoxin. In such report, serum concentrations of TNF- α rose within one hour *post inoculation* (*p.i.*) to reach peak values at 1–2 h *p.i.* and then declined rapidly to baseline levels 3–5 h *p.i.*; however, hp showed a delayed but prolonged increase in its values from 12 h *p.i.* and reached a plateau at 60 h *p.i.* It was reported that high concentrations of pro-inflammatory cytokines correlate with the prognosis of sepsis and the development of multiple organ dysfunction syndromes (Simmons et al. 2004). Likewise in animal models, the infusion of high concentrations of pro-inflammatory cytokines can lead directly to the development of multi-organ system failure (Kusawa et al. 1988). Several, but not all, studies have sug-

gested that levels of circulating pro-inflammatory cytokines can be used as a measure of the severity of illness and/or as a prognostic marker for patients with sepsis and multiple organ system failure (Makimura and Suzuki 1982; Rauchhaus et al. 2000; Orro et al. 2011; El-Ashker et al. 2013; El-Sebaei et al. 2013). It was also proposed that the quantification of blood APP levels can serve as both a diagnostic and prognostic tool when appropriate sampling schedules are performed (Murata, et al. 2004; Cray et al. 2009; Orro et al. 2011). Similarly, Jafarzadeh et al. (2004) demonstrated that serial interpretation of total plasma protein and plasma Fb with suggested cut-off points is useful for differentiating TRP from other gastrointestinal problems in cattle. Several studies have previously demonstrated the significance of Fb, Hp, and CRP as useful biochemical parameters for evaluating the incidence and severity of inflammatory reactions in bovine species with inflammatory conditions (McSherry et al. 1970; Makimura and Suzuki 1982; Makimura and Usui 1990; Lee et al. 2003; Jafarzadeh et al. 2004, Orro et al. 2011). CRP was also found to be both a useful marker for evaluating the health status of a herd, and a parameter to assess individual stress levels. CRP may be useful in early surveillance of intra-herd TRP (Lee et al. 2003), and may also fulfil an important function in defence against infection and control of the inflammatory response (Mold et al. 2002). It is suggested that CRP can be used as a sensitive biomarker of TRP in water buffalo.

The results of our study suggest that the dysregulation of pro-inflammatory and anti-inflammatory cytokine networks may occur in parallel in buffalo with TRP, and imply that the overall degree of cytokine network disruption may be an important prognostic indicator. Therapeutic strategies to modulate inflammation must take into account the complex cytokine biology in buffalo patients with TRP.

Acknowledgement

The authors would like to thank Prof. Dr. Sabry El-Khodery (Faculty of Veterinary Medicine, Mansoura University, Egypt) for his kind and valuable assistance during sonographic examination. Special thanks are given to Dr. Mahmoud El-Sebaei (Faculty of Veterinary Medicine, Mansoura University, Egypt) for his kind help during SSA measurement. The authors of this paper hold no financial or per-

sonal relationships with individuals or organisations that would inappropriately influence or bias the findings of this paper.

REFERENCES

- Abdelaal A, Floeck M, El Maghawry S, Baumgartner W (2009): Clinical and ultrasonographic differences between cattle and buffalo with various sequelae of traumatic reticuloperitonitis. *Veterinarni Medicina* 9, 399–406.
- Abouelnasr K, Mosbah E, Karrouf G, Zaghloul A (2012): Comparative ultrasonographic findings of traumatic reticulitis, perireticular abscess and diaphragmatic hernia in buffalo (*Bubalus Bubalis*). *Journal of American Science* 12, 590–595.
- Abouelnasr K, Mosbah E, Karrouf G, Zaghloul A (2013): Ultrasonography of normal reticulum in 30 healthy buffalo (*Bubalus bubalis*). *Journal of Applied Animal Research*, DOI: 10.1080/09712119.2013.823435.
- Aref N, Abdel-Hakim M (2013): Clinical and diagnostic methods for evaluation of sharp foreign body syndrome in buffaloes. *Veterinary World* 9, 586–591.
- Braun U (2009): Ultrasonography of the gastrointestinal tract in cattle. *Veterinary Clinics of North America: Food Animal Practice* 25, 567–590.
- Cray C, Zaias J, Altman N (2009): Acute phase response in animals: A Review. *Comparative Medicine* 59, 517–526.
- Ducharme NG, Fubini SL (2004): Surgery of the forestomach compartments. In: Fubini SL, Ducharme NG. (eds): *Farm Animal Surgery*. Elsevier Science, Saunders, USA. 184–194.
- El-Ashker M, Salama M, El Boshy M (2013): Traumatic reticuloperitonitis in water buffalo (*Bubalus bubalis*): Clinical findings and the associated inflammatory response. *Journal of Veterinary Medicine*. <http://dx.doi.org/10.1155/2013/808656> (Accepted 7 July 2013)
- El-Deeb W, Iacob O (2012): Serum acute phase proteins in control and *Theileria annulata* infected water buffaloes (*Bubalus bubalis*). *Veterinary Parasitology* 190, 12–18.
- El-Sebaei M, El-Ashker M, El-Boshy M (2013): The role of acute phase cytokines in the recovery and disease progress of *Theileria annulata*-infected cattle. *Comparative Clinical Pathology*. DOI 10.1007/s00580-013-1812-7.
- Fiorentino D, Zlotnik A, Mosmann T, Howard M, O'Garra A (1991): IL-10 inhibits cytokine production by activated macrophages. *Journal Immunology* 147, 3815–3822.
- Horadagoda NU, Hodgson JC, Moon GM, Wijewardana TG, Eckersall PD (2002): Development of a clinical syndrome resembling haemorrhagic septicaemia in the buffalo following intravenous inoculation of *Pasteurella multocida* serotype B:2 endotoxin and the role of tumour necrosis factor-alpha. *Research in Veterinary Science* 72, 194–200.
- Jafarzadeh S, Nowrouzian I, Khaki Z, Ghamsari M, Adib-hashemi F (2004): The sensitivities and specificities of total plasma protein and plasma fibrinogen for the diagnosis of traumatic reticuloperitonitis in cattle. *Preventive Veterinary Medicine* 65, 1–7.
- Johnson RW (1997): Inhibition of growth by pro-inflammatory cytokines; an integrated view. *Journal Animal Science* 75, 1244–1255.
- Koj A (1997): Initiation of acute phase response and synthesis of cytokines. *Biochimica et Biophysica Acta* 1317, 84–94.
- Kusawa S, Gelfand JA, Ikejima T (1988): Interleukin 1 induces a shock-like state in rabbits. Synergism with tumor necrosis factor and the effect of cyclooxygenase inhibition. *Journal of Clinical Investigation* 81, 1162–1172.
- Lee WC, Hsiao HC, Wu YL, Lin JH, Lee YP, Fung HP, Chen HH, Chen YH, Chu RM (2003): Serum C-reactive protein in dairy herds. *Canadian Journal Veterinary Research* 67, 102–107.
- Makimura S, Suzuki N (1982): Quantitative determination of bovine serum haptoglobin and its elevation in some inflammatory diseases. *Japanese Journal Veterinary Science* 44, 15–21.
- Makimura S, Usui M (1990): Correlation between haptoglobin and sialic acid or mucoprotein in diseased bovine serum. *Japanese Journal Veterinary Science* 52, 1245–1250.
- McSherry BJ, Horney FD, deGroot JJ (1970): Plasma fibrinogen levels in normal and sick cows. *Canadian Journal Comparative Medicine* 34, 191–197.
- Mingala C, Odbilega R, Konnaia S, Ohashia K, Onumaa M (2007): Molecular cloning, sequencing and phylogenetic analysis of inflammatory cytokines of swamp type buffalo contrasting with other bubaline breeds. *Comparative Immunology, Microbiology and Infectious Diseases* 30, 119–131.
- Misk N, Semieka M, Ali S (2001): Varieties and sequelae of ingested foreign bodies in buffalo and cattle. *Assiut Veterinary Medical Journal* 46, 250–273.
- Mold C, Rodriguez W, Rodic-Polic B, Du Clos TW (2002): C-reactive protein mediates protection from lipopolysaccharide through interactions with Fc-gammaR. *Journal of Immunology* 169, 7019–7025.
- Morimatsu M, Sakai H, Yoshimatsu K, Minowa O, Yamamoto S, Yatomi K, Fujinaga T, Naiki M (1989): Isolation and characterization of C-reactive protein and serum

- amyloid P component from bovine serum. Japanese Journal Veterinary Science 50, 723–732.
- Mousavi G, Hassanpour A, AmoghliTabrizi B, Rezaie A (2007): Electrocardiographic changes in buffalo with Traumatic reticuloperitonitis. Italy Journal Animal Science 6, 1029–1031.
- Murata H, Shimada N, Yoshioka M (2004): Current research on acute phase proteins in veterinary diagnosis: an overview. Veterinary Journal 168, 28–40.
- Orro T, Pohjanvirta T, Rikula U, Huovilainen A, Alasuutari S, Sihvonon L, Pelkonen S, Soveri T (2011): Acute phase protein changes in calves during an outbreak of respiratory disease caused by bovine respiratory syncytial virus. Comparative Immunology, Microbiology and Infectious Diseases 34, 23–29.
- Radostits OM, Gay CC, Hinchclif KW, Constable PD (2007): Manifestations of circulatory failure. In: Veterinary Medicine: A Textbook of the Diseases of Cattle, Horses, Sheep, Pigs and Goats. 10th ed. Elsevier Health Sciences, Philadelphia, PA, USA. 402–431.
- Rauchhaus M, Doehner W, Francis D, Davos C, Kemp M (2000): Plasma cytokine parameters and mortality in patients with chronic heart failure. Circulation 102, 3060–3067.
- Saleh M, Rateb H, Misk N (2008): Comparison of blood serum proteins in water buffalo with traumatic reticuloperitonitis and sequellae. Research in Veterinary Science 85, 208–213.
- Shamberger RJ (1984): Serum sialic acid in normals and in cancer patients. Journal Clinical Chemistry and Clinical Biochemistry 22, 647–651.
- Simmons E, Himmelfarb J, Sezer T, Chertow G, Mehta R, Mehta RL, Paganini EP, Soroko S, Freedman S, Becker K, Spratt D, Shyr Y, Ikizler TA (2004): Clinical nephrology –epidemiology – clinical trials. Plasma cytokine levels predict mortality in patients with acute renal failure. Kidney International, 65, 1357–1365.
- Suzuki S, Konnai S, Okagawa T, Githaka N, Kariuki N, Gakuya F, Kanduma E, Shirai T, Ikebuchi R, Ikenaka Y, Ishizuka M, Murata S, Ohashi K (2011): Molecular cloning and characterization of Th1 and Th2 cytokines of African buffalo (*Syncerus caffer*). International Journal of Immunology. DOI: 10.1111/j.1744-313X.2011.01062.x
- Yoshioka M, Watanabe A, Shimada N, Murata H, Yokomizo Y, Nakajima Y (2002): Regulation of haptoglobin secretion by recombinant bovine cytokines in primary cultured bovine hepatocytes. Domestic Animal Endocrinology 23, 425–433.

Received: 2014–03–24

Accepted after corrections: 2014–06–18

Corresponding Author:

Maged El-Ashker, Mansoura University, Faculty of Veterinary Medicine, Department of Internal Medicine and Infectious Diseases, Mansoura 35516, Egypt
E-mail: maged_elashker@yahoo.com
