

State and development of phytocenoses on research plots in the Krkonoše Mts. forest stands

S. VACEK¹, K. MATĚJKA²

¹Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic

²IDS, Prague, Czech Republic

ABSTRACT: The paper assesses the state and development of phytocenoses in beech, mixed and spruce stands on permanent research plots (PRP) 1–32 in the Krkonoše (Giant) Mts. in the years 1980–2005, i.e. during the air-pollution calamity and afterwards. Dynamics (the extent of change) of the vegetation structure has been expressed as the overall change of species composition in comparison with the year 1980. The change was quantified using the Euclidean distance or as the change of the several first ordination axes (DCA 1–DCA 4). Species composition was significantly changing on all 32 PRP stands in the period 1980–2005; some species completely disappeared (e.g. *Cicerbita alpina*, *Lamium maculatum*, *Phyteuma spicatum*, *Viola biflora*) or their ratio was reduced (e.g. *Blechnum spicant*, *Dentaria enneaphyllos*, *Homogyne alpina*). From the viewpoint of diversity, the most significant change is the reduction of species in the moss layer, which was observed in all types of stands; in beech and mixed forests the average number of species dropped from 4.8 to 2.7 (44%), in non-declining spruce stands the number fell from 11.6 to 5.5 (53%), and in declining spruce stands it dropped from 10.4 to 3.3 (68%). The overall reduction of species diversity ranged between 31% and 43%; the highest reduction was recorded at species with lower representation. The most significant factors influencing the species composition were altitude and exposition of plot.

Keywords: beech; mixed and spruce stands; classification; Krkonoše Mts.; ordination; state and development of phytocenoses

The state of vegetation of the Krkonoše (Giant) Mts. in the 13th century, i.e. before the impact of human activities on landscape, was the result of flora quaternary development. Mikyska in their vegetation reconstruction (MIKYŠKA et al. 1968) reported that the natural basis of the stands in the lowest parts of the Krkonoše Mts. were flowery and woodrush meadows, followed upward by acidophilic mountain beech forests, climax spruce forests, and the highest altitudes were covered with mountain pine and subalpine communities. Alluvial plains, alder carrs, scree forests, moist spruce forests, upland moors, and valley bogs occurred in a mosaic pattern. A map of potential vegetation in the Krkonoše National Parks (Krkonošský národní park and Karkonozsky Park Narodowy) was constructed earlier (Fig. 1; see VACEK et al. 2006).

Species composition of the Krkonoše Mts. forests has changed significantly, especially during the last four centuries. Original beech, spruce-fir-beech and partly spruce forests were destroyed by human activities, such as chalet farming, mining and glass industry. The forests were slowly restored, yet, their species, genetic, age and location structure was significantly changed. Original montane spruce forests were preserved only in a thin belt along the upper forest boundary, on steep slopes and near the glacial cirques. The remains of original mixed and beech forests may be found mostly in the natural reserves (zone I) and partly in inaccessible locations of protective forests.

As to the forests vegetation zones, the largest are: FVZ 6 – spruce with beech (42.5%), FVZ 8 –

Supported by the Ministry of Education, Youth and Sports of the Czech Republic, Project No. 2B06012 .

spruce (20.7%) and FVZ 7 – beech with spruce (19.2%). Dominant are forest types of following groups (FTG): 6K – acid spruce with beech (18.8%) and 7K – acid beech with spruce (10.6%). Acid sites significantly predominate (acid and extreme ecological series) – 72% over fertile (fertile and humus-enriched ecological series) sites – 17%, and moist (moist ecological series, water-enriched, gleyic or peat) sites – 11% (MIKESKA et al. 2007).

Forests of the Krkonoše Mountains are an important landscape component of the Czech Republic. They are object of a special interest in relation to the natural environment protection, stabilisation of natural processes and the overall landscape homeostasis. Moreover, they serve for several production and non-production functions. Attention paid to mountain forest ecosystems during the last two decades is undoubtedly rightful. Mountain forest ecosystems are generally very sensitive, and even in natural environmental conditions they are exposed to many stress factors. The most important stressors are impacts of extreme climate (low temperature, drought periods, intensive air flows), but also soil conditions may not be often favourable (mostly it is the case of relatively poorly developed soils and soils naturally considerably acid with low biological activity). For these reasons, dynamics of forest ecosystems is often a subject of interest of environmentalists and silviculturists. Attention is usually paid to natural or near-natural forests; however, these are often more or less afflicted with imission-ecological stress, and thus in different stage of forest development cycle or in different level of decline. Generally, a spatially limited community is observed; usually it is a research plot that sufficiently represents the community (mostly plant), not the whole forest ecosystem; the latter is composed of smaller or larger areas that represent different stages of the forest development. Yet, this is not the last problem. During the whole period of development of contemporary forest generation, significant changes of conditions under which these forests grew have been going on. To put it simple, let us mention only some of the most important ones:

- (1) change of the air pollution load with an increase up to the early 1990s, and subsequent irregular decrease of this load and a change of proportional representation of individual pollutants;
- (2) change of silvicultural technologies and preference of certain woody species;
- (3) forest restoration at sites that had not been afforded for periods of variable length (MATĚJKA, VACEK 2007).

Long-term dynamics of vegetation changes at similar sites and stand conditions has been studied by

many authors. Classification and ordination methods at beech, mixed (spruce with beech, beech with spruce and spruce with fir and beech) and spruce forests were studied for instance by SÝKORA (1971), SOFRON (1981), VACEK and LEPŠ (1991, 1992, 1999), AMBROS and MÍCHAL (1992), VIEWIGH (1994), ELENBERG (1996), BOHN et al. (2000), POKORNÝ (2002), HÉDL (2004), VACEK and MATĚJKA (2003), ŠAMONIL and VRŠKA (2008), PODLASKI (2008), who dealt with suitability of use of different input data for numerical analysis. In numerous cases, the impact of imission load on acceleration of vegetation changes was proven (VACEK, LEPŠ 1996; MOLDAN, HAK 2007; VACEK et al. 2007). Yet, long-term vegetation observations under natural conditions with almost minimal air-pollution load are rather few (MATUSZKIEWICZ et al. 1960, 1967; STÖCKER 1968; MIKYŠKA 1972; MÁLEK 1970, 1973; PIŠTA 1972, 1978; MORAVEC 1974; VACEK 1981; PRŮŠA 1985; UJHÁZY et al. 2005).

Evaluation of recorded changes in concrete site and stand conditions and their interpretation is considered as one of the most critical issue of phytocenological monitoring (SAMEK 1986). He underlines that the assessment of the rate of anthropogenic impact must necessarily stem from comparison and quantification of the observed impact. These conditions are often not fulfilled. Therefore, the above mentioned monitoring type should be set at sites of some other experimental measurements or autecological studies. According to BRANG (2005), it is important for further implementation of these findings in forest ecosystem management. Our research of forest ecosystem on permanent research plots in the Krkonoše Mts. fulfilled this important condition.

The aim of this paper was to evaluate the state and development of phytocenoses in beech, mixed and spruce stands on permanent research plots in the Krkonoše Mts. in the years 1980–2005, i.e. during the air-pollution calamity and afterwards.

MATERIAL AND METHODS

Characteristics of permanent research plots

As in the research on state and development of soils, the state and development of phytocenoses in beech, mixed (beech with spruce to spruce to beech) and spruce stands was carried out on research plots 1–32 in the Krkonoše Mts.; their characteristics are given in our previous paper (MATĚJKA et al. 2010).

Phytocenological relevés were recorded in the years 1980 (1976) to 2005 (with 5-year intervals: 1980, 1985, 1990, 1995, 2000 and 2005) by usual means, using the combined eleven-point Domin-Hadač scale for abundance and dominance. Phytocenological relevés were collected in database of the DBreleve software. It is a system developed by the IDS company for unified editing, storage and further processing of the records of species structure of communities (MATĚJKA 2000). For calculations, the species representations were transformed: scale values were replaced by the values of average cover of individual scale levels and such values were multiplied by a constant (specific for the relevé and the etage), thus sum of representations of all species participating in the etage of the relevé should be equal to the total etage coverage.

To compute ordination (using the DCA method) and divisive classification (the TWINSpan procedure; HILL 1979) the relevé data were exported. Ordination and classification results were again downloaded to the DBreleve programme. Both numerical methods were applied to process the data of herb (E1) and moss (E0) layer.

The evaluation of relevés was done based on the following methods:

- Evaluation of species diversity indices in the course of the period of observation: total species diversity (Shannon-Wiener's diversity index, H), species richness (S) and evenness (e).
- Classification of all phytocenological relevés using the TWINSpan procedure.
- Agglomerative hierarchical classification of relevés, method of average linkage; Euclidean distance was used as a dissimilarity measure.
- Ordination of relevés using the DCA method using Canoco, version 4.5; data from all plots and all sampling periods were used together. Results were plotted within the PlotOA programme (MATĚJKA 2009).

Extent of vegetation change throughout the period of monitoring (starting in the year 1980) was evaluated on the basis of the sum of variance of the first (one to four) DCA axes (the results of ordination from all plots were used) – using the formula

$$v_i = \sqrt{\sum_{j=1}^i \text{var}(DCA_{j,r})} \quad (1)$$

where:

- j – index of ordination axis,
- r – index of year.

Random fluctuations were thus eliminated. Compare MATĚJKA (2009), where $V_i = v_i^2$

Total difference (E_{r_1,r_2}) in species structure between two years (between a year and the original state in 1980, specially) was evaluated on the basis of the Euclidean distance between vectors of individual species representation in both phytocenological relevés compared. Maximal change in the vegetation structure during the whole observation period (1980–2005) was specified as $E_{\max} = \max(E_{1980,r})$.

Total cover of woody plants was evaluated using the combination of total cover of tree and shrub layers. Both layers produce a shelter for herbal vegetation and during the regeneration of the declined cover, the plants gradually pass from shrub to tree layer. Canopy coverage (D) was expressed as follows:

$$D = 1 - [(1 - E2) \times (1 - E3)] \quad (2)$$

where:

E3, E2 – the cover of tree and shrub layer.

RESULTS AND DISCUSSION

Numeric classification

Classification using the TWINSpan procedure (Table 1) distinguished individual basic community classification classes. It is possible to describe them on the basis of their specific species composition.

The communities were divided as supramontane spruce forests (group *00; typical is occurrence of *Streptopus amplexifolius*, *Athyrium distentifolium*, *Sphagnum girgensohnii*, *Deschampsia cespitosa* and *Myliia taylorii*). Within this group, species composition on plots 13 and 14 are rather specific (group *0010 with *Adenostyles alliariae* and *Rhizomnium punctatum*); in the course of development, the canopy completely or partially declined, and specific was also the plant community on plot 23 (group *000 with *Carex limosa* as indicator species); all of them appear stable.

Among typical supramontane spruce stands belong communities of the classification group *0011 on plots 11, 12, 5, 4, 10, 2, and 21, with a stable canopy species composition. At the beginning of the 1980's, this group comprised also the plots 3, 25, 15, and 17; at these plots, however, the canopy sooner or later declined and as a result, the classification changed to montane spruce forest (*010). Opening of canopy probably resulted in a change of the temperature regime. A similar change was observed at plot 20, which may indicate its position at the verge between montane and supramontane spruce forests.

Typical montane spruce forests are gathered in the classification group *010 (or subgroup *0100),

Table 1. Results of the classification using the TWINSPAN procedure. At each classification group the indicator species and their representation according to "cut-levels" [0.00% (1)–3.12% (2)–6.25% (3)–12.50% (4)–25.00% (5)–50.00% (6)] are given. The last but one column contains labels of relevés belonging to respective classification group (No. of plot – one or more years [the last two digits of the year] of the relevé). The years following the stand decline are marked red. The relevés dated from 1980 or before are typed in bold. The last column provides the stand type, based on the apportionment of spruce (sp) and beech (be)

		*0000					23–80,85,90	
	*000	<i>Bar gra</i> 1						
	<i>Car lim</i> 1	*0001					23–95,00,05	
	*00		*001000				14–76,80,85	
	<i>Str amp</i> 1		*00100	<i>Ace pse</i> 1				
	<i>Ath dis</i> 1		<i>Vio bif</i> 1					
	<i>Sph gir</i> 1		*001001				13–76,80,85	
	<i>Des ces</i> 1							
	<i>Myl tay</i> 1	*0010		*0010100			14–90,05	
		<i>Ade all</i> 1	*001010					
		<i>Rhi pun</i> 1		*0010101			14–95,00	
			*00101					
				*0010110			13–90,95	
				<i>Phe con</i> 1	*0010111		13–00,05	
					<i>Bet pen</i> 1			
				*00110000			11–76,80,85	
				<i>Bar gra</i> 1				
					*001100010		12–76,80,85	
					<i>Pel nee</i> 1			
				*0011000				
				*00110001		*0011000110	12–90,95,00,05	
			*001100			<i>Phe con</i> 1		
			<i>Ath dis</i> 1		*001100011			
						*0011000111	11–95,00,05	
*0								
<i>Ave fle</i> 2			*0011001	*001100100			05–80,85	
<i>Tri eur</i> 1			<i>Sph gir</i> 3	<i>Luz luz</i> 1	*001100101			sp
					<i>Vac myr</i> 2		05–90,95,00,05	
	*001				*001100110		03–80	
							04–80,85,90	
					*001100111			
					<i>Jun fil</i> 1		04–95,00,05	
			*00110					
*01					*001101000		10–80,85	
<i>Vac myr</i> 5					*00110100	<i>Luz pil</i> 1		
		*0011			<i>Des ces</i> 1			
					*001101001		10–90,95,00,05	
				*0011010			20–80	
							24–90,95,00,05	
					*001101010			
			*001101				25–80,85,90	
			<i>Oxa ace</i> 1		*0011010101	<i>Pol com</i> 1		
			<i>Gen asc</i> 2					
					*001101011		24–80,85	
					<i>Ant odo</i> 1			
					*001101100		21–90,95	
				*00110110				
			*0011011		*001101101		21–00,05	
			<i>Pol ver</i> 1		<i>Ath dis</i> 1			
					*00110111		21–80,85	
					<i>Ant odo</i> 1			
			*00111	*001110			15–90	
			<i>Dry dil</i> 4				17–80,85	
				*001111				
				<i>Vac vit</i> 1			15–76,80,85	

Table 1 to be continued

		*010000			15-95,00,05	
		<i>Ade all 1</i>				
		*010000	*0100010		22-80,85	
		<i>Poh nut 1</i>	<i>Cam lat 1</i>			
		*010001	*01000110		22-00,05	
			*0100011	*01000111	22-90,95	
				<i>Sph squ 1</i>		
			*0100100	*01001000	18-80,85	
			<i>Myl tay 1</i>	<i>Lyc ann 1</i>		
				*010010001	18-90,95	
				*010010010		
				<i>Sca und 1</i>		
			*01001001	<i>Epi ang 1</i>	16-80,85,90,95	
				<i>Sen her 1</i>		
	*0100				19-80,85,90	
		*010010	*010010011	*0100100110	20-85	
*010		<i>Dry dil 1</i>	<i>Oxa ace 1</i>	<i>Pol for 2</i>	25-95	
<i>Ave fle 4</i>					26-80	
<i>Tri eur 1</i>	*01001	<i>Oxa ace 1</i>	<i>Sph gir 1</i>			
<i>Ave fle 5</i>	<i>Ave fle 5</i>			*0100100111	20-90,95,00,05	sp
					16-00	
			*0100101	*010010100	25-00,05	
			<i>Gal har 1</i>	*010010101	18-00,05	
			<i>Gen asc 1</i>	<i>Des ces 1</i>		
			<i>Epi ang 1</i>	*010010110	17-90,95,00,05	
				*010010111		
			<i>Dic sco 2</i>	*010010111	11-90	
		*010011	*0100110		26-85,90,95	
		<i>Bet pen 1</i>	*0100111		26-00,05	
*0		<i>Cal vul 1</i>	<i>Gal har 1</i>			
<i>Ave fle 2</i>	*01				16-05	
<i>Tri eur 1</i>	<i>Vac myr 5</i>				19-00,05	
		*0101	*010100			
		<i>Epi mon 1</i>	*010101		03-95,00,05	
		<i>Cal vil 6</i>	<i>Luz luz 1</i>			
		<i>Gen asc 1</i>	*01011		03-85,90	
			<i>Pla und 1</i>			
			*011000		06-80,85	
			<i>Ble spi 1</i>			
		*01100			06-00	
		<i>Cal vil 5</i>	*0110010		19-95	
			*011001		06-90,95	
			<i>Hom alp 1</i>			
		*0110	*0110100		09-00,05	be-sp
		<i>Gen asc 1</i>	<i>Gym dry 1</i>			(sp)
			<i>Ave fle 1</i>		09-90,95	
			*0110101	*01101010		
		*01101		*01101011	09-80,85	
				<i>Cic alp 1</i>		
			*011011		08-95	
					02-00,05	
			*0111000		28-95	
				*011100100	27-90,95,00	
		*011100	*01110010	<i>Tri eur 1</i>		
		<i>Sor auc 1</i>	*0111001	<i>Pic abi 1</i>	27-05	be-sp
			<i>Luz luz 1</i>	*011100101	28-80,90	to
	*0111	*01110			27-80,85	be
		<i>Mai bif 1</i>		*01110011	28-85	
					02-90,95	
			*0111010			
			*011101	*0111011	02-80,85	
			<i>Ace pse 1</i>			

Table 1 to be continued

		*011				*0111100	01–00,05	
*0		<i>Fag syl</i> 1		*01111	*011110	*0111101	01–90,95	be-sp
<i>Ave fle</i> 2	*01	<i>Dic het</i> 1	*0111	<i>Ste nem</i> 1		<i>Bet pen</i> 1		to
<i>Tri eur</i> 1	<i>Vac myr</i> 5	<i>Pol ver</i> 1		<i>Pel nee</i> 1	*011111		01–80,85	be
		<i>Pre pur</i> 1			<i>Ath fil</i> 1			
			*1000	*10000			06–05	
			<i>Par qua</i> 1	<i>Ble spi</i> 1				
			<i>Gen asc</i> 1		*100010		07–80,85	
			<i>Phe con</i> 1			*1000110	07–90,95	be-sp
			<i>Ste nem</i> 1	*10001	*100011	*1000111		
			<i>Rub ida</i> 1		<i>Ath fil</i> 1	*1000111	07–00,05	
			<i>Cal vil</i> 2			<i>Asp odo</i> 1		
	*10	*100		*10010			08–00,05	
				<i>Ble spi</i> 1				
					*100110		28–00,05	
			*1001		<i>Vac myr</i> 3		29–90	be
				*10011		*1001110	29–95,00,05	to
						*1001111	31–05	be(sp)
						<i>Bra syl</i> 1	32–05	
*1							29–80,85	be(sp)
<i>Gym dry</i> 1		*101	*1010					
<i>Dry fil</i> 1		<i>Myc mur</i> 1	*1011				08–80,85,90	be-sp
<i>Ath fil</i> 1		<i>Gym dry</i> 2	<i>Ble spi</i> 1					
				*11000			31–80,85	
				<i>Asp odo</i> 1				
				*1100	*110010		31–95	
					<i>Dry fil</i> 4			
	*11			*11001			31–90,00	
<i>Mil eff</i> 2		*110			*110011		32–95	be
					<i>Asp odo</i> 1			
			*1101	*11010			32–90,00	
			<i>Lun red</i> 1					
				*11011			32–80,85	
							30–95,00,05	
			*1110					
		*111					30–80,85,90	
		<i>Act spi</i> 1	*1111					
			<i>Cir x i</i> 1					

where the predominant species is *Avenella flexuosa*, and *Trientalis europaea* and *Epilobium angustifolium* occur. These communities were reported on plots 22, and 26 that were rather stable during the observation period, and on plots 6, 18, and 19, where the canopy decline occurred. This classification group includes also the communities developed after the canopy decline, with classification subgroups *0101 (characterised by dominant *Calamagrostis villosa* and further *Gentiana asclepiadea* and *Epilobium montanum*) and *0100101 (with appearance of the species *Galium saxatile*, *Gentiana asclepiadea* and *Epilobium angustifolium*).

Beech-spruce forests of the 5th to 7th vegetation zone belong into the classification group *011. They contain many indicators of spruce forests together with beech types. The structure of these communities (plots 1, 2, 6, 9, 27) is rather stable in the course of time; exceptionally the community may be shift-

ed into the group *10 (plot 28), which may be interpreted as a change resulting in warming of the stand.

Communities of beech to spruce-beech stands at plots 7, 8, and 29 are similar to those of the previous group, however, the indicators of typical forest stand miss here. Stands correspond to the 6th to 7th vegetation zone and are classified in the group *10.

Typical beech stands on plots 30, 31, and 32 have communities of the classification group *11.

Ordination analysis

The interpretation of the first two ordination axes (Figs. 2 and 3) is relatively easy, compared to the third and four ordination axes (Figs. 4 and 5). High values along the first axis are observed in beech forest (plots 30, 31, 32); they usually show significant changes during the time, as well. Plots 7, 8, and 29 represent




Fig. 1. Potential vegetation of the Giant Mountains (VACEK et al. 2006)




Fig. 2. Position of all relevés from PRP 1–32 in the DCA ordination space of the first two axes. Each series of relevés from one plot is connected with a trajectory with a plot number at its beginning (corresponding to the year 1980; at plots 11 to 15, to the year 1976)




Fig. 3. Position of species in ordination space of the DCA 1 and DCA 2 axes




Fig. 4. Position of all phytocenological relevés from PRP 1–32 in the DCA ordination space of the 3rd and 4th axes. Each series of relevés from one plot is connected with a trajectory with a plot number at its beginning (corresponding to the year 1980; at plots 11 to 15, to the year 1976)

typical spruce-beech stands, followed, with the decreasing score, by stands of higher altitudes.

The second ordination axis discerns supramontane spruce communities with occurrence of subalpine tall-forb species; typical examples are plots 13 and 14, whereas the plot 23 lies at the opposite.

Gradual movement of the plots along the trajectories to the centre of ordination space is evident. It confirms reduction of forest community diversity in the Krkonoše Mts. area. There is, however, another distinguishable trend, characteristic for beech stands (plots 7, 8, 1, 2, 28, 6, 9); it results in a certain break of trajectory in the direction of increasing values of both the first and the second ordination axis.

Vegetation development on particular PRP

The development of various vegetation characteristics studied from 1980, on plots 11–15 from 1976, was presented in a graphic form; each plot was sorted in one of the basic classification groups based on the 1980 relevé.

Spruce stands – group *0000: PRP 23.

Spruce stands – group *0010: On both plots classified here (PRP 13 and 14), canopy decline was reported.

Spruce stands – group *0011: PRP 10, 21, 20, 24, 4, 12, 15, 5, and 11 (in respective order rises the extent of species composition change); next, this group includes PRP 3, 25 and 17, where canopy decline was reported. Except for PRP 17, these plots have autochthonous spruce vegetation. The change of floor layer vegetation is usually weak, medium or strong (even in the case of canopy stability).

Spruce stands – group *0100: this group comprises PRP 22 and 18 with vegetation classified as autochthonous, and PRP 16, 26 and 19 with allochthonous vegetation. The canopy decline was observed at all the plots, except for PRP 22.

Comparison of all spruce stands shows that the canopy decline is not a cause of the greatest changes in floor layers, if these changes are evaluated quantitatively using the change in cover of present species (e.g. Euclidean distance or score along the ordination axes). The changes may be noticed more in the vegetation classification, especially in the data such as presence – absence of species.

Spruce-beech stands – group *0110: PRP 9, 6.

Spruce-beech to beech stands – group *0111: PRP 2, 27, 1 and 28.

Spruce-beech stands – group *1000 (sp-be): PRP 7.

Beech stands with the admixture of spruce – group *1010: PRP 29.




Fig. 5. Position of species in ordination space of the DCA 3 and DCA 4 axis

Table 2. Basic characteristics of plant community dynamics on studied plots

PRP	Stand type	Autochthonous vegetation	Canopy decline	TWINSPAN 1980	Basic classification group	Classification level change	Classification				
							v_1	v_2	v_3	v_4	E_{max}
23	sp	+		*00001	*0000	5	0.000	0.300	0.321	0.369	31.4
14	sp	+	+	*0010000	*0010	6	0.095	0.313	0.329	0.359	38.0
13	sp	+	+	*0010010	*0010	6	0.032	0.257	0.303	0.321	52.5
10	sp	+		*001101000	*0011	10	0.032	0.045	0.055	0.134	17.1
21	sp	+		*001101111	*0011	9	0.032	0.071	0.105	0.286	27.7
20	sp	+		*0011010100	*0011	3	0.000	0.055	0.100	0.210	30.7
24	sp	+		*001101011	*0011	10	0.000	0.000	0.084	0.219	34.5
4	sp	+		*0011001101	*0011	10	0.032	0.077	0.122	0.179	47.4
12	sp	+		*0011000100	*0011	10	0.045	0.239	0.257	0.283	51.9
15	sp	+		*00111111	*0011	3	0.084	0.130	0.268	0.802	60.5
5	sp	+		*001100100	*0011	10	0.000	0.207	0.232	0.265	68.3
11	sp	+		*001100001	*0011	3	0.063	0.395	0.420	0.442	76.1
3	sp	+	+	*0011001100	*0011	3	0.077	0.077	0.105	0.145	18.1
25	sp	+	+	*0011010101	*0011	3	0.032	0.045	0.077	0.155	37.4
17	sp		+	*0011101	*0011	3	0.141	0.167	0.365	0.660	53.1
22	sp	+		*0100010	*0100	8	0.084	0.089	0.100	0.164	35.7
18	sp	+	+	*010010000	*0100	8	0.032	0.045	0.055	0.241	23.1
16	sp		+	*0100100101	*0100	5	0.095	0.095	0.167	0.184	31.7
26	sp		+	*0100100110	*0100	7	0.000	0.055	0.130	0.265	38.1
19	sp		+	*0100100110	*0100	4	0.071	0.077	0.155	0.219	51.0
9	be-sp	+		*01101011	*0110	8	0.243	0.332	0.630	0.792	29.1
6	be-sp	+		*011000	*0110	2	0.261	0.307	0.449	0.508	33.5
2	be-sp	+		*0111011	*0111	7	0.297	0.387	0.837	0.901	29.1
27	be(sp)	+		*011100111	*0111	9	0.055	0.071	0.118	0.237	29.8
1	be-sp			*0111111	*0111	7	0.321	0.415	0.740	0.764	34.6
28	be	+		*0111001011	*0111	2	0.241	0.283	0.446	0.480	35.1
7	be-sp	+		*100010	*1000	7	0.311	0.404	0.909	0.931	34.8
29	be(sp)	+		*1010	*1010	4	0.342	0.346	0.567	0.673	7.5
8	be-sp	+		*10111	*1011	2	0.290	0.438	1.706	1.875	70.7
31	be	+		*11000	*1100	3	1.107	1.142	1.889	1.932	45.7
32	be	+		*11011	*1101	3	1.203	1.245	1.472	1.479	29.0
30	be	+		*11111	*1111	5	0.152	0.152	0.190	0.259	30.3
AVG	sp						0.047	0.137	0.188	0.295	41.2
	be-+						0.402	0.460	0.829	0.903	34.1

AVG – average, sp – spruce; be-sp – beech-spruce; be(sp) – beech with admixed spruce; be – beech; The “basic classification group” was distinguished in 1980 using the TWINSPAN procedure. “Classification level change” determines the highest TWINSPAN classification level recorded after 1980. Variables v_1 to v_4 are calculated on the base of DCA ordination (see equation 1); E_{max} is the maximum Euclidean distance of a relevé with respect to the 1980 relevé

Spruce-beech stands – group *1011: PRP 8.

Beech stands – group *1100: PRP 31.

Beech stands – group *1101: PRP 32.

Beech stands – group *1111: PRP 30.

Plant communities in beech and mixed stands usually show medium to high change of the species composition, which is caused by higher species diversity in herb layer (Fig. 5).

Global trends in the plant community development

Species composition of vegetation on all 32 permanent research plots was significantly changing during the period 1980(1976)–2005, and some of the species completely disappeared (e.g. *Cicerbita alpina*, *Lamium maculatum*, *Phyteuma spicatum*, *Viola biflora*) or their apportionment decreased (e.g. *Blechnum spicant*, *Dentaria enneaphyllos*, *Homogyne alpina*). The number of species was greatly reduced especially in the moss layer.

To evaluate the trends in community development on studied research areas, the plots were

divided to spruce, beech and beech-spruce stands (Table 2). The first group was then subdivided into the plots with canopy decline and the others. Canopy decline was reported when the overall canopy cover of woody plants during the observation reduced to 25% or less.

Average canopy cover of woody plants in beech and mixed stands decreased only insignificantly – by 8% in 25 years. A similar tendency was observed in spruce stands where the decline did not occur. Canopy decline was marked at phytocenological relevés since 1985 and the average canopy cover decreased until 2000. The canopy regeneration was observed in the last sampling year 2005 (Table 3).

Total herb layer coverage in beech and mixed forests changed insignificantly. In spruce stands, an increase in the total herb layer coverage was observed. At the beginning of the observation this cover was higher in communities that later went through the wooden canopy decline.

The change in the number of species richness in the Krkonoše Mountains was significant. In beech and mixed forests, an average decrease of species was 27% of those present in 1980. In non-declining

Table 3. Trends in the phytocenose features according to the dominant woody plants on PRP 1–32 in the years 1976–2005, with relation to the occurrence of canopy decline

Type	Canopy decline	Year	N	Canopy closure D (%)			Total cover of E1 (%)			Richness E1			Richness E0		
				MIN	AVG	MAX	MIN	AVG	MAX	MIN	AVG	MAX	MIN	AVG	MAX
be	NO	1980	12	67	89	98	25	68	90	12	22.3	37	3	4.8	7
		1985	12	67	88	98	20	65	90	11	21.7	40	3	3.8	5
		1990	12	73	86	96	15	61	95	11	18.7	36	2	3.3	5
		1995	12	71	82	96	10	58	95	10	17.7	34	2	2.8	4
		2000	12	69	80	94	5	62	95	9	18.1	33	2	3.2	4
		2005	12	66	81	93	5	64	100	5	16.3	29	1	2.7	4
sp	NO	1976	3	53	60	65	70	75	80	10	12.7	14	10	14.3	21
		1980	11	42	60	75	55	78	100	12	17.2	24	8	11.6	20
		1985	11	40	56	75	60	82	100	11	16.8	23	7	10.0	17
		1990	11	40	55	80	65	86	100	9	14.9	20	3	6.9	11
		1995	11	35	54	80	70	89	100	8	14.1	20	3	5.7	10
		2000	11	40	54	75	75	91	100	10	15.9	22	3	6.4	9
sp	YES	1976	2	53	65	76	65	75	85	19	21.0	23	15	16.5	18
		1980	9	54	67	80	65	87	100	7	13.4	22	7	10.4	16
		1985	9	0	49	71	75	89	95	12	15.2	22	3	7.9	13
		1990	9	0	39	71	80	92	100	11	13.6	19	2	5.7	8
		1995	9	5	39	76	75	93	100	9	13.0	17	0	4.3	8
		2000	9	3	19	38	80	95	100	10	15.0	22	0	3.9	8
		2005	9	5	35	68	85	96	100	11	14.7	22	0	3.3	6

be – beech; sp – spruce; N – number of plots

spruce forests, this change was observed more as a fluctuation, with the minimum around 1995. In declining spruce stands, the changes in total number of species were not observed in herb layer; it does not necessarily mean that the species composition did not change (Table 3). These findings are comparable with another results (MATĚJKA 2007).

The most significant change was a reduced number of species in moss layer in all types of stands. In beech and mixed forests the average number of species dropped from 4.8 to 2.7 (44%), in non-declining spruce stands the number fell from 11.6 to 5.5 (53%), and in declining spruce stands it dropped from 10.4 to 3.3 (68%). Similarly, the reduction of total species diversity in these three groups was 36%, 31% and 43%, respectively. Lower percentage, compared to evaluation of the number of species, indicates that the reduction was recorded mainly at species with lower representation. Within the last group, the plots with no moss layer were observed (PRP 3 and 26) (Table 3).

CONCLUSION

The existing results of the phytocenoses study on 32 plots (beech, mixed and spruce stands) in the years 1980–2005 imply a relatively accelerated vegetation dynamics, mainly in the period of significant air-pollution impact of the range in the period from the 1970s to 1990s; and, on the contrary, important revitalisation processes after the year 2000. Great imission-ecological stress caused not only destruction or even dieback of wooden components of the ecosystems, but also significant changes in biodiversity of herbal and moss layer and the recess of lichens. Concerning the biodiversity, the most important change is a massive reduction of a number of species in moss layer (44 to 68%), at all types of stands (beech, mixed and spruce). Similarly, the overall species diversity at these three forest types decreased by 31% to 43%; the reduction concerned mainly the species with lower apportionment.

The greatest influence on differentiation of species composition was that of altitude and exposition of the plot. The influence of time on the decrease of species richness was also highly distinctive. The number of species that vanished was significantly higher than that of newly found species.

Some site changes were observed. This might be interpreted as indicators of shift of community to lower forest vegetation zones in the sense of global climate changes. A significant increase in the ratio of natural regeneration of beech was observed in the 6th and 7th forest altitudinal zone, too.

References

- AMBROS Z., MÍČHAL I. (1992): Phytoindication of changes in the natural forests of the Moravian-Silesian Beskids in the course of the years 1952–1986. *Ekologia*, **11**: 205–214.
- BOHN U., GOLLUB G., HETTEWER CH. (2000): Karte der natürlichen Vegetation Europas – 1:2 500 000. Bonn, Bundesamt für Naturschutz.
- BRANG P. (2005): Virgin forests as a knowledge source for central European silviculture: reality or myth? *Forest Snow and Landscape Research*, **79**: 19–32
- ELENBERG H. (1996): Vegetation Mitteleuropas mit den Alpen in ökologischer, dynamischer und historischer Sicht. Stuttgart, Verlag Eugen Ulmer: 1095.
- HÉDL R. (2004): Vegetation of beech forests in the Rychlebské Mountains, Czech Republic re-inspected after 60 years with assessment of environmental changes. *Plant Ecology*, **170**: 243–265.
- HILL M.O. (1979): TWINSpan – a FORTRAN Program for Arranging Multivariate Data in an Ordered Two Way Table by Classification of Individuals and Attributes. Ithaca, Cornell University: 48.
- MÁLEK J. (1970): Sphagnum fir-spruce forests (*Abieto-Piceetum sphagnosum*) in the Czech-Moravian highlands. *Vlastivědný sborník Vysočiny, Oddělení věd přírodních*, **6**: 61–70. (in Czech)
- MÁLEK J. (1973): Fir and spruce forests in the southern part of the Czech-Moravian highlands. *Lesnictví*, **19**: 37–58. (in Czech)
- MATĚJKA K. (2000): DBreleve Software Help. Available at <http://infodatasys.cz/software/dbreleve.htm> (accessed on December 16, 2009)
- MATĚJKA K. (2007): New methods of evaluation of community diversity (with examples of forest ecosystems). In: KRÍŽOVÁ E., UJHÁZY K. (eds): Dynamics, Stability and Diversity of Forest Ecosystems. Zvolen, Technical University in Zvolen, 161–170. (in Czech)
- MATĚJKA K. (2009): PlotOA software help (Plotting of ordination diagrams and cartograms). Available at: http://www.infodatasys.cz/software/hlp_PlotOA/PlotOA.htm (accessed on December 2, 2009)
- MATĚJKA K., VACEK S. (2007): Examples of forest ecosystems dynamics and their evaluation. In: KRÍŽOVÁ E., UJHÁZY K. (eds): Dynamics, stability and diversity of forest ecosystems. Zvolen, TU in Zvolen: 7–13. (in Czech)
- MATĚJKA K., VACEK S., PODRÁZSKÝ V. (2010): Development of forest soils in the Krkonoše Mts. in the period 1980–2009. *Journal of Forest Science*, **56**: 485–504.
- MATUSZKIEWICZ W., MATUSZKIEWICZ A. (1960): Pflanzensoziologische Untersuchungen der Waldgesellschaften des Riesengebirges. *Acta Societatis Botanicorum Poloniae*, **29**: 499–530.
- MATUSZKIEWICZ W., MATUSZKIEWICZ A. (1967): Groupments végétaux du Parc national Karkonosze (Sudètes occidentales). Partie 1. Groupments forestiers. *Prace Wrocławskie Towarzystwo Naukowe, serie B No. 135*: 1–100. (in Polish)
- MIKESKA M., VACEK S., PODRÁZSKÝ V. (2007): Concept of forestry typology in the bilateral biospheric reserve in the

- Krkonoše/Karkonosze mountains. *Opera Corcontica*, **44**: 471–483. (in Czech)
- MIKYŠKA R. (1968): Geobotanical map of the ČSSR. 1 Bohemia. Vegetation of the ČSSR. Praha, Academia: 204. (in Czech)
- MIKYŠKA R. (1972): Die Wälder der böhmischen mittleren Sudeten und ihrer Vorberge. *Rozpravy Československé akademie věd, Řada MPV*, **82**(3): 1–162.
- MOLDAN B., HAK, T. (2007): Environment in the Czech Republic: a positive and rapid change. *Environmental Science & Technology*, **41**: 359–362.
- MORAVEC J. (1974): Zusammensetzung und Verbreitung der *Dentario enneaphylli-Fagetum* in der Tschechoslowakei. *Folia geobotanica phytotaxonomica bohemoslovaca*, **9**: 113–152.
- PIŠTA F. (1972): Forest communities of the Šumava primeval forest. *Lesnictví*, **18**: 415–437. (in Czech)
- PIŠTA F. (1978): Spruce-beech stands in the southern part of the Šumava mountains. *Lesnictví*, **24**: 219–236. (in Czech)
- POKORNÝ P. (2002): Role of man in the development of Holocene vegetation in Central Bohemia. *Preslia*, **77**: 113–128.
- PODLASKI R. (2008): Dynamics in Central European near-natural *Abies-Fagus* forests: Does the mosaic-cycle approach provide an appropriate model? *Journal of Vegetation Science*, **19**: 173–182.
- PRŮŠA E. (1985): Die böhmischen und mährischen Urwälder – ihre Struktur und Ökologie. Praha, Academia Verlag: 578.
- SAMEK V. (1986): Problems and typisation of the monitoring of the influence of human activities on phytocenoses and the issues of experimental testing of phytotechnical measures. In: Preventive and remedial measures in endangered phytocenoses. Praha, ČSVTS – lesnická společnost, Botanický ústav ČSAV: 10–16. (in Czech)
- SOFRON J. (1981): Natural spruce forests in West and Southwest Bohemia. Study of the Czechoslovak Academy of Sciences, **7**: 127. (in Czech)
- STÖCKER G. (1968): Das *Anastrepto-Piceetum* im Harz und Riesengebirge (Krkonoše). *Opera Corcontica*, **5**: 135–155.
- SÝKORA T. (1967): Contribution to the study of mountain beech stands in the Krkonoše mountains. *Opera Corcontica*, **4**: 43–53. (in Czech)
- ŠAMONIL P., VRŠKA T. (2008): Long-term vegetation dynamics in the Šumava Mts. Natural spruce-fir-beech forests. *Plant Ecology*, **196**: 197–214.
- UJHÁZY K., KRIŽOVÁ E., VANČO M., FREŇÁKOVÁ E., ONDRUŠ M. (2005): Herb layer dynamics of primeval fir-beech forests in central Slovakia. In: CAMMARMOT B., HAMOR F.D. (eds): Natural Forests in the Temperate Zone of Europe – Values And Utilisation. Birmensdorf, Rakhiv, Swiss Federal Research Institute WSL, Carpatien Biosphere Reserve: 193–202.
- VACEK S. (1984): Phytocenoses analysis on the Strmá stráň location in the Krkonoše mountains. *Opera Corcontica*, **21**: 67–101. (in Czech)
- VACEK S., LEPŠ J. (1991): Analysis of the vegetation changes in the beech stands of the Orlické mountains. *Lesnictví*, **37**: 993–1007. (in Czech)
- VACEK S., LEPŠ J. (1992): Analysis of the vegetation changes in the spruce stands of the Orlické mountains. *Lesnictví*, **38**: 773–749. (in Czech)
- VACEK S., LEPŠ J. (1996): Spatial dynamics of forest decline: the role of neighbouring trees. *Journal of Vegetation Science*, **7**: 789–798.
- VACEK S., LEPŠ J. (1999): Vegetation dynamics in forest ecosystems in the Krkonoše mountains. *Zprávy České botanické společnosti*, **17**: 89–101. (in Czech)
- VACEK S., MATĚJKA K. (1999): The state of forest stands on permanent research plots in the Krkonoše Mts. in years 1976–1997. *Journal of Forest Science*, **45**: 291–315.
- VACEK S., MATĚJKA K. (2003): Vegetation changes in beech and spruce stands in the Orlické hory Mts. in 1951–2001. *Journal of Forest Science*, **49**: 445–473.
- VACEK S., PODRÁZSKÝ V., MATĚJKA K. (2006): Dynamics of the health status of forest stands and its prediction on research plots in the Šumava Mts. *Journal of Forest Science*, **52**: 457–473.
- VACEK S., PODRÁZSKÝ V., MIKESKA M., SIMON J. (2006): Forests and Ecosystems above the Upper Forest Boundary in the National Parks of the Krkonoše Mountains. *Kostelec nad Černými lesy, Lesnická práce*: 112. (in Czech)
- VACEK S., MATĚJKA K., SIMON J., MALÍK V., SCHWARZ O., PODRÁZSKÝ V., MINX T., TESAŘ V., ANDĚL P., JANKOVSKÝ, L., MIKESKA M. (2007): State and dynamics of forest ecosystems in the Krkonoše mountains under air pollution-induced stress. *Kostelec nad Černými lesy, Lesnická práce*: 216. (in Czech)
- VIEWEGH J. (1994): Changes in the vegetation in some nature reserves of the Moravian-Silesian Beskids. *Lesnictví- Forestry*, **40**: 523–536. (in Czech)

Received for publication April 24, 2010
Accepted after corrections June 26, 2010

Corresponding author:

Prof., RNDr. STANISLAV VACEK, DrSc., Česká zemědělská univerzita, Fakulta lesnická a dřevařská, Kamýcká 129, 165 21 Praha 6-Suchbát, Česká republika
tel.: + 420 224 382 870, fax: + 420 234 381 860, e-mail: vacekstanislav@fld.czu.cz
