

## Effect of *DGAT1* polymorphisms on the estimated breeding values of Czech Simmental sires

L. HANUSOVÁ<sup>1</sup>, A. MÍKOVÁ<sup>1</sup>, L. VEČEREK<sup>1</sup>, D. SCHROEFFELOVÁ<sup>2</sup>, V. ŘEHOUT<sup>1</sup>,  
L. TOTHOVÁ<sup>1</sup>, K. VERNEROVÁ<sup>1</sup>, B. HOSNEDLOVÁ<sup>1</sup>, J. ČÍTEK<sup>1</sup>

<sup>1</sup>Department of Genetics and Animal Breeding, Faculty of Agriculture,  
University of South Bohemia in České Budějovice, České Budějovice, Czech Republic

<sup>2</sup>Czech-Moravian Breeders Association, Hradištko, Czech Republic

**ABSTRACT:** The aim was to evaluate the effect of polymorphisms in the promoter and in the coding region of the *DGAT1* gene on the estimated breeding values (EBV) of Czech Simmental sires. The *K232A* polymorphism ( $n = 191$ ) in the coding region was genotyped by PCR/RFLP, and the *KU* and *SA* polymorphisms in the promoter ( $n = 203$ ) were identified in an automatic sequencer. In the *K232A* polymorphism, the frequency of the genotype *AA* (Alanine) was greater than that of the genotype *KA*, the homozygous genotype *KK* (Lysine) was not found. Similarly, the allele *A* predominated over the *K* allele (0.945 and 0.055). The EBV for milk performance have been assigned to the genotypes, and the associations quantified. For the *AA* genotype and *A* allele, positive association with EBV of milk yield and protein yield was found, and negative association with the breeding values of fat percentage and yield, and protein percentage, but only the value of fat content was found to be significant. The positive non-significant association of the *A* variant with the protein yield was caused by the high milk yield. In the *KU* polymorphism, the *CC* genotype was associated significantly with lower EBV for the fat percentage, both the *C* allele and the *CC* genotype were associated with higher EBV for milk yield, so both the fat and the protein yield were non-significantly increased. For the diplotypes *K232A/KU*, there was a significant association with the fat percentage. The *AACC* combination seemed to have some breeding potential. The *K232A* polymorphism explained maximum of 6.2% of EBV variability, the *KU* polymorphism of 4.4%, and the *SA* polymorphism of 4.2%. The diplotypes *K232A/KU* explained maximum of 7.4% of variability. The highest proportion of variability was explained for fat percentage. The results confirmed the important role of the BTA14 region in controlling milk performance.

**Keywords:** cattle; acyl-CoA diacylglycerol transferase1; *K232A*; promoter; milk performance; breeding; genotypes

### INTRODUCTION

The *DGAT1* gene encodes the DGAT1 enzyme, which catalyzes the final step of triglyceride synthesis (Sanders et al. 2006).

Several studies in cattle have described a quantitative trait locus (QTL) with impact on milk production traits, and on milk fat percentage in particular, in the 3-cM region in the centromeric part of *Bos taurus* autosome 14 (BTA14) (Riquet et al. 1999; Looft et al. 2001). Grisart et al. (2002,

2004) and Winter et al. (2002, 2004) have identified a nonconservative dinucleotide substitution (*K232A*) in the acyl-CoA diacylglycerol acyltransferase1 (*DGAT1*) gene at position 10433 and 10434 in exon number 8 as the most likely mechanism underlying the QTL on this chromosome.

However, apparent differences in the effect observed between families and across populations could not be fully explained by this diallelic polymorphism alone. It was reported that genetic variation additional to the DGAT1 *K232A* mutation affecting milk fat

Supported by the Grant Agency of the University of South Bohemia in České Budějovice (GAJU) (Projects No. 011/2013/Z and No. 022/2013/Z).

content should be present in the same QTL (Bennewitz et al. 2004; Kuhn et al. 2004). Winter et al. (2002) considered alleles of the *DGAT1* promoter region, which comprise a variable number of tandem repeats (VNTR), as likely candidates. In the German Holstein population, Kuhn et al. (2004) described 5 alleles at a VNTR polymorphism in the *DGAT1* promoter, which showed an effect on fat content additional to the *DGAT1* K232A mutation. The most frequent allele in the *DGAT1* promoter VNTR was allele 3. Multiple regression analysis of the *DGAT1* promoter VNTR alleles in sons of genotype *DGAT1* 232A/232A revealed that the allele substitution effect on the milk fat content of allele 5 was higher than of all the other alleles. The regression analysis of the *DGAT1* promoter VNTR alleles revealed significant effects for the allele 5 enhancing the milk fat content percentage as well as the milk protein percentage but decreasing the milk yield and milk protein yield (Kuhn et al. 2004).

Sanders et al. (2006) reported 6 alleles found in the *DGAT1* promoter VNTR in the Angeln population, which were denoted as VNTR alleles *A*, *B*, *C*, *D*, *E*, and *F*, respectively. The allele *F* was present in two unrelated daughters only. The most frequent *DGAT1* allele was *E*. In their study, they observed that the VNTR allele *E* showed significant effects for some milk production indicators compared with all other alleles in the *DGAT1* promoter. The same results were reported by Kuhn et al. (2004) for the *DGAT1* VNTR allele 5. However, in contrast to Kuhn et al. (2004), the VNTR allele *E* was mainly linked to the *K* variant at *DGAT1* K232A (Sanders et al. 2006), whereas the *DGAT1* VNTR allele 5 showed up with the *A* variant in the German Holstein Friesian population (Kuhn et al. 2004). The VNTR allele *E* of Sanders et al. (2006) probably corresponds to the *DGAT1* VNTR allele 5 of Kuhn et al. (2004).

In the paper, we report on our analysis of the effect of the genotypes and alleles in the promoter, and in the coding region at nucleotide positions 10433 and 10434 of the *DGAT1* gene on the estimated breeding values (EBV) of Czech Simmental sires.

## MATERIAL AND METHODS

The analyzed group was made of the Czech Simmental sires born in the period 2000–2004, the sires were selected randomly. The DNA was isolated from frozen sperm. The respective parts of the

*DGAT1* locus carrying the polymorphisms studied were amplified in the PCR. The primer sequences and PCR conditions were as in Kuhn et al. (2004) and Sanders et al. (2006) for the polymorphisms in the promoter and as in Winter et al. (2002) for the polymorphisms at nucleotide positions 10433 and 10434 of the *DGAT1* gene (*K232A* polymorphism). In promoter, the polymorphisms *SA* and *KU* were studied (Kuhn et al. 2004; Sanders et al. 2006). The PCR was carried out on the Biometra TGradient Thermocycler (Biometra GmbH, Göttingen, Germany). The VNTR polymorphisms *KU* and *SA* in the promoter of the *DGAT1* gene were distinguished in the automatic sequencer ABI PRISM<sup>®</sup>310 Genetic Analyzer (Applied Biosystems, Foster City, USA). The polymorphism *K232A* causing *K* to *A* substitution was distinguished by a restriction fragment length polymorphism using the restriction endonuclease *CfrI* (PCR/RFLP method). An alternative genotyping method was published by Abdolmohammadi et al. (2011). For the *K232A* polymorphism, 191 sires were genotyped, and 203 sires for the polymorphisms in the promoter.

The actual breeding values estimated in 2011 (Plemdat; www.plemdat.cz) have been assigned to the genotypes, and the associations quantified. The genotypes with low frequency were left out in the association analysis. So, for the *K232A* polymorphism, 180 sires were involved, namely purebred Simmentals ( $n = 54$ ), crossbreds of Simmental with Holstein and Ayrshire with the proportion of Simmental above 75% ( $n = 93$ ), crossbreds with the proportion of Simmental of 50–74% ( $n = 21$ ), and purebred Montbeliardes ( $n = 12$ ). For the promoter polymorphisms, the counts were the same, just the number of purebred Simmentals was 55, and so the total number of sires was 181.

We analyzed the relation between the detected genotypes and estimated breeding values for the milk production traits: milk yield (kg), relative breeding value for milk yield, fat content (%), fat yield (kg), protein content (%), relative breeding value for protein content, protein yield (kg), and relative breeding value for protein yield. The loci were analyzed independently. Generally, the using of daughter yield deviations (DYD) is preferred to EBV, because EBV contain information from other relatives than the bull's daughters. However, the large number of daughters make the difference between the DYD and EBV negligible (Viitala et al. 2006). The reliabilities of EBV were equable, which enabled their direct comparison (Table 1).

Table 1. Reliability of estimated breeding values (EBV) of Czech Simmental sires with different *DGAT1* genotypes

| Polymorphism | Genotype       | Reliability of EBV |
|--------------|----------------|--------------------|
| <i>K232A</i> | <i>AA</i>      | 89 ± 3.36          |
|              | <i>KA</i>      | 89 ± 3.08          |
| <i>KU</i>    | <i>CC</i>      | 89 ± 3.69          |
|              | <i>CD</i>      | 89 ± 2.89          |
|              | <i>CE</i>      | 89 ± 4.25          |
|              | <i>DD</i>      | 90 ± 2.64          |
|              | <i>DE</i>      | 88 ± 2.14          |
| <i>SA</i>    | <i>221/221</i> | 89 ± 3.73          |
|              | <i>221/239</i> | 89 ± 2.87          |
|              | <i>221/256</i> | 89 ± 4.25          |
|              | <i>239/239</i> | 90 ± 2.65          |
|              | <i>239/256</i> | 88 ± 2.14          |

The statistical evaluation was based on the model equation:

$$EBV = \mu + G_i + e_{ijk}$$

where:

EBV = estimated breeding value for partial milk production parameter

$\mu$  = overall mean

$G_i$  = fixed effect of genotype/allele of polymorphic sites in the promoter, or in *K232A*

$e_{ijk}$  = residual effect

Assumptions for the Analysis of Variance (ANOVA) were tested by using the Bartlett test of homogeneity of variance. The estimated breeding values were evaluated by ANOVA depending on the genotype on the *DGAT1* locus; similarly, the differences between the alleles were quantified. The differences were evaluated at the significance levels of  $P < 0.05^*$ , and  $P < 0.01^{**}$ . The software STATISTICA (Version 10, 2013) and the ANOVA/MANOVA procedure were used. The Hardy-Weinberg equilibrium was tested by the  $\chi^2$  test.

## RESULTS AND DISCUSSION

**Genotypes and alleles frequencies.** In the *K232A* polymorphism, the genotype *AA* prevailed substantially over the *KA*, while the homozygous genotype *KK* was not found. Similarly, the allele *A* predominated over the *K* allele in the group analyzed (0.945 and 0.055, respectively) (Table 2). This is different from frequencies found in German Holsteins in our own previous study (Hradecka

et al. 2008), where the frequencies of 0.660 and 0.340 were found, but the population mentioned was upgraded with the Jersey breed. Weller et al. (2003) in Israeli Holstein cows gave a frequency of *K* allele of 0.09, in sires of 0.16. By contrast, Thaller et al. (2003) gave the frequency of the allelic variant coding for *K* in the German Holstein of 0.548. As in Grisart et al. (2002), *K* is probably the ancestral allele; its frequency is indirectly influenced by selection and decreased while selecting for high milk yield. The low frequency of the *K* allele found in this paper implies indirect selection as a consequence of the preference of the protein yield in Czech Simmentals in the last decades. Moreover, the Czech Simmental was upgraded by using Ayrshire and Red Holstein cattle with the aim of bettering the milk performance, and the crossing could have influenced the

Table 2. Genotype and allele frequencies of *DGAT1* polymorphisms

| Genotype            | <i>n</i> | Frequency | $c^2$              | Allele     | Frequency |
|---------------------|----------|-----------|--------------------|------------|-----------|
| <b><i>K232A</i></b> |          |           |                    |            |           |
| <i>AA</i>           | 170      | 0.890     |                    | <i>A</i>   | 0.945     |
| <i>KA</i>           | 21       | 0.110     | 0.662 <sup>a</sup> | <i>K</i>   | 0.055     |
| <i>KK</i>           | 0        | 0.00      |                    |            |           |
| <b><i>KU</i></b>    |          |           |                    |            |           |
| <i>BD</i>           | 2        | 0.010     |                    | <i>B</i>   | 0.005     |
| <i>CC</i>           | 45       | 0.222     |                    | <i>C</i>   | 0.480     |
| <i>CD</i>           | 65       | 0.320     |                    | <i>D</i>   | 0.318     |
| <i>CE</i>           | 39       | 0.192     |                    | <i>E</i>   | 0.192     |
| <i>CF</i>           | 1        | 0.005     | 3.750 <sup>a</sup> | <i>F</i>   | 0.005     |
| <i>DD</i>           | 20       | 0.099     |                    |            |           |
| <i>DE</i>           | 21       | 0.103     |                    |            |           |
| <i>DF</i>           | 1        | 0.005     |                    |            |           |
| <i>EE</i>           | 9        | 0.044     |                    |            |           |
| <b><i>SA</i></b>    |          |           |                    |            |           |
| <i>205/239</i>      | 2        | 0.010     |                    | <i>205</i> | 0.005     |
| <i>221/221</i>      | 44       | 0.217     |                    | <i>221</i> | 0.478     |
| <i>221/239</i>      | 66       | 0.325     |                    | <i>239</i> | 0.320     |
| <i>221/256</i>      | 39       | 0.192     |                    | <i>256</i> | 0.192     |
| <i>221/273</i>      | 1        | 0.005     | 4.305 <sup>a</sup> | <i>273</i> | 0.005     |
| <i>239/239</i>      | 20       | 0.099     |                    |            |           |
| <i>239/256</i>      | 21       | 0.103     |                    |            |           |
| <i>239/273</i>      | 1        | 0.005     |                    |            |           |
| <i>256/256</i>      | 9        | 0.044     |                    |            |           |

<sup>a</sup>non significant

$c^2$  = test for evaluation of average variation from expected frequencies by Hardy-Weinberg equilibrium

Table 3. Estimated breeding values according to genotypes and alleles of *DGATI K232A* polymorphism

| Genotype/allele | n   | Trait      |          |                           |                |               |                 |              |                |     |
|-----------------|-----|------------|----------|---------------------------|----------------|---------------|-----------------|--------------|----------------|-----|
|                 |     | milk yield |          | fat yield (kg)            |                | protein yield |                 |              |                |     |
|                 |     | (kg)       | RBV      | fat (%)                   | fat yield (kg) | protein (%)   | protein (%) RBV | (kg)         | protein yield  | RBV |
| AA              | 161 | 166 ± 490  | 109 ± 12 | -0.01 ± 0.20 <sup>a</sup> | 6.09 ± 18.81   | -0.01 ± 0.12  | 98.66 ± 11.42   | 5.19 ± 15.18 | 109.08 ± 12.11 |     |
| KA              | 19  | -4 ± 618   | 105 ± 15 | 0.18 ± 0.30 <sup>a</sup>  | 9.21 ± 20.47   | 0.01 ± 0.12   | 100.05 ± 11.55  | 0.16 ± 19.92 | 105.21 ± 15.63 |     |
| F-test          |     | 1.93       | 1.77     | 4.32*                     | 0.46           | 0.35          | 0.25            | 1.74         | 1.63           |     |
| R <sup>2</sup>  |     | 0.005      | 0.004    | 0.062                     | -0.003         | -0.003        | -0.004          | 0.004        | 0.003          |     |
| A               | 341 | 157 ± 498  | 109 ± 13 | 0.00 ± 0.21 <sup>b</sup>  | 6.27 ± 18.87   | 0.00 ± 0.12   | 98.74 ± 11.40   | 4.91 ± 15.47 | 108.87 ± 12.32 |     |
| K               | 19  | -4 ± 618   | 105 ± 15 | 0.18 ± 0.30 <sup>b</sup>  | 9.21 ± 20.47   | 0.01 ± 0.12   | 100.05 ± 11.55  | 0.16 ± 19.92 | 105.21 ± 15.63 |     |
| R <sup>2</sup>  |     | 0.002      | 0.002    | 0.029                     | -0.002         | -0.002        | -0.002          | 0.002        | 0.001          |     |

RBV = relative breeding value (%)

<sup>a</sup>values in one column with identical letters differ significantly ( $P < 0.05$ )\* $P < 0.05$ Table 4. Estimated breeding values according to genotypes and alleles of *DGATI KU* polymorphism

| Genotype/allele | n   | Trait      |          |                           |                |               |                             |              |                |     |
|-----------------|-----|------------|----------|---------------------------|----------------|---------------|-----------------------------|--------------|----------------|-----|
|                 |     | milk yield |          | fat yield (kg)            |                | protein yield |                             |              |                |     |
|                 |     | (kg)       | RBV      | fat (%)                   | fat yield (kg) | protein (%)   | protein (%) RBV             | (kg)         | protein yield  | RBV |
| CC              | 44  | 306 ± 418  | 113 ± 11 | -0.06 ± 0.20 <sup>a</sup> | 8.86 ± 18.28   | -0.04 ± 0.11  | 95.25 ± 10.61 <sup>Aa</sup> | 8.55 ± 13.76 | 112.00 ± 10.87 |     |
| CD              | 62  | 101 ± 584  | 107 ± 15 | 0.07 ± 0.24 <sup>a</sup>  | 7.48 ± 20.74   | 0.02 ± 0.13   | 101.06 ± 11.45 <sup>A</sup> | 4.19 ± 18.02 | 108.02 ± 14.14 |     |
| CE              | 36  | 107 ± 481  | 108 ± 12 | 0.04 ± 0.23               | 6.27 ± 14.77   | 0.02 ± 0.14   | 101.06 ± 12.86 <sup>a</sup> | 4.31 ± 14.93 | 108.53 ± 11.75 |     |
| DD              | 19  | 223 ± 483  | 111 ± 12 | -0.03 ± 0.21              | 7.79 ± 19.65   | -0.03 ± 0.08  | 95.89 ± 7.67                | 5.31 ± 14.57 | 109.63 ± 11.57 |     |
| DE              | 20  | 105 ± 342  | 108 ± 8  | -0.06 ± 0.19              | 0.75 ± 12.64   | -0.02 ± 0.13  | 97.20 ± 12.28               | 2.35 ± 10.33 | 107.00 ± 8.25  |     |
| F-test          |     | 1.21       | 1.36     | 2.65*                     | 0.60           | 1.90          | 2.08                        | 0.76         | 1.00           |     |
| R <sup>2</sup>  |     | 0.009      | 0.011    | 0.044                     | -0.006         | 0.027         | 0.032                       | -0.005       | -0.001         |     |
| C               | 186 | 199 ± 498  | 109 ± 13 | 0.00 ± 0.23               | 7.90 ± 18.43   | -0.01 ± 0.13  | 98.31 ± 11.63               | 6.27 ± 15.56 | 110.00 ± 12.27 |     |
| D               | 120 | 140 ± 517  | 108 ± 13 | 0.02 ± 0.22               | 6.46 ± 19.22   | 0.00 ± 0.12   | 98.78 ± 10.72               | 4.24 ± 15.77 | 108.36 ± 12.44 |     |
| E               | 70  | 98 ± 412   | 107 ± 11 | 0.02 ± 0.22               | 5.10 ± 14.78   | 0.00 ± 0.13   | 98.86 ± 12.27               | 3.00 ± 13.32 | 107.16 ± 11.10 |     |
| R <sup>2</sup>  |     | 0.001      | 0.003    | -0.004                    | -0.002         | -0.005        | -0.005                      | 0.002        | 0.003          |     |

RBV = relative breeding value (%)

<sup>a</sup>values in one column with identical letters differ significantly ( $P < 0.05$ )<sup>A</sup>values in one column with identical letters differ significantly ( $P < 0.01$ )\* $P < 0.05$

frequencies in the *DGAT1* locus. Spelman et al. (2002) recorded variable frequencies in Holstein sires based on the origin of the genetic material. The frequency was higher in the homebred (New Zealander) population, where it was similar to the Jersey breed, and it indicates changes due to indirect selection. They report a low frequency of 0.22 in Ayrshire. In Montbeliarde, the *A* variant is practically fixed (Gautier et al. 2007), and this is concordant with our frequencies in phylogenetic related Simmental cattle.

In the promoter region, two possible polymorphisms, *SA* and *KU*, were studied. In the *SA* polymorphism, the most frequent allele 221 had a frequency of 0.478 and the least frequent alleles 205 and 273 of 0.005. Using the same allele designation as Sanders et al. (2006), the most frequent allele in our study was allele *B* and the least frequent were alleles *A* and *E*. These results are in contrast to results of the authors mentioned, who found allele *E* to be the most frequent. The allele with the lowest frequency was allele *A* in both studies. In the *KU* polymorphism, we studied 5 alleles. The most frequent was allele *C* (0.480) and the least frequent were alleles *B* and *F* (0.005). Our results are also not completely concordant with Kuhn et al. (2004). They described allele 3 (*D*) as the most frequent, and allele 1 (*B*) as the least frequent. In our paper, the frequencies as compared among the purebred Simmentals, purebred Montbeliarde, and crosses were not significantly different both in *K232A* and promoter polymorphisms.

**Association analysis of the *K232A* polymorphism.** For the coding region of the *DGAT1* gene (Table 3), significant differences in estimated breeding values of fat content both for genotypes and alleles were found; the *K* variant was associated with higher values. The results are in accordance with those previously found in German Holsteins (Citek et al. 2007). However, also in Holsteins, the estimated breeding values of milk yield, fat yield, protein yield, and protein content were significant. In this paper, the trend of estimated breeding values for genotypes and alleles was the same, i.e. the homozygous *AA* genotype and *A* allele were associated with higher milk yield, protein yield, and with lower fat percentage, fat yield, and protein percentage, but significance was found only in fat content. The positive non-significant link of the *A* variant on protein yield was caused by high milk yield. The highest portion of variability was explained in EBV for fat content (6.20% by

genotype; 2.90% by allele), the other values did not exceed 1%.

Because there was a high breeding value for milk yield and a negative value for fat content in the homozygotes for the Alanine variant, the *K232A* polymorphism may contribute to the negative correlations between the traits. The group has been in Hardy-Weinberg equilibrium, that implies that the selection on breeding value for protein yield did not importantly affect frequencies on the *K232A* polymorphism. The possible reasons may be seen in the rational selection on the fat and protein yield in the breeding programme, which could steady the frequencies; and also in the fact that the *DGAT1* itself does not directly influence the protein synthesis (compare Citek et al. 2007). Our results correspond well with many other authors (Grisart et al. 2002; Weller et al. 2003; Sanders et al. 2006), and are applicable in breeding practice.

**Association analysis of the promoter polymorphisms.** The *CC* genotype in the *KU* polymorphism was associated significantly with lower EBV for fat percentage compared with *CD*, and with the protein percentage compared with *CD* and *CE* genotypes, but non-significantly with the others (Table 4). But both *C* allele and *CC* genotype were linked to higher EBV for milk yield, so the fat and protein yield were increased non-significantly. For the *EE* genotype, the lowest EBV for milk yield of 65 kg was stated, the highest increase of fat percentage of 0.09%, and low protein yield of 0.57 kg; this was similar to the allele *E* effect, although the number of animals was very low ( $n = 7$ ), and thus the data are not shown in Table 4. Again, the highest portion of variability was explained in EBV for fat content, 4.40% for genotype. Evidently, the *SA* polymorphism (Table 5) is the same as that of *KU*, as noted by other authors (Kuhn et al. 2004, Sanders et al. 2006).

The relationship between the combined genotypes of *K232A* and *KU* polymorphisms and the estimated breeding values was quantified (Table 6). The significant association with the fat percentage was noted. At the *KU* polymorphism, the combination of the *AA* genotype in *K232A* and the *CC*, *DE*, and *DD* in *KU* had the negative link, whereas *LA/CD* had the positive. For fat content, the combined genotype explained 7.4% of variability. The *AA/CC* combination seems to have some potential, as it was related with the higher estimated breeding values of the milk, fat and protein yield, but the differences were not significant for the traits mentioned.

Table 5. Estimated breeding values according to genotypes and alleles of *DGATI SA* polymorphism

| Genotype/<br>allele   | <i>n</i> | Trait              |                           |                |              |                       |                      |                |
|-----------------------|----------|--------------------|---------------------------|----------------|--------------|-----------------------|----------------------|----------------|
|                       |          | milk yield<br>(kg) | fat (%)                   | fat yield (kg) | protein (%)  | protein yield<br>(kg) | protein yield<br>RBV |                |
| 221/221               | 43       | 308 ± 423          | -0.06 ± 0.20 <sup>a</sup> | 9.19 ± 18.37   | -0.04 ± 0.11 | 95.53 ± 10.56         | 8.77 ± 13.84         | 112.12 ± 10.92 |
| 221/239               | 63       | 103 ± 579          | 0.06 ± 0.23 <sup>a</sup>  | 7.29 ± 20.63   | 0.02 ± 0.13  | 100.78 ± 11.58        | 4.11 ± 17.89         | 107.95 ± 14.03 |
| 221/256               | 36       | 107 ± 481          | 0.04 ± 0.23               | 6.28 ± 14.77   | 0.02 ± 0.14  | 101.06 ± 12.86        | 4.31 ± 14.93         | 108.53 ± 11.75 |
| 239/239               | 19       | 224 ± 483          | -0.03 ± 0.21              | 7.79 ± 19.65   | -0.03 ± 0.08 | 95.89 ± 7.67          | 5.32 ± 14.57         | 109.63 ± 11.57 |
| 239/256               | 20       | 105 ± 342          | -0.06 ± 0.19              | 0.75 ± 12.64   | -0.02 ± 0.13 | 97.20 ± 12.28         | 2.35 ± 10.33         | 107.00 ± 8.25  |
| <i>F</i> -test        |          | 1.21               | 2.57*                     | 0.63           | 1.65         | 1.81                  | 0.82                 | 1.07           |
| <i>R</i> <sup>2</sup> |          | 0.009              | 0.042                     | -0.005         | 0.020        | 0.024                 | -0.003               | 0.000          |
| 221                   | 185      | 199 ± 499          | 0.00 ± 0.23               | 7.97 ± 18.45   | -0.01 ± 0.13 | 98.39 ± 11.61         | 6.31 ± 15.59         | 110.03 ± 12.30 |
| 239                   | 121      | 141 ± 515          | 0.01 ± 0.22               | 6.36 ± 19.17   | -0.01 ± 0.12 | 98.65 ± 10.77         | 4.20 ± 15.71         | 108.32 ± 12.39 |
| 256                   | 70       | 98 ± 412           | 0.02 ± 0.22               | 5.10 ± 14.78   | 0.00 ± 0.13  | 98.86 ± 12.27         | 3.00 ± 13.32         | 107.16 ± 11.10 |
| <i>R</i> <sup>2</sup> |          | 0.001              | -0.004                    | -0.001         | -0.005       | -0.005                | 0.002                | 0.004          |

RBV = relative breeding value (%)

<sup>a</sup>values in one column with identical letters differ significantly ( $P < 0.05$ ); \* $P < 0.05$ Table 6. Estimated breeding values according to combined genotypes of *K232A* and *KU* polymorphisms

| Genotype              | <i>n</i> | Trait              |                            |                |              |                       |                      |                |
|-----------------------|----------|--------------------|----------------------------|----------------|--------------|-----------------------|----------------------|----------------|
|                       |          | milk yield<br>(kg) | fat (%)                    | fat yield (kg) | protein (%)  | protein yield<br>(kg) | protein yield<br>RBV |                |
| AA/CC                 | 44       | 306 ± 418          | -0.06 ± 0.20 <sup>A</sup>  | 8.86 ± 18.28   | -0.04 ± 0.11 | 95.25 ± 10.61         | 8.55 ± 13.76         | 112.00 ± 10.87 |
| AA/CE                 | 32       | 140 ± 474          | 0.02 ± 0.21                | 6.63 ± 15.52   | 0.01 ± 0.14  | 100.28 ± 12.98        | 5.03 ± 14.89         | 109.06 ± 11.72 |
| AA/CD                 | 53       | 125 ± 552          | 0.04 ± 0.22                | 6.87 ± 19.64   | 0.02 ± 0.13  | 101.04 ± 11.56        | 4.96 ± 16.56         | 108.58 ± 13.01 |
| AA/DE                 | 18       | 74 ± 346           | -0.05 ± 0.20 <sup>b</sup>  | 0.11 ± 13.20   | -0.01 ± 0.13 | 98.50 ± 12.00         | 1.94 ± 10.84         | 106.67 ± 8.66  |
| AA/DD                 | 16       | 263 ± 461          | -0.06 ± 0.15 <sup>c</sup>  | 8.19 ± 20.45   | -0.03 ± 0.09 | 96.18 ± 8.10          | 6.69 ± 13.50         | 110.81 ± 10.73 |
| AA/EE                 | 6        | 63 ± 372           | 0.03 ± 0.17                | 4.50 ± 16.21   | -0.06 ± 0.11 | 93.67 ± 10.25         | -0.33 ± 14.80        | 102.83 ± 14.39 |
| KA/CE                 | 4        | -153 ± 524         | 0.19 ± 0.30                | 3.50 ± 6.76    | 0.09 ± 0.13  | 107.25 ± 11.44        | -1.50 ± 16.13        | 104.25 ± 12.84 |
| KA/CD                 | 9        | -38 ± 771          | 0.24 ± 0.27 <sup>Abc</sup> | 11.11 ± 27.51  | 0.02 ± 0.12  | 101.22 ± 11.40        | -0.33 ± 25.83        | 104.67 ± 20.24 |
| KA/DD                 | 3        | 17 ± 658           | 0.10 ± 0.45                | 5.67 ± 18.15   | -0.05 ± 0.06 | 94.33 ± 5.78          | -2.00 ± 21.17        | 103.33 ± 16.44 |
| <i>F</i> -test        |          | 1.126              | 2.845*                     | 0.472          | 1.417        | 1.521                 | 0.788                | 0.496          |
| <i>R</i> <sup>2</sup> |          | 0.005              | 0.074                      | -0.024         | 0.018        | 0.022                 | -0.009               | -0.002         |

RBV = relative breeding value (%)

values in one column with identical letters differ significantly <sup>A</sup>( $P < 0.01$ ), <sup>b</sup>( $P < 0.05$ )\* $P < 0.01$

Table 7. Estimated breeding values for different *KU* alleles in 232A/232A homozygous sires

| <i>KU</i> allele <sup>1</sup> | <i>n</i> | Trait      |          |              |                |              |                 |               |                |
|-------------------------------|----------|------------|----------|--------------|----------------|--------------|-----------------|---------------|----------------|
|                               |          | milk yield |          | fat (%)      | fat yield (kg) | protein (%)  | protein (%) RBV | protein yield |                |
|                               |          | kg         | RBV      |              |                |              |                 | kg            | RBV            |
| <i>C</i>                      | 173      | 220 ± 477  | 110 ± 12 | -0.02 ± 0.21 | 7.83 ± 18.11   | -0.01 ± 0.13 | 97.95 ± 11.61   | 6.80 ± 14.86  | 110.41 ± 11.73 |
| <i>D</i>                      | 103      | 159 ± 493  | 109 ± 12 | -0.01 ± 0.20 | 6.10 ± 18.90   | -0.01 ± 0.12 | 99.08 ± 10.78   | 4.97 ± 14.68  | 108.94 ± 11.60 |
| <i>E</i>                      | 62       | 106 ± 414  | 107 ± 11 | 0.00 ± 0.20  | 4.32 ± 15      | -0.01 ± 0.13 | 98.48 ± 12.21   | 3.10 ± 13.64  | 107.16 ± 11.41 |
| <i>R</i> <sup>2</sup>         |          | 0.020      | 0.025    | 0.033        | 0.005          | 0.030        | 0.037           | 0.010         | 0.019          |

RBV = relative breeding value (%)

<sup>1</sup>differences among alleles were not significant

Finally, the effect of the alleles at the *KU* site within the group of 232A/232A sires was evaluated (Table 7). The differences were not significant; the allele *E* showed to be linked to lower EBV of milk, fat, and protein yield, and the relation to the fat and protein content was neutral. This is in concordance with Sanders et al. (2006) who found the interallelic differences to be insignificant but the influence of haplotypes to be significant.

Together with other markers analyzed in Czech Simmental (Boleckova et al. 2012), the *DGATI* locus should be studied intensively, as the breeding potential is high. Then, the genomic approach (Schopen et al. 2009; Pribyl et al. 2010, 2012, 2013; Matejickova et al. 2013; Szyda et al. 2013) could be completed by the known major genes concerning milk performance and by the biometric approach (Sigl et al. 2012; Meszaros et al. 2013; Zavadilova and Stipkova 2013; Zavadilova and Zink 2013). When different effects depending on the breed are found, e.g. Suchocki et al. (2010) noted a stronger effect in Jersey than in Holstein, the gene polymorphisms should be evaluated regarding the breed. In the paper, we have analyzed the dual-purpose Czech Simmental cattle, as in most cases the dairy breeds, predominantly Holstein, are in the spotlight.

## CONCLUSION

Concludingly, in the *K232A* polymorphism the *K* allele was associated significantly with higher estimated breeding values of fat content and *vice versa* for the *AA* genotype and *A* allele. The *CC* genotype in the *KU* polymorphism was associated significantly with lower EBV for fat and protein percentage. Both *C* allele and *CC* genotype were linked to higher EBV for milk yield, so the fat and protein yield were linked positively, but non-signif-

icantly. The combined genotypes of *K232A* and *KU* polymorphisms and the EBV for the fat percentage were linked significantly, and the combination *AA/CC* may have some breeding potential. The *K232A* polymorphism explained at the most of 6.2% of variability of estimated breeding value, the *KU* polymorphism of 4.4%, and the *SA* polymorphism of 4.2%. The combined genotypes *K232A* and *KU* explained at the most of 7.4% of variability. In all polymorphisms, the highest proportion of EBV variability was explained for fat percentage. Definitely, there is promising that repeated analyses in different breeds show the important role of the BTA14 region in controlling milk performance.

**Acknowledgement.** The authors thank John P. McAvoy for the language revision.

## REFERENCES

- Abdalmohammadi A., Atashi H., Zamani P., Bottema C. (2011): High resolution melting as an alternative method to genotype diacylglycerol o-acyltransferase 1 (*DGATI*) *K232A* polymorphism in cattle. Czech Journal of Animal Science, 56, 370–376.
- Bennewitz J., Reinsch N., Paul S., Looft C., Kaupe B., Weimann C., Erhardt G., Thaller G., Kuhn C., Schwerin M., Thomsen H., Reinhardt F., Reents R., Kalm E. (2004): The *DGATI K232A* mutation is not solely responsible for the milk production quantitative trait locus on the bovine chromosome 14. Journal of Dairy Science, 87, 431–442.
- Boleckova J., Matejickova J., Stipkova M., Kyselova J., Barton L. (2012): The association of five polymorphisms with milk production traits in Czech Fleckvieh cattle. Czech Journal of Animal Science, 57, 45–53.
- Citek J., Rehout V., Hradecka E., Vecerek L., Panicke L. (2007): The breeding values of German Holstein sires and the *DGATI* polymorphism. Archiv für Tierzucht, 50, 136–146.

- Gautier M., Capitan A., Fritz S., Eggen A., Boichard D., Druet T. (2007): Characterization of the *DGATI K232A* and variable number of tandem repeat polymorphisms in French dairy cattle. *Journal of Dairy Science*, 90, 2980–2988.
- Grisart B., Coppieters W., Farnir F., Karim L., Ford C., Berzi P., Cambisano N., Mni M., Reid S., Simon P., Spelman R., Georges M., Snell R. (2002): Positional candidate cloning of a QTL in dairy cattle: identification of a missense mutation in the bovine *DGATI* gene with major effect on milk yield and composition. *Genome Research*, 12, 222–231.
- Grisart B., Farnir F., Karim L., Cambisano N., Kim J.J., Kvasz A., Mni M., Simon P., Frere J.M., Coppieters W., Georges M. (2004): Genetic and functional confirmation of the causality of the *DGATI K232A* quantitative trait nucleotide in affecting milk yield and composition. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 2398–2403.
- Hradecka E., Citek J., Panicke L., Rehout V., Hanusova L. (2008): The relation of *GHI*, *GHR* and *DGATI* polymorphisms with estimated breeding values for milk production traits of German Holstein sires. *Czech Journal of Animal Science*, 53, 238–245.
- Kuhn C., Thaller G., Winter A., Bininda-Edmonds O.R.P., Kaupe B., Erhardt G., Bennewitz J., Schwerin M., Fries R. (2004): Evidence for multiple alleles at the *DGATI* locus better explains a quantitative trait locus with major effect on milk fat content in cattle. *Genetics*, 167, 1873–1881.
- Looft C., Reinsch N., Karall-Albrecht C., Paul S., Brink M., Thomsen H., Brockmann G., Kuhn C., Schwerin M., Kalm E. (2001): A mammary gland EST showing linkage disequilibrium to a milk production QTL on bovine chromosome 14. *Mammalian Genome*, 12, 646–650.
- Matejickova J., Stipkova M., Sahana G., Kott T., Kyselova J., Matejicek A., Kottova B., Sefrova J., Krejcová M., Melcova S. (2013): QTL mapping for production traits in Czech Fleckvieh cattle. *Czech Journal of Animal Science*, 58, 396–403.
- Meszáros G., Kadlecik O., Kasarda R., Solkner J. (2013): Analysis of longevity in the Slovak Pinzgau population – extension to the animal model. *Czech Journal of Animal Science*, 58, 289–295.
- Pribyl J., Rehout V., Citek J., Pribylova J. (2010): Genetic evaluation of dairy cattle using a simple heritable genetic ground. *Journal of the Science of Food and Agriculture*, 90, 1765–1773.
- Pribyl J., Haman J., Kott T., Pribylova J., Simeckova M., Vostry L., Zavadilova L., Cermak V., Ruzicka Z., Splichal J., Verner M., Motycka J., Vondrasek L. (2012): Single-step prediction of genomic breeding value in a small dairy cattle population with strong import of foreign genes. *Czech Journal of Animal Science*, 57, 151–159.
- Pribyl J., Madsen P., Bauer J., Pribylova J., Simeckova M., Vostry L., Zavadilova L. (2013): Contribution of domestic production records, Interbull estimated breeding values, and single nucleotide polymorphism genetic markers to the single-step genomic evaluation of milk production. *Journal of Dairy Science*, 96, 1865–1873.
- Riquet J., Coppieters W., Cambisano N., Arranz J.J., Berzi P., Davis S.K., Grisart B., Farnir F., Karim L., Mni M., Simon P., Taylor J.F., Vanmanshoven P., Wagenaar D., Womack J.E., Georges M. (1999): Fine-mapping of quantitative trait loci by identity by descent in outbred populations: application to milk production in dairy cattle. *Proceedings of the National Academy of Sciences of the United States of America*, 96, 9252–9257.
- Sanders K., Bennewitz J., Reinsch N., Thaller G., Prinzenberg E.M., Kuhn C., Kalm E. (2006): Characterization of the *DGATI* mutations and the *CSN1S1* promoter in the German Angeln dairy cattle population. *Journal of Dairy Science*, 89, 3164–3174.
- Schopen G.C.B., Koks P.D., van Arendonk J.A.M., Bovenhuis H., Visker M.H.P.W. (2009): Whole genome scan to detect quantitative trait loci for bovine milk protein composition. *Animal Genetics*, 40, 524–537.
- Sigl T., Meyer H.H.D., Wiedemann S. (2012): Gene expression of six major milk proteins in primary bovine mammary epithelial cells isolated from milk during the first twenty weeks of lactation. *Czech Journal of Animal Science*, 57, 469–480.
- Spelman R.J., Ford C.A., McElhinney P., Gregory G.C., Snell R.G. (2002): Characterization of the *DGATI* gene in New Zealand dairy population. *Journal of Dairy Science*, 85, 3514–3517.
- Suchocki T., Komisarek J., Szyda J. (2010): Testing candidate gene effects on milk production traits in dairy cattle under various parameterizations and modes of inheritance. *Journal of Dairy Science*, 93, 2703–2717.
- Szyda J., Zukowski K., Kaminski S., Zarnecki A. (2013): Testing different single nucleotide polymorphism selection strategies for prediction of genomic breeding values in dairy cattle based on low density panels. *Czech Journal of Animal Science*, 58, 136–145.
- Thaller G., Kramer W., Winter A., Kaupe B., Erhardt G., Fries R. (2003): Effects of *DGATI* variants on milk production traits in German cattle breeds. *Journal of Animal Science*, 81, 1911–1918.
- Viitala S., Szyda J., Blott S., Schulman N., Lidauer M., Maki-Tanila A., Georges M., Vilkki J. (2006): The role of the bovine growth hormone receptor and prolactin receptor genes in milk, fat and protein production in Finnish Ayrshire dairy cattle. *Genetics*, 173, 2151–2164.
- Weller J.I., Golik M., Seroussi E., Ezra E., Ron M. (2003): Population-wide analysis of a QTL affecting milk-fat



- production in the Israeli Holstein population. *Journal of Dairy Science*, 86, 2219–2227.
- Winter A., Kramer W., Werner F.A.O., Kollers S., Kata S., Durstewitz G., Buitkamp J., Womack J.E., Thaller G., Fries R. (2002): Association of lysine 232 alanine polymorphism in bovine gene encoding acyl-CoA:diacylglycerol acyltransferase (*DGATI*) with variation at a quantitative trait locus for milk fat content. *Proceedings of the National Academy of Sciences of the United States of America*, 99, 9300–9305.
- Winter A., Alzinger A., Fries R. (2004): Assessment of the gene content of the chromosomal regions flanking bovine *DGATI*. *Genomics*, 83, 172–180.
- Zavadilova L., Stipkova M. (2013): Effect of age at first calving on longevity and fertility traits for Holstein cattle. *Czech Journal of Animal Science*, 58, 47–57.
- Zavadilova L., Zink V. (2013): Genetic relationship of functional longevity with female fertility and milk production traits in Czech Holsteins. *Czech Journal of Animal Science*, 58, 554–565.

Received: 2013–05–14

Accepted after corrections: 2014–03–04

---

*Corresponding Author*

prof. Ing. Jindřich Čítek, CSc., University of South Bohemia in České Budějovice, Faculty of Agriculture, Department of Genetics and Animal Breeding, Studentská 13, 370 05 České Budějovice, Czech Republic  
Phone: +420 387 772 591, e-mail: citek@zf.jcu.cz

---