

Effect of dietary supplementation of some antioxidants on growth performance, carcass composition and breast meat characteristics in quails reared under heat stress

H. IMIK¹, M. AYDEMIR ATASEVER², M. KOC³, M. ATASEVER², K. OZTURAN⁴

¹Department of Animal Science, Faculty of Veterinary Sciences, Ataturk University, Erzurum, Turkey

²Department of Food Hygiene and Technology, Faculty of Veterinary Medicine, Ataturk University, Erzurum, Turkey

³Department of Biology, Faculty of Art and Sciences, Bozok University, Yozgat, Turkey

⁴9th Army cops “A” Type Food Control Detachment Commandery, Erzurum, Turkey

ABSTRACT: This research investigates the effects of adding vitamin E, vitamin C, vitamin E+C, and alpha lipoic acid to feed rations for Japanese quails (*Coturnix coturnix japonica*) exposed to heat stress. The aspects studied were growth performance, carcass composition and breast meat characteristics. Five groups of quails, containing 50 birds each (250 Japanese quails: 150 female and 100 male) were used. The 21-days-old birds were fed for a period of 21 days, and they were kept in a controlled environment with a temperature of 34°C between 08:00 and 17:00 and a temperature of 24°C for the remaining part of the day. The five groups under study included: a control group without any additive (BS), a group fed diets with vitamin E (BSE), with vitamin C (BSC), with vitamin E+C (BSEC) and with lipoic acid (BSLA). The supplement additions to the diets did not affect the growth performance and carcass composition of the birds. The TBA (malonaldehyde) value of the BS group was significantly higher ($P < 0.001$) than in the other groups. L^* , a^* , and b^* values in muscle samples, superficialis pectoralis muscle (SPM) and deep pectoralis muscle (DPM), were determined. In the samples from the SPM, the L^* value of the BS group was higher than in the BSC group; the a^* value of the BSE group was higher than in the BS group; and the b^* value of the BSC group was higher than in the BS group ($P < 0.05$). In the samples from the DPM, the L^* value of the BS group was higher than in the BSEC group ($P < 0.05$); the a^* values of the BSE and BSEC groups were higher than in the BS group ($P < 0.05$); and the b^* values of the BSC, BSEC and BSLA groups were higher than in the BS group ($P < 0.01$). In the microbiological analysis of meat, total aerobic mesophilic bacterial counts of the BS and BSE groups were higher than the counts in the BSC, BSEC, and BSLA groups ($P < 0.01$); coliform bacterial counts were higher in the BSE group than in the BSC group ($P < 0.05$); and lactic acid was higher in the BSE and BSEC groups than in the BSC and BSLA groups ($P < 0.01$). In conclusion, the supplemented antioxidants did not exhibit any significant effect on growth performance, but they significantly decreased lipid oxidation in the meat.

Keywords: quail; vitamin E; vitamin C; lipoic acid; growth performance; meat quality

It is known that a balance in the physiological activities of living beings is required to generate a suitable environment for the removal of free radicals by antioxidants. This balance is disturbed under certain

conditions, such as increasing or decreasing ambient temperature, environmental pollution, old age, and illness. Some substances, having strong antioxidant characteristics, such as vitamin E, vitamin C

and alpha lipoic acid, have been well-known to play important roles in protecting this balance (Azzi and Stocker, 2000; Padayatty et al., 2003). Vitamin C and lipoic acid are synthesized while Vitamin E is not synthesized in organisms of animals (NRC, 1994; Patel and Vettakkorumakankav, 1995). Vitamin C exhibits an antioxidant effect by removing free radicals from the environment (Packer et al., 1979; Tanaka et al., 1997) while the effective mechanism of vitamin E in its reactions with active radicals is to break chains, to exert pressure, to renovate, and to increase endogen defence (Packer et al., 1979; Azzi and Stocker, 2000). Alpha-lipoic acid, also known as thioctic acid, is a strong antioxidant substance having the scavenger capacity of reactive oxygen species and preventive effect on lipid oxidation (Dinçer et al., 2002; Gonzalez-Perez and Gonzalez-Castaneda, 2006).

Sahin et al. (2003) proposed that only additions of vitamin C and folic acid as antioxidants to diets of Japanese quails under heat stress do not affect body weights after feeding and that their combinations positively affect the fattening performances of animals. It has been observed in some studies (David Peebles and Brake, 1985; Puthpong siriporn et al., 2001; Sahin et al., 2002) that vitamin E and vitamin C added to diets of laying poultry under heat stress improved performance related with egg production. Imik et al. (2000) stated that vitamin E and vitamin C did not affect the performances of early-weaned, 21-days-old Ankara goat kids.

Studies show that antioxidants added to the diets of poultry (Lauridsen et al., 1997; Sahin et al., 2002; Young et al., 2003; Kennedy et al., 2005), ruminants (Liu et al., 1995; Schaefer et al., 1995) and pigs (Asghar et al., 1991; Buckley et al., 1995; Lahučký et al., 2005) prevent lipid oxidation.

The reasons for recruitment of quails as experimental model in this research are their easy management and higher resistance to stress than in other poultry. Thus, the effects of antioxidants, added to quail rations, on growth performance, carcass composition and breast meat characteristics of Japanese quail (*Coturnix coturnix japonica*) subjected to heat stress are investigated.

MATERIAL AND METHODS

Birds and feeding

This study was performed on the Research and Practice Farm at the Faculty of Veterinary Medicine,

Ataturk University. A total of 250 (150 female and 100 male) 21-days-old Japanese quails (*Coturnix coturnix japonica*) were divided into five groups. Each group had 50 quails in 5 cages. Each cage contained a random selection of 6 female and 4 male birds. Lots were drawn to determine the placement of groups in the cages. The quail cages were placed in temperature-controlled rooms during the study, receiving 17 h of daylight. A room temperature of 34°C was maintained between 8:00 and 16:00. For the rest of the time the birds were kept at a temperature of 24°C. The study spanned a total of 21 days.

The feed rations for the studied groups of quails were prepared as follows: a basal ration containing no additional antioxidant (BS); a basal diet + vitamin E (250 mg DL- α -tocopheryl-acetate/kg of diet) (BSE); a basal ration + vitamin C (250 mg L-ascorbic acid/kg of diet) (BSC); a basal ration + vitamin E + C (250 mg DL- α -tocopheryl-acetate/kg of diet + 250 mg L-ascorbic acid/kg of diet) (BSEC); and a basal diet + lipoic acid (250 mg α -lipoic acid/kg of diet) (BSLA).

The formulation and chemical composition of floury formed experimental diet, which was mixed according to NRC, are presented in Table 1. The crude protein proportion was determined by analysis (AOAC, 1990) and metabolizable energy, Ca, P were calculated (Jurgens, 1996). Small amounts of the basal diet were first mixed with the respective amounts of vitamin E and C as small batch, then with a larger amount of the basal diet until the total amounts of the respective diets were homogeneously mixed. All formulations of the diet were prepared freshly and stored at 15–21°C before feeding to quails.

Growth performance

The birds were weighed at the beginning of experiment and on the 7th, 14th and 21st day of the experiment. They were fed at the same hour each day during research. The daily feed prepared for the consumption by birds as well as any remaining feed were weighed. Water was supplied in automatic bowls. Feed and water were administered *ad libitum* during research.

Carcass composition

A total of 100 birds (5 groups \times 20 = 100) from each group (including 2 males and 2 females chosen

from each subgroup at random) were slaughtered to determine the carcass characteristics. After the quails were fasted for ten hours, the blood of the birds was allowed to flow for 120 seconds. The birds were plucked manually. The internal organs of the birds were extracted and weighed. The carcasses were kept at +4°C for 24 h, and then the weights of the cold carcasses were determined.

Meat characteristics

A total of 50 (25 male and 25 female) birds were randomly chosen for the analysis of fresh breast meat quality, including one male and one female from each of 5 subgroups. Then moisture, pH, total ash, total protein, TBA, colour analysis (L^* (relative

lightness), a^* (relative redness), b^* (relative yellowness)) and numbers of aerobic mesophilic microorganisms, coliforms and lactic acid bacteria were determined.

Analyses for moisture, total ash and total protein contents of samples were performed according to the methods declared by Gokalp et al. (2001). The pH value was determined by directly inserting a pH metre probe into the chest muscle. The protein content of the samples was determined with a Kjeldahl device.

TBA (thiobarbituric acid) was performed according to the methods suggested by Turk Standards Institute 2409 (2007). In this regard, 10 g of homogenized sample was broken into pieces and mixed with 50 ml of distilled water. This mixture was washed with 47.5 ml of distilled water and poured

Table 1. The formulation and chemical composition of the basal ration

Ingredients (%)	Starter phase (0–3 weeks)	Finisher phase (4–6 weeks)
Wheat	27.90	30.40
Maize	28.20	31.70
Maize gluten, 43% CP	15.00	15.00
Soybean meal, 44% CP	21.20	15.00
Vegetable oil	0.50	0.50
Wheat bran*	3.80	3.80
Dicalcium phosphate	1.80	2.00
D,L-Methionine	0.10	0.10
Limestone	0.70	0.70
L-Lysine hydrochloride	0.10	0.10
Salt	0.20	0.20
Vitamin-mineral premix**	0.50	0.50
Nutritional levels		
Metabolizable energy (MJ/kg)	11.91	12.15
Crude protein (%)	22.20	20.00
Ca (%)	0.81	0.84
P (%)	0.43	0.45

*the additives (vitamin E, vitamin C, vitamin E+C and lipoic acid) were added in place of wheat bran

**the vitamin-mineral premix provides the following (per kg): all-trans-retinyl acetate 1.8 mg; all-rac- α -tocopherol acetate 1.25 mg; menadione sodium bisulphate 1.1 mg; riboflavin 4.4 mg; thiamine (thiamine mononitrate) 1.1 mg; vitamin B6 2.2 mg; niacin 35 mg; Ca-pantothenate 10 mg; vitamin B12 0.02 mg; folic acid 0.55 mg; d-biotin 0.1 mg; choline chloride 175 mg; manganese (from manganese oxide) 40 mg; iron (from iron sulphate) 12.5 mg; zinc (from zinc oxide) 25 mg; copper (from copper sulphate) 3.5 mg; iodine (from potassium iodide) 0.3 mg; selenium (from sodium selenite) 0.15 mg

into a ball. 2.5 ml of hydrochloric acid was added to make a pH of 1.5. This ball was then placed on a distillation unit and 50 ml of distillate was produced. 3 distillates of 5 ml were prepared. 5 ml of 0.02M-thiobarbituric acid standard solution, prepared with 90% glacial acetic acid, was added into each tube and then the tubes were placed in a double boiler for 35 seconds. 5 ml of distilled water and 5 ml of TBA standard solution were prepared as a blank and underwent the same procedures. The tubes were then cooled. The optic density of the solution in each of the three tubes was measured with a spectrometer adjusted to a wavelength of 538 nanometres against the blank. Averages were calculated as follows:

TBA amount in the sample is determined as $\mu\text{g MA/g}$.

$$\text{TBA} = a \times 7.8$$

where:

a = absorbance of the sample in a spectrometer

Colour measurements (L^* , a^* , b^*) on samples from the superficialis pectoralis muscle (SPM) and deep pectoralis muscle (DPM) (Fitzgerald, 1969) were done using a tristimulus colorimeter (Minolta Chroma Meter Measuring Head CR-200, Minolta, Osaka, Japan).

Microbiological analysis of the samples was performed with respect to the methods suggested by Baumgart et al. (1993). 25 g of the meat sample was homogenized in 225 ml of sterilized Ringer's solution.

Then sterilized solutions were prepared. The pour plate method was used for the microbiological analysis. Plate Count Agar (PCA, Merck) was used for the enumeration of total aerobic mesophilic microorganisms. Plates were evaluated after being incubated at $30 \pm 1^\circ\text{C}$ for 72 ± 1 hours. Rogosa Acetate Agar (RAA, Merck) was used for the enumeration of lactic acid bacteria. These plates were incubated anaerobically for 5 days at $30 \pm 1^\circ\text{C}$. All colonies produced following the incubation were evaluated. Violet Red Bile Agar (VRBA, Merck) was used for the enumeration of coliform bacteria. The plates were incubated anaerobically at $30 \pm 1^\circ\text{C}$ for 24 hours. Red colonies larger than 1 mm were counted after incubation. Bacteria counts were expressed as logarithmic colony-forming units per gram of sample (log CFU/g).

Statistical analyses

The effect of antioxidants on the fattening performance and meat quality was statistically analyzed by one-way ANOVA (total). The effect of sex on meat quality and carcass characteristics was analyzed by Student's t -test for independent-samples. Finally, male and female birds were divided into two different groups and the effect of the antioxidants on meat quality was determined by one-way ANOVA. Differences between vitamin groups were determined by Duncan's multiple range test. All statistical analyses were done with SPSS. 10.00 package software programme (SPSS, 1996).




Figure 1. Body weight gains of groups during research

Table 2. Effect of dietary antioxidant addition on growth performance of quails exposed to heat stress for 21 days

Treatment*	BW initial (g)	BW final (g)	BW change (g)	BW daily (g)	Daily feed intake (g)	FCR
BS	69.20	156.30	87.10	4.15	12.43	3.02
BSE	69.00	159.80	90.80	4.32	12.49	2.90
BSC	69.90	159.44	89.54	4.27	12.21	2.86
BSEC	69.80	158.80	89.00	4.24	12.36	2.92
BSLA	69.90	162.40	92.50	4.40	13.00	2.96
SEM	0.50	2.37	1.22	0.06	0.16	0.30

BS = basal diet; BSE = diet BS plus 250 mg DL- α -tocopheryl-acetate/kg; BSC = diet BS plus 250 mg of L-ascorbic acid/kg; BSEC = diet BS plus 250 mg DL- α -tocopheryl-acetate/kg + 250 mg of L-ascorbic acid/kg; BSLA = diet BS plus 250 mg α -lipoic acid/kg; BW = body weight; FCR = BW gain/feed intake; SEM = standard error of the mean

*nonsignificant; SEM = standard error of the mean

RESULTS AND DISCUSSION

Growth performance

Weekly body weight gains of the birds are presented in Figure 1. While body weight gains of quails in the control group were highest in the first week, during the second and third week their gains declined. Body weights in the groups containing antioxidants regularly increased in the course of

research ($P < 0.05$), (Figure 1). High temperatures are required for better growth performance in early periods for poultry. Therefore, the temperature of 34°C did not affect the body weights of the BS group negatively in the first week of the study; but it had a negative effect in the second and third week of research. Antioxidants added to the feed rations of the other groups removed the negative effect of heat stress in the second and third week and positively affected the growth performance ($P < 0.05$),

Table 3. Effect of dietary antioxidant addition on carcass characteristics of quails exposed to stress for 21 days

Treatment*	Cold carcass weight (g)	Carcass yield (%)	Total weight of internal organs (g)
BS	98.70	64.68	20.80
BSE	104.10	63.76	20.40
BSC	99.70	63.87	19.30
BSEC	102.00	63.50	21.00
BSLA	102.90	62.24	21.10
SEM	3.05	0.96	1.24
Sex			
Male	99.40	64.02	20.00
Female	103.56	63.20	21.04
SEM	1.98	0.63	0.78

total weight of internal organs = liver, heart, gizzard, offal

SEM = standard error of the mean

*nonsignificant

Table 4. Effect of dietary antioxidant addition on breast meat characteristics of quails exposed to heat stress for 21 days

Treatment	Moisture (%)	pH	Total ash (%)	Total protein (%)	TBA	SPM		DPM			
						L*	a*	L*	a*		
BS	71.36	6.14	1.79	20.82	2.07 ^a	40.76 ^a	12.11 ^b	4.54 ^b	39.79 ^a	14.35 ^b	3.43 ^c
BSE	72.89	6.12	1.89	21.56	0.71 ^b	39.18 ^{ab}	14.89 ^a	5.50 ^{ab}	37.91 ^{ab}	18.60 ^a	4.87 ^{bc}
BSC	70.66	6.14	1.78	21.06	0.74 ^b	37.1 ^b	14.34 ^{ab}	7.34 ^a	37.08 ^{ab}	15.75 ^{ab}	9.00 ^a
BSEC	71.82	6.08	1.77	21.11	0.47 ^b	37.00 ^b	14.26 ^{ab}	6.70 ^{ab}	36.20 ^b	17.91 ^a	6.96 ^{ab}
BSLA	71.42	6.12	1.98	21.06	0.51 ^b	38.49 ^{ab}	13.79 ^{ab}	6.12 ^{ab}	37.22 ^{ab}	17.36 ^a	6.64 ^{ab}
SEM	1.49	0.03	0.17	0.60	0.26 ^{**}	0.86 [*]	0.79 [*]	0.71 [*]	0.90 [*]	0.84 [*]	0.97 ^{**}
Sex											
Male	71.55	6.13	1.73	21.20	1.08	38.09	13.64	5.33	37.37	16.53	6.45
Female	71.71	6.10	1.95	21.05	0.71	38.89	14.12	6.75	37.91	17.05	5.91
SEM	0.95	0.20	0.54	0.40	0.02	0.61	0.53	0.46 [*]	0.61	0.65	0.72

TBA = μg malonaldehyde/g; SPM = superficialis pectoralis muscle; SEM = standard error of the mean; ^{a-c} values within the same column having the same letters in the same sections are significantly different at * $P < 0.05$; ** $P < 0.01$

Table 5. Effect of dietary antioxidant addition on breast meat characteristics of male and female quails exposed to heat stress for 21 days

	Moisture (%)	pH	Total ash (%)	Total protein (%)	TBA	SPM		DPM		
						L*	a*	L*	a*	
Male										
BS	74.08 ^a	6.09	1.53	20.93	2.90 ^a	40.07 ^a	12.20	40.43 ^a	13.40 ^b	3.39
BSE	71.19 ^{ab}	6.13	2.01	22.03	0.91 ^b	38.94 ^{ab}	14.55	37.83 ^{ab}	19.38 ^a	5.38
BSC	69.24 ^b	6.18	1.76	21.46	0.26 ^b	35.26 ^b	14.39	34.91 ^b	15.86 ^{ab}	7.55
BSEC	72.26 ^{ab}	6.10	1.77	21.19	0.66 ^b	37.91 ^{ab}	13.37	36.67 ^{ab}	17.63 ^{ab}	7.97
BSLA	71.00 ^{ab}	6.16	1.60	20.38	0.70 ^b	38.28 ^{ab}	13.68	37.00 ^{ab}	16.38 ^{ab}	7.98
SEM	1.20*	0.05	0.16	0.65	0.30***	1.23*	1.10	1.29*	1.32*	1.33
Female										
BS	68.65	6.19 ^a	2.06	20.72	1.23	41.45 ^a	12.02	39.15	15.30	3.46 ^b
BSE	74.58	6.10 ^{ab}	1.77	21.09	0.52	39.42 ^{ab}	15.23	37.00	17.81	4.36 ^b
BSC	72.09	6.09 ^{ab}	1.80	20.66	1.22	38.94 ^{ab}	14.28	39.25	15.64	10.54 ^a
BSEC	71.38	6.06 ^b	1.77	21.04	0.28	36.07 ^b	15.15	35.72	18.18	5.94 ^b
BSLA	71.84	6.08 ^{ab}	2.36	21.73	0.32	38.56 ^{ab}	13.90	37.43	18.33	5.31 ^b
SEM	2.54	0.03*	0.27	1.03	0.28	1.12*	1.19	1.14	1.08	1.33*

TBA = μg malonaldehyde/g; SPM = superficialis pectoralis muscle; DPM = deep pectoralis muscle; SEM = standard error of the mean; ^{a-c} values within the same column having the same letters in the same sections are significantly different at * $P < 0.05$; ** $P < 0.01$

(Figure 1). These results are coherent with Sahin et al. (2003) findings on the same issue. Finding no differences between fattening performances of the groups may be a result of the short interval of the research period.

Carcass composition

At the end of the experiment, the performance (Table 2) and carcass characteristics (Table 3) of the groups were found to be similar ($P > 0.05$). In this research, the body weight of the cut male quails was 155.20 g and the body weight of the cut female quails was 164.00 g (SEM:2.77) ($P < 0.05$). Carcass weights of male quails in groups BS, BSE, BSC, BSEC and BSLA were 95.8, 96.4, 100.6, 103.4 and 100.8 g, respectively (SEM:3.89); carcass weights of female quails were 101.6, 111.8, 98.8, 100.6 and 105.0 g, respectively (SEM:4.18). It was observed that there was no statistically significant difference in carcass yield between the males (62.59%–65.51% /SEM:1.31/) and the females (61.48%–64.92% SEM:1.24/) in the groups. There was no effect on carcass composition produced by the addition of vitamin E, C, and lipoic acid to the feed rations of quails under heat stress ($P > 0.05$).

Breast meat characteristics

The effect of the antioxidants added to quail diets on meat quality is presented in Table 4. Average moisture, pH, total ash, and total protein values of the groups were determined in this study and no differences between groups and sexes were found.

TBA values of the BS group were found to be higher ($P < 0.01$) than in other groups (Table 4). Therefore it is noted that antioxidants did not affect the TBA values in the meat of females, while antioxidants decreased the TBA values in the meat of males ($P < 0.001$), (Table 5). The TBA values of the quail groups that consumed diets containing vitamin E, vitamin C, vitamin E+C, and lipoic acid were found to be lower than in the control group in the present study ($P < 0.01$), (Table 4). Most studies show that various antioxidants added to feed rations of poultry prevented lipid oxidation (Lauridsen et al., 1997; Nam and Ahn, 2003; Young et al., 2003; Kennedy et al., 2005; Ryu et al., 2005). Some studies reported that

lipoic acid decreased lipid oxidation (Dinçer et al., 2002; Shen et al., 2005).

The L^* value of the BS group was higher than in the BSC and BSEC groups; the relative redness a^* value of the BSE group was higher than in the BS group; the relative yellowness b^* value of the BSC group was higher than in the BS group ($P < 0.05$) in colour analyses from the SPM. The relative lightness L^* value of the BS group was higher than in the BSEC group; the relative redness a^* values of the BSE, BSEC and BSLA groups were higher than in the BS group ($P < 0.05$). The relative yellowness b^* values of the BSE, BSEC and BSLA groups were determined to be significantly higher than in the BS group ($P < 0.01$) in colour analyses from the DPM (Table 4).

Previous studies explained the effect of antioxidants on L , a^* and b^* values of meat in poultry (Liu et al., 1995; Nam and Ahn, 2003; Young et al., 2003) and pigs (Asghar et al., 1991; Buckley et al., 1995). Literary sources also indicate that the meat of poultry fed antioxidants lasted longer (Liu et al., 1995; Nam and Ahn, 2003). Findings in both the SPM and DPM make it clear that antioxidants affect colour parameters of the meat to a different extent (Table 4). These findings are coherent with the literature. Antioxidants added to diets of the experimental groups produced limited effects on the meat colour of female quails (SPM = L^* ; DPM = b^*) while antioxidants added to diets of the experimental groups affected the meat colour of male quails (SPM = L^* , b^* ; DPM = L^* , a^*) ($P < 0.05$) (Table 5). L^* , a^* , and b^* values of these studies by Du et al. (2000) are 53.6–58.5, 12.9–20.1 and 17.6–23.0, respectively; in Du et al. (2001) they are 80.46–85.42, 7.40–11.56 and 19.21–22.20; in Young et al. (2003) they are 50.3–52.4, 2.5–2.9 and 2.9–4.1; in the pectoralis major muscle of female broilers they are 50.5–51.0, 3.9–4.2 and 2.7–4.7 in iliobtibialis muscle; in Aksu et al. (2007) they are 51.61, 7.16 and 3.63 for male quails, and 49.20, 7.01 and 4.33 for female quails; 50.34, 5.35 and 3.87 in fresh breast; and 50.47, 8.82 and 4.10 in fresh drumstick.

The L^* value of the BSC group was found to be lower than in the BS group in colour analyses from the SPM of male quails ($P < 0.05$). The b^* values of the BSC and BSEC groups were determined to be higher than in the BS group ($P < 0.05$). The L^* value of the BSC group was lower than in the BS group; the a^* value of the BS group was found to be lower than in the BSE group ($P < 0.05$) in the DPM. The L^* value of the BS group was found to be higher than

Table 6. Effect of dietary antioxidant addition on carcass total aerobic mesophilic, coliform and lactic acid bacteria in meat samples of quails exposed to heat stress for 21 days

	Total aerobic mesophilic bacteria (log CFU/g)	Coliform bacteria (log CFU/g)	Lactic acid bacteria (log CFU/g)
Treatment			
BS	5.17 ^a	2.59 ^{ab}	2.81 ^{ab}
BSE	4.89 ^a	3.70 ^a	3.85 ^a
BSC	3.63 ^b	1.80 ^b	1.83 ^b
BSEC	3.90 ^b	1.97 ^{ab}	3.34 ^a
BSLA	3.85 ^b	1.88 ^{ab}	1.47 ^b
SEM	0.29 ^{**}	0.59 [*]	0.46 ^{**}
Sex			
Male	4.50	2.44	3.03
Female	4.08	2.34	2.29
SEM	0.23	0.39	0.34

SEM = standard error of the mean; ^{a,b}values within the same column having the same letters in the same sections are significantly different at * $P < 0.05$; ** $P < 0.01$

Table 7. Effect of dietary antioxidant addition on carcass total aerobic mesophilic, coliform and lactic acid bacteria in meat samples of male and female quails exposed to heat stress for 21 days

	Total aerobic mesophilic bacteria (log CFU/g)	Coliform bacteria (log CFU/g)	Lactic acid bacteria (log CFU/g)
Male			
BS	5.3 ^a	2.85	2.88 ^{ab}
BSE	5.45 ^a	2.66	4.31 ^a
BSC	3.41 ^b	1.95	1.51 ^b
BSEC	3.58 ^b	2.87	4.64 ^a
BSLA	4.71 ^a	1.88	1.84 ^b
SEM	0.23 ^{***}	0.92	0.46 ^{**}
Female			
BS	5.00 ^a	2.33 ^b	2.75
BSE	4.33 ^{ab}	4.77 ^a	3.39
BSC	3.85 ^{ab}	1.66 ^b	2.16
BSEC	4.21 ^{ab}	1.08 ^b	2.03
BSLA	2.99 ^b	1.88 ^b	1.11
SEM	0.43 [*]	0.69 [*]	0.64

SEM = standard error of the mean; ^{a,b}values within the same column having the same letters in the same sections are sig-

in the BSEC group in colour analyses from the SPM of female quails ($P < 0.05$). The b^* value of the BSC group was determined to be higher than in the other groups in the DPM ($P < 0.05$) (Table 5).

The effects of the addition of antioxidants to quail diets on the counts of microorganisms are presented in Table 6. Total aerobic mesophilic bacteria counts in the BSC, BSEC, and BSLA groups were found to be lower than in the BS and BSE groups ($P < 0.01$). Coliform counts in the BSC group were determined to be lower than those in the BSE group ($P < 0.05$). Lactic acid bacteria counts were found to be lower in quail muscles containing vitamin C and lipoic acid ($P < 0.01$) than in meat from quails whose diets contained vitamin E and E+C. Microbiological results of antioxidant addition with respect to sex are shown in Table 7.

Vitamin C added to the feed ration decreased the total aerobic mesophilic, coliform and lactic acid bacteria counts; and lipoic acid decreased the total mesophilic, aerobic and lactic acid bacteria counts. Vitamin E produced no effect (Table 6). Aksu et al. (2005) reported that total aerobic mesophilic bacterial counts were 3.99 and 3.83 log CFU/g in the breast and drumstick of broilers, respectively, and that lactic acid bacterial counts were 3.04 and 2.80 log CFU/g. Manousaridis et al. (2005) found out that the total aerobic mesophilic bacterial count was 3.4 log CFU per g and that the lactic acid bacterial count was 1.2 log CFU/g. Aksu et al. (2007) stated that the broiler lactic acid bacterial count was 3.62 CFU/g in male quails and that the broiler lactic acid bacterial count was log 3.47 CFU/g in female quails. Karpińska-Tymoszyk (2007) also pointed out that the total aerobic mesophilic bacterial count was 2.6×10^5 CFU/g and the coliform bacterial count was 1.3×10^4 CFU/g in turkey meat. Saucier et al. (2000) found that lactic acid bacterial counts were 3.67 log CFU/g and 4.82 log CFU/g when they were kept in stock under different conditions; and that total aerobic mesophilic and coliform bacteria increased in chicken meat more than in turkey meat.

CONCLUSION

The antioxidants (vitamin E, vitamin C, lipoic acid) and antioxidant combinations (vitamin E+C) added to the feed rations did not have any significant effect on growth performance, but they decreased lipid oxidation in meat and modified the

colour characteristics of meat in Japanese quails under heat stress. The effects on growth performance with increased amounts of antioxidants and a longer fattening period should be a focus of further studies.

REFERENCES

- Aksu M.I., Karaoglu M., Kaya M., Esenbuga N., Macit M. (2005): Effect of dietary humate on the pH, TBARS and microbiological properties of vacuum- and aerobic-packed breast and drumstick meats of broilers. *Journal of the Science of Food and Agriculture*, 85, 1485–1491.
- Aksu M.I., Imik H., Karaoglu M. (2007): Influence of dietary sorghum (*Sorghum vulgare*) and corn supplemented with methionine on cut-up pieces weights of broiler carcass and quality properties of breast and drumsticks meat. *Food Science and Technology International*, 13, 361–367.
- AOAC (1990): Official Methods of Analysis. Vol. I. 15th ed. Association of Official Analytical Chemists, Arlington, USA.
- Asghar A., Gray J.I., Miller E.R., Ku P.K., Booren A.M., Buckley D.J. (1991): Influence of supranutritional vitamin E supplementation in the feed on swine growth performance and deposition in different tissues. *Journal of the Science of Food and Agriculture*, 57, 19–29.
- Azzi A., Stocker A. (2000): Vitamin E: non-antioxidant roles. *Progress in Lipid Research*, 39, 231–255.
- Baumgart J., Firnhaber J., Spcher G. (1993): Mikrobiologische untersuchung von lebensmitteln behr's verlag. Hamburg, Germany.
- Buckley D.J., Morrissey P.A., Gray J.I. (1995): Influence of dietary vitamin E on the oxidative stability and quality of pig meat. *Journal of Animal Science*, 73, 3122–3130.
- David Peebles E., Brake J. (1985): Relationship of dietary ascorbic acid to broiler breeder performance. *Poultry Science*, 64, 2041–2048.
- Dinçer Y., Telci A., Kayali R., Yilmaz I.A., Cakatay U., Akcay T. (2002): Effect of α -lipoic acid on lipid peroxidation and anti-oxidant enzyme activities in diabetic rats. *Clinical and Experimental Pharmacology and Physiology*, 29, 281.
- Du M., Ahn D.U., Nam K.C., Sell J.L. (2000): Influence of dietary conjugated linoleic acid on volatile profiles, color and lipid oxidation of irradiated raw chicken meat. *Meat Science*, 56, 87–395.

- Du M., Hur S.J., Nam K.C., Ismail H., Ahn D.U. (2001): Volatiles, color, and lipid oxidation of broiler breast fillets irradiated before and after cooking. *Poultry Science*, 80, 1748–1753.
- Fitzgerald T.C. (1969): The coturnix quail anatomy and histology. Iowa State University Press, IOWA, USA, 132–133.
- Gokalp H.Y., Kaya M., Tulek Y., Zorba O. (2001): Guide for quality control and laboratory application of meat products, 4th ed. Atatürk University Publication, Erzurum, Turkey, No. 751.
- Gonzalez-Perez O., Gonzalez-Castaneda R.E. (2006): Therapeutic perspectives on the combination of α -lipoic acid and vitamin E. *Nutrition Research*, 26, 1–5.
- Imik H., Aytac M., Coskun B., Fidanci H. (2000): Effects of E and C vitamins on the growth and immunity of Angora goat kids exposed to stress. *Turkish Journal of Veterinary and Animal Sciences*, 24, 51–58.
- Jurgens M.H. (1996): *Animal Feeding and Nutrition*, 8th ed. Kendall/Hunt Pub. Co., Dubuque, IA.
- Karpińska-Tymoszczyk M. (2007): Effects of sage extract (*Salvia officinalis* L.) and a mixture of sage extract and sodium isoascorbate on the quality and shelf life of vacuum-packed turkey meatballs. *Journal of Muscle Foods*, 18, 420–434.
- Kennedy O.B., Stewart-Knox B.J., Mitchell P.C., Thurnham D.I. (2005): Vitamin E supplementation, cereal feed type and consumer sensory perceptions of poultry meat quality. *British Journal of Nutrition*, 93, 333–338.
- Lahučký R., Bahelka I., Novotná K., Vašíčková K. (2005): Effects of dietary vitamin E and vitamin C supplementation on the level of α -tocopherol and L-ascorbic acid in muscle and on the antioxidative status and meat quality of pigs. *Czech Journal of Animal Science*, 50, 175–184.
- Lauridsen C., Buckley D.J., Morrissey P.A. (1997): Influence of dietary fat and vitamin E supplementation on alpha-tocopherol levels and fatty acid profiles in chicken muscle membranous fractions and on susceptibility to lipid peroxidation. *Meat Science*, 46, 9–22.
- Liu Q., Lanari M.C., Schaefer D.M. (1995): A review of dietary vitamin E supplementation for improvement of beef quality. *Journal of Animal Science*, 73, 3131–3140.
- Manousaridis G., Nerantzaki A., Paleologos E.K., Tsiuotsias A., Savvaidis I.N., Kontominas M.G. (2005): Effect of ozone on microbial, chemical and sensory attributes of shucked mussels. *Food Microbiology*, 22, 1–9.
- Nam K.C., Ahn D.U. (2003): Use of double packaging and antioxidant combinations to improve color, lipid oxidation, and volatiles of irradiated raw and cooked turkey breast patties. *Poultry Science*, 82, 850–857.
- NRC (1994): National Research Council. *Nutrient Requirements of Poultry*, 9th revised edition. NRC, National Academy Press, Washington, USA.
- Packer J.E., Slater T.F., Willson R.L. (1979): Direct observation of a free radical interaction between vitamin E and vitamin C. *Nature*, 278, 737–738.
- Padayatty S.J., Kkatz A., Wang Y.H., Eck P., Kwon O., Lee J.H., Chen S.L., Corpe C., Dutta A., Dutta S.K., Levine M. (2003): Vitamin C as an antioxidant: Evaluation of its role in disease prevention. *Journal of the American College of Nutrition*, 22, 18–35.
- Patel M.S., Vettakkorumakankav N.N. (1995): Lipoic acid-requiring proteins: recent advances. In: Packer L., Cadenas E. (eds.): *Biothiols in Health and Disease*. Marcel Dekker Inc., New York, USA, 373–388.
- Puthongsiriporn U., Scheideler S.E., Sell J.L., Beck M.M. (2001): Effects of vitamin E and C supplementation on performance, *in vitro* lymphocyte proliferation, and antioxidant status of laying hens during heat stress. *Poultry Science*, 80, 1190–1200.
- Ryu Y.C., Rhee M.S., Lee K.M., Kim B.C. (2005): Effects of different levels of dietary supplemental selenium on performance, lipid oxidation, and color stability of broiler chicks. *Poultry Science*, 84, 809–815.
- Sahin K., Sahin N., Onderci M. (2002): Vitamin E supplementation can alleviate negative effects of heat stress on egg production, egg quality, digestibility of nutrients and egg yolk mineral concentrations of Japanese quails. *Research in Veterinary Science*, 73, 307–312.
- Sahin K., Onderci M., Sahin N., Gursu M.F., Kucuk O. (2003): Dietary vitamin C and folic acid supplementation ameliorates the detrimental effects of heat stress in Japanese quail. *Journal Nutrition*, 133, 1882–1886.
- Saucier L., Gendron C., Garipey C. (2000): Shelf life of ground poultry meat stored under modified atmosphere. *Poultry Science*, 79, 1851–1856.
- Schaefer D.M., Liu Q., Faustman C., Yin M.C. (1995): Supranutritional Administration of Vitamins E and C Improves Oxidative Stability of Beef. *Journal Nutrition*, 125, 1792–1798.
- Shen Q.W., Jones C.S., Kalchayanand N., Zhu M.J., Du M. (2005): Effect of dietary alpha-lipoic acid on growth, body composition, muscle pH, and AMP-activated protein kinase phosphorylation in mice. *Journal of Animal Science*, 83, 2611–2617.
- Statistical Packages for the Social Sciences (1996): SPSS for Windows release 10.01. SPSS Inc., Chicago, USA.
- Tanaka K., Hashimoto T., Tokumaru S., Iguchi H., Kojo S. (1997): Interactions between vitamin C and vitamin

E are observed in tissues of inherently scorbutic rats. Journal Nutrition, 127, 2060–2064.
Turkish Standart Instiýtute (2007): Chicken carcass meat. Chemical properties. Ankara, Turkey.
Young J.E., Stagsted J., Jensen S.K., Karlsson A.H., Henckel P. (2003): Ascorbic acid, alpha-tocopherol,

and oregano supplements reduce stress-induced deterioration of chicken meat quality. Poultry Science, 82, 1343–1351.

Received: 2009–06–05

Accepted after corrections: 2009–10–14

Corresponding Author

Asst. Prof. Dr. Halit Imik, Department of Animal Nutrition and Nutritional Disorders, Faculty of Veterinary Medicine, Atatürk University, 25200 Erzurum, Turkey
Tel. +90 442 236 0880, fax: +90 442 236 0881, e-mail: Halitimik60@hotmail.com; himik@atauni.edu.tr

The journal publishes original scientific papers and board-invited review articles in English. The author is fully responsible for the originality of the paper and its subject and formal correctness. The author's declaration that the paper has not been published anywhere else should be enclosed. The Board of Editors decides on the publication of papers, taking into account peer reviews, scientific importance, and manuscript quality. Good laboratory practice and ethical rules must be followed. The SI international system of measurement units should be used. Manuscripts must be grammatically and linguistically correct and authors whose native language is not English are advised to seek the help of native English-speaker. Manuscript containing language errors are prejudiced in the reviewing process and may be returned to the author for rewriting before peer review and/or before acceptance.

Only manuscripts assessed by leading experts in the field will be published. If such reviews are not available within four months after registration of the manuscript, the peer review process is ceased, and the authors are notified. They can resubmit the manuscript, after its thorough revision and/or update, either to Czech Journal of Animal Science for a new assessment or to another journal. This should eliminate a long waiting period, and likely rejection of the manuscript.

If a revision of manuscript following the recommendation of the reviewers is requested, the modified manuscript must be re-submitted within six weeks. The authors can, however request an extension of the re-submission deadline, if necessary. All parts of the manuscript, including tables and figures (even unchanged) must be re-submitted. A detailed reply by the authors to every point of reviewer's recommendations must be attached to the revision manuscript (cover letter). It is not necessary to accept all requests of the reviewers, but a clear explanation of why reviewers' comments were not accepted, has to be provided. If the deadline for re-submission is missed, the paper will be removed from the reviewing process.

The proof reading must be returned within three days. Only errors originating during preparation of the document for printing can be corrected. Standard proof marks will be used. No changes in the manuscript after acceptance for publication can be allowed.

Czech Journal of Animal Science **do not accept hard-copy paper manuscripts**; all manuscripts must be submitted electronically to the journal website. This method of submission results in much faster handling of your manuscript, fewer handling errors, and allows you to track the handling progress of your manuscript at any time. Manuscripts must be submitted as a Microsoft Word or other word processing document (filetype ".doc" or ".rtf").

Copyright. The journal is protected by copyright held by the publisher after the manuscript has been accepted for publication. As concerns the transfer of rights, the corresponding author takes over responsibility for all authors. No part of this publication may be reproduced, stored, or transmitted in any form or by any means, without the written permission of the publisher.

Manuscript layout. The Microsoft (MS) Word for Windows word-processing software should be used for creating the text in non-formatted style strictly following the journal layout. The manuscript must be typed (Times New Roman, font 12) with double spacing throughout. Lines must be numbered in a consecutive manner starting on the first page, in the left-hand margin. All pages of the manuscript must also be numbered consecutively.

If any abbreviations are used in the paper, they have to be explained appropriately when they are used in the text for the first time. It is not advisable to use any abbreviation in the paper title or in the abstract. Tables, graphs and other MS Word documents are to be submitted on separate pages appended to the article. The document must not be formatted in columns, heading styles, etc. This unique MS Word file must be saved under the first author's surname only. Graphs should be provided in MS Excel and they should be stored with original data. Photographs and autotypes should be submitted in high resolution (min. 300 dpi) JPG or TIFF format. All tables, graphs and photos should be numbered in the order in which they are included in the text, using Arabic numerals.

Manuscript length. Originals papers, including figures, tables and references, should not exceed 25 typed or computer-written pages. The number of figures and tables should be kept to a minimum.

The Title of the Paper should be short and informative, no subtitles or numbering the "serial" articles (Part I, Part II etc.) should be used.

The Abstract should not have more than 2,500 keystrokes (characters plus spaces). It should contain important information on methods used to solve the problem, clear description of results and their statistical significance, and brief and unambiguous conclusions drawn from the results. References and discussion of results should not be included in the abstract.

Keywords should not repeat nouns used in the title and should describe the studied problem as best as possible.

The Introduction section should provide information on the present state of research in the field concerned. It briefly justifies the research, specifies the hypotheses to be tested and gives the objective (s). References to literary sources document such

present findings that are used by the authors, not all that have been published until now. References in the text should agree with those in the list of references. It is recommended to include references to papers from peer periodicals only. Citations from non-available sources (reports, national journals, proceedings etc.) should be omitted. Papers published by one or two authors are to be cited by their names, those published by three or more authors by the name of the first one et al. If more than one paper by the same author(s) published in the same year and are cited, they should be differentiated by YEAR_{a,b,c} both in the text and the list of References. Names and year of publication are to be cited by including them in the text directly, e.g. "...as published by Brown (1995)" or indirectly – citing authors and year of publication in parenthesis (Green and Grey, 1996), (Jakl et al., 2002). Several papers cited together should be arranged according to the year of publication starting with the oldest one.

Material and Methods. All preliminary material, conducted experiments, their extent, conditions and course (experimental design) should be described in detail in this section. All original procedures that were used for the processing of experimental material and all analytical methods used for evaluation should also be detailed. Data verifying the quality of acquired data should be indicated for the used methods. The whole methodology is to be described only if it is an original one, in other cases it is sufficient to cite the author of the method and to mention any particular differences. Methods of statistical processing including the software used should also be listed in this section.

Results and Discussion. The results obtained from the experiments including their statistical evaluation and any commentary should be presented graphically or in tables in this section. The author should confront partial results with data published by other authors, whose names and year of publication are to be cited by including them in the text directly or indirectly.

References should be arranged in alphabetical order according to the surname and initials of authors. The year of publication cited in parenthesis, the full title of the paper in English with the language of publication in parenthesis, e.g. (in Czech) should follow. The title of the periodical should be typed in full.

Only papers cited in the text should be included in the list of references. All names of the authors must be printed in English transcription without national letters. Authors are responsible for the accuracy of their references.

Examples of references in the list:

- **Journal article**

Brown J. (1995): Estradiol determination in post-partum sows. *Journal of Endocrinology*, 198, 155–169.

- **Electronic journal article:**

Steinkraus K.H. (2002). Fermentation in world food processing. *Comprehensive Reviews in Food Science and Food Safety* [serial online]. 1, 23–32. Available from www.ift.org (accessed Apr 1, 2002).

- **Books and article within edited books, proceedings**

Spally M.R., Morgan S.S. (1989): *Methods of food analysis*. 2nd Ed. Elsevier, New York, 682 pp.

Kalab J. (1995): Changes in milk production during the sexual cycle. Hekel K. (ed.): *Lactation in Cattle*. Academic Press, London, UK, 876–888.

Janson L., Ahlin K.A. (1992): Postpartum reproductive performance in cattle selected for high and low fat content. In: *Proc. 43rd Annu. Mtg. European Association for Animal Production (EAAP)*. Madrid, Spain, 93–95.

- **Patent:**

Harred J.E., Knight A.R., McIntyre J.S., inventors; Dow Chemical Co., assignee. 1972 Apr 4. Epoxidation process. U.S. patent 3, 654, 317.

- **Dissertation:**

Smith D.E. (1988): Lipid oxidation at very low water activities. PhD Diss. Ithaca, NY: Cornell University. Available from: University Microfilms, Ann Arbor, MI: ABD62-83. 210 p.

Corresponding Author should include his or her full name including all academic, scientific and pedagogic titles and detailed address of the institution with postal code, phone and fax numbers, e-mail address. The author who is responsible for any correspondence with the journal should be indicated clearly.

Declaration of the authors must be carefully completed and signed by the first author, and then scanned in .pdf format to be uploaded in the submission of your manuscript, along side with the other manuscript files.

Offprints: Free reprint in Portable Document Format (PDF) sent via e-mail as an attachment.

Compliance with these instructions is obligatory for all authors. If a manuscript does not comply exactly with the above requirements, the editorial office will not accept it for a consideration and will return it to the authors without reviewing.

Revised February 2010