

## ***Lr39* + *Pm21*: a New Effective Combination of Resistance Genes for Leaf Rust and Powdery Mildew in Wheat**

ALEKSANDRA PIETRUSIŃSKA<sup>1</sup>, PAWEŁ CZĘSTOBOR CZEMBOR<sup>2</sup> and JERZY HENRYK CZEMBOR<sup>1</sup>

<sup>1</sup>Department of Plant Breeding and Genetics and <sup>2</sup>Department of Plant Pathology, Plant Breeding and Acclimatization Institute, National Research Institute at Radzików, Błonie, Poland

### **Abstract**

PIETRUSIŃSKA A., CZEMBOR P.CZ., CZEMBOR J.H. (2013): *Lr39* + *Pm21*: a new effective combination of resistance genes for leaf rust and powdery mildew in wheat. Czech J. Genet. Plant Breed., 49: 109–115.

Two effective resistance genes were introduced, one for leaf rust (*Lr39*) and the other for powdery mildew (*Pm21*), into the susceptible German wheat cv. Lexus. Molecular selection of plant material was carried out using closely linked markers to the introduced genes (foreground selection). In addition, for the BC<sub>1</sub>F<sub>1</sub> population, background selection was carried out using AFLP markers that were distributed randomly throughout the wheat genome. Moreover, resistance tests were conducted using natural pathogen populations of *Puccinia triticina* and *Blumeria graminis*. The use of molecular markers for foreground selection in combination with pathology tests enabled 66 homozygous lines to be obtained that were simultaneously resistant to leaf rust and powdery mildew.

**Keywords:** *Blumeria graminis*; gene pyramiding; *Puccinia triticina*

Despite the knowledge of resistance genes, it is still important to look for new and effective sources of resistance to fungal diseases, especially because the resistance of cultivars can be overcome by new virulent pathogen races. One of the best strategies to obtain effective and durable resistance in cultivars is gene pyramiding, namely the selection of multiple resistance genes. Since the expression of individual resistance genes is difficult to monitor in offspring populations, marker-assisted selection (MAS) became a powerful alternative. There are several published examples of resistance breeding in common wheat using gene pyramiding, which include: *Pm2* + *Pm4a*, *Pm2* + *Pm21*, *Pm4a* + *Pm21* in the Chinese cv. Yang 158 (LIU *et al.* 2000), *Lr16* + *Lr34* and *Lr22a* + *Lr52* in wheat populations (HIEBERT *et al.* 2010).

The ultimate aim of our work was to increase wheat resistance to powdery mildew and leaf rust

by pyramiding two highly effective resistance genes *Lr39* and *Pm21*. The 6VS/6AL translocation line of Yangmai 5, which carries the *Pm21* gene derived from the wild wheat relative *Dasypyrum villosum*, was used as the donor of resistance to powdery mildew. The line KS90WGRC10, which carries the *Lr39* (formerly *Lr41*) gene derived from the diploid wild wheat *Triticum tauschii*, was used as the donor of resistance to leaf rust (SINGH *et al.* 2004).

### **MATERIAL AND METHODS**

#### **Plant materials**

The wheat line KS90WGRC10, which carries the *Lr39* gene, was used as the source of resistance to leaf rust. This line was developed at the

Wheat Genetic and Genomic Resources Center at Kansas State University, USA. The translocation line 6VS/6AL Yangmai5, which carries the *Pm21* gene, was used as the donor of resistance to powdery mildew. The Yangmai5 line was developed at the Cytogenetics Institute, Nanjing Agricultural University (CINAU) (CAO *et al.* 2006). The German wheat cv. Lexus, which was provided by Dr Andreas Jacobi from W. von Borries-Eckendorf (Leopoldshöhe, Germany), was used as the recurrent parent susceptible to both diseases.

### Experimental design

To introduce the two resistance genes, *Lr39* and *Pm21*, into the cv. Lexus, the breeding scheme shown in Figure 1 was used.

### Screening for leaf rust and powdery mildew

Screening for resistance to leaf rust and powdery mildew was carried out in a greenhouse on plants at the three-leaf stage. The greenhouse conditions were 16–22°C with a light:dark cycle of 16:8 h. The inoculum sources were natural populations of *P. triticina* and *B. graminis*. The chosen inocula for both pathogens were avirulent to line KS90WGRC10 and virulent to Lexus. Plants were inoculated initially with *P. triticina* and then after 3–4 days the same plants were inoculated with the Bgt28 *B. graminis*. Reactions to infection were assessed 8–10 days later using a scale in which 0–2 indicates resistant plants and 3–4 indicates susceptible plants, with the latter displaying symptoms of leaf rust (LEVINE & CHEREWICK 1952) or powdery mildew (MAINS & DIETZ 1930).

### DNA extraction

For the detection of resistance genes in plant materials, DNA was extracted from fresh leaves of plants using the method described by HIGGINS *et al.* (2000). For background selection, total DNA was isolated using a spin column-based method (DNeasy Plant Mini Kit, Qiagen, Valencia, USA).

### Foreground selection (FS)

***Lr39*, a leaf rust resistance gene.** In foreground selection to detect the *Lr39* gene, five microsatellites (Simple Sequence Repeats, SSRs) were used: *Gdm35* (170 bp), *Barc124* (250 bp), *Gwm261* (160 to

200 bp), *Gwm296* (135 bp), and *Gwm210* (182 bp) (RÖDER *et al.* 1998; PESTSOVA *et al.* 2000; SINGH *et al.* 2004; SOMERS *et al.* 2004; SONG *et al.* 2005). Polymerase chain reactions (PCRs) were performed in low-profile PCR tube strips (BIO-RAD, Hercules, USA) in a Mastercycler ep thermocycler (Eppendorf, Hamburg, Germany). PCRs were carried out under the following conditions when the total reaction volume of 8 µl contained: 1.5 µl of DNA solution, 1× buffer (MBI Fermentas, Vilnius, Latvia), 2.5mM MgCl<sub>2</sub>, 0.2mM dNTPs, 0.5µM primer, and 1 U of *Taq* DNA polymerase (MBI Fermentas, Vilnius, Latvia). Amplification products were separated on denaturing polyacrylamide gels (Long Ranger Gel Solution, Rockland, USA) on an ABI Prism 377 automated DNA sequencer for 2 h.

***Pm21*, a powdery mildew resistance gene.** In foreground selection to detect the *Pm21* gene in plant material, three molecular markers were used: *SCAR*<sub>1250</sub> (1250 bp), *SCAR*<sub>1400</sub> (1400 bp) (LIU *et al.* 1999; YILDIRIM *et al.* 2004), and *NAU/xibao* (902 bp) (CAO *et al.* 2006). The 10-µl reaction mixture contained the following components: 4 µl of DNA solution, 1 × buffer (MBI Fermentas), 2.5mM MgCl<sub>2</sub>, 0.2mM dNTPs, 0.5µM primer, and 1 U of *Taq* DNA polymerase (MBI Fermentas). Amplification products were separated on 1.6% agarose gels in 1 × TBE buffer for 4 h, and visualised under UV light after the addition of ethidium bromide.

### AFLP analysis and estimation of the percentage of recurrent parent genome

Amplified fragment length polymorphism (AFLP) analysis was performed in accordance with the protocol described by Vos *et al.* (1995) at KeyGene (Wageningen, the Netherlands). After a prescreening of the parental lines, the 19 most informative *Pst*I+3/*Taq*I+3 primer combinations were chosen: *P38/T43*, *P41/T41*, *P39/T41*, *P35/T43*, *P36/T46*, *P38/T44*, *P41/T42*, *P36/T45*, *P41/T40*, *P42/T41*, *P39/T42*, *P35/T44*, *P39/T40*, *P44/T40*, *P37/T44*, *P37/T45*, *P35/T46*, *P36/T44* and *P40/T40*. On the basis of visual inspection of the fingerprints obtained in the prescreening, the five best *Pst*I-*Taq*I AFLP primer combinations (*P38/T43*, *P41/T41*, *P39/T41*, *P35/T43*, *P36/T46*) were selected for the background analysis of BC<sub>1</sub>F<sub>1</sub> plants to detect the AFLP fragments on a MegaBace 1000 capillary platform (GE Healthcare, Pittsburgh, USA). The percentage of the recurrent parent genome (% RP) in the BC samples was estimated

using the formula developed by KeyGene (Wageningen, the Netherlands).

### Genetic map construction.

Genetic maps for the BC<sub>1</sub>F<sub>3</sub> generation were constructed with 5 microsatellites that flanked the *Lr39* gene using JoinMap 4.0 (Kyazma B.V., Wageningen, the Netherlands) at LOD > 3.0 (LANDER *et al.* 1987) and the Kosambi mapping function (KOSAMBI 1943).

## RESULTS AND DISCUSSION

First, two crosses were carried out between Lexus and the line KS90WGRC10, and between Lexus and the translocation line 6VS/6AL Yangmai5. Next, we performed crosses between (Lexus × *Lr39*) and (Lexus × *Pm21*) and analysed 39 plants (Table 1). Three molecular markers were used to detect the *Lr39* gene in the F<sub>1</sub> plants: *Gdm35*, *Barc124*, and *Gwm210*. A 170 bp allele for the *Gdm35* marker was amplified in 24 plants, whereas in 13 plants both a 250 bp allele for *Barc124* and a 182 bp allele for *Gwm210* were amplified (Table 1). Two specific molecular markers were used to detect the *Pm21* gene in the F<sub>1</sub> plants: *SCAR*<sub>1250</sub> (1250 bp) and *SCAR*<sub>1400</sub> (1400 bp). Seventeen plants yielded the expected 1250 bp band for *SCAR*<sub>1250</sub> and 1400 bp band for *SCAR*<sub>1400</sub>. When the results were compared, 15 plants carried both resistance genes (*Lr39* + *Pm21*) (Table 1).

The plants that carried both *Lr39* and *Pm21* were used in backcrosses with the Lexus cultivar as the recurrent parent to obtain a BC<sub>1</sub>F<sub>1</sub> generation. Ninety-four BC<sub>1</sub>F<sub>1</sub> plants were tested for the presence or absence of the resistance genes by foreground selection (Table 1). To detect the *Lr39* gene in the BC<sub>1</sub>F<sub>1</sub> plants, the same 3 microsatellites described above were used. With the primers for *Gdm35*, a 170 bp fragment was amplified in 53 plants, whereas the primers for *Barc124* amplified a 250 bp fragment in 46 plants, and the primers for *Gwm210* amplified a 182 bp fragment in 12 plants (Table 1). This marker *NAU/xibao* was used to detect the presence of the *Pm21* gene in the BC<sub>1</sub>F<sub>1</sub> generation, and the expected 902 bp fragment was amplified in 44 plants (Table 1). Overall, 24 plants showed the presence of both (*Lr39* + *Pm21*).

Next, these 24 plants were subjected for BS analysis using 5 combinations of AFLP primers: *P38/*

Table 1. Molecular analysis (foreground and background selection) and resistance of the ((Lexus × *Lr39*) × (Lexus × *Pm21*)) × Lexus population to leaf rust and powdery mildew

Generation	Population size	Foreground selection						Background selection		Resistance test		
		No. of plants with <i>Lr39</i>			No. of plants with <i>Pm21</i>			Background selection range % RP	No. of resistant plants	<i>P. triticina</i>	<i>B. graminis</i>	No. of plants selected
<i>Gdm35</i> 170 bp	<i>Barc124</i> 250 bp	<i>Gwm210</i> 182 bp	<i>Gwm261</i> 165 bp	<i>Gwm296</i> 135 bp	<i>SCAR</i> <sub>1250</sub> 1250 bp	<i>SCAR</i> <sub>1400</sub> 1400 bp	<i>NAU/xibao</i> 902 bp					
F <sub>1</sub>	39	24	13	13	–	–	17	17	–	–	–	15
BC <sub>1</sub> F <sub>1</sub>	94	53	46	12	–	–	–	44	32.1–75.0	–	–	15
BC <sub>1</sub> F <sub>2</sub>	188	47	42	27	76	15	–	111	–	151	162	45
BC <sub>1</sub> F <sub>3</sub>	742	184	200	135	150	165	–	262	–	471	642	66

– not tested

*T43*, *P41/T41*, *P39/T41*, *P35/T43*, and *P36/T46*. These primers generated a total of 28 polymorphic AFLP markers. On the basis of the results, the % RP for the BC<sub>1</sub>F<sub>1</sub> was estimated. The % RP was in the range of 32.1–75% (Table 1). Fifteen plants which were in the range from 50 to 75% were used to produce the BC<sub>1</sub>F<sub>2</sub> generation by self-pollination.


In total, 188 BC<sub>1</sub>F<sub>2</sub> plants were analysed by foreground selection and phytopathology tests (Table 1). To detect the *Lr39* gene in the BC<sub>1</sub>F<sub>2</sub> plants, 5 molecular markers were used: *Gdm35*, *Barc124*, *Gwm261*, *Gwm296*, and *Gwm210*. With the primers for marker *Gdm35*, a 170 bp DNA fragment from the line KS90WGRC10 was amplified in 47 plants (Table 1). The primers for *Barc124* amplified a 250 bp fragment in 42 plants, whereas the primers for *Gwm261* amplified a 165 bp fragment in 76 BC<sub>1</sub>F<sub>2</sub> plants. The primers for *Gwm296* amplified a 135 bp fragment in 15 plants, and *Gwm210* a 182 bp fragment in 27 plants (*Lr39* + *Pm21*) (Table 1). The marker for the *Pm21* gene, *NAU/xibao*, was detected in 111 plants (Table 1). Phytopathology tests using a natural population of *P. triticina* revealed that 151 plants were resistant to this pathogen, whereas 162 BC<sub>1</sub>F<sub>2</sub> plants were resistant to *B. graminis*. On the basis of the results from the foreground selection and phytopathology tests, 45 plants that carried both *Lr39* and *Pm21* were selected for production of the BC<sub>1</sub>F<sub>3</sub> generation by self-pollination (Table 1).

In total, 742 BC<sub>1</sub>F<sub>3</sub> plants were analysed by phytopathology tests using natural populations of *P. triticina* and *B. graminis* (Table 1). Four hundred and seventy-one BC<sub>1</sub>F<sub>3</sub> plants were found to be resistant to leaf rust. In addition, 642 plants were found to be resistant to powdery mildew (Table 1). A total of 382 plants were analysed by foreground selection to detect the *Lr39* and *Pm21* genes. The primers for the marker *Barc124* amplified a 250 bp fragment in 200 plants, those for *Gdm35* amplified a 170 bp fragment in 184 plants, whereas the primers for *Gwm261* amplified a 165 bp fragment in 150 plants (Table 1). In addition, the primers for *Gwm296* amplified a 135 bp fragment in 165 plants and a 182 bp fragment for *Gwm210* was amplified in 135 plants (Table 1). Furthermore, the primers for the marker *NAU/xibao* amplified a 902 bp fragment in 262 plants (Table 1). On the basis of the molecular and phenotypic characteristics, a total of 66 homozygous plants carried both *Lr39* and *Pm21* in a single genotype, and were selected for field evaluation (Table 1).

Linkage analysis showed that the *Lr39* gene was linked to the 5 microsatellite markers *Gdm35*, *Barc124*, *Gwm261*, *Gwm296*, and *Gwm210*. Mapping of these markers showed that *Barc124* was linked most closely to *Lr39*, at a distance of 9.7 cM. The order of the other markers on the 2DS chromosome was *Gdm35*, *Gwm261*, *Gwm296*, and *Gwm210* at 14.8 cM, 20.6 cM, 26.0 cM, and 33.2 cM, respectively, from the locus of resistance *Lr39* (Figure 2).

The results described herein demonstrate that the success of gene pyramiding depends on the combination of molecular selection on the basis of DNA markers with phenotypic selection, both of which contribute significantly to the efficiency of the selection process. During foreground selection to detect the *Lr39* gene in the composite F<sub>1</sub> generation (crosses 2) (Figure 1), 3 microsatellites were used: *Gdm35*, *Barc124*, and *Gwm210*. On the basis of the results obtained for the composite F<sub>1</sub> generation, we applied additional polymorphic markers that are linked more closely to locus *Lr39*, according to the microsatellite consensus map constructed by SOMERS *et al.* (2004), namely *Gwm261* and *Gwm296*. Therefore, subsequently, 5 microsatellite markers were used for foreground selection: *Gdm35*, *Barc124*, *Gwm210*, *Gwm261*, and *Gwm296*. The use of these specific molecular markers to detect *Lr39* in breeding material was based on previous reports. SINGH *et al.* (2004) used only 3 markers (*Gdm35*, *Barc124*, *Gwm210*) to perform successful foreground selection for *Lr39*. On the basis of the findings of SUN *et al.* (2009), other combinations of microsatellites could be used potentially for marker-assisted selection (MAS), such as *Barc124*, *Gwm210*, *Gdm35*, and *Cfd36*. From these earlier reports of the identification of *Lr39* resistance genes in breeding material, it can be concluded that there might be no universal marker that is suitable for MAS of *Lr39* in breeding programs.

The effectiveness of MAS and the suitability of individual markers depend on the parents used in the breeding program. SUN *et al.* (2009) proposed two SSR markers that can be used in MAS separately, namely *Gdm35* and *Barc124*. When the American cultivar Thunderbolt or line KS93U62 (Century\*3/TA2460) is used as the source of the *Lr39* resistance gene, both above-mentioned markers can be applied, but when the line TX01V5719 is used, only *Barc124* can be applied. If the *Lr39* donor is another cultivar, such as Cardinal, Duster



<sup>1</sup>FS = Foreground Selection; BS = Background Selection; RT = Resistance Test(s)

Figure 1. Experimental design for the pyramiding of two resistance genes for leaf rust and powdery mildew

or Foster etc., *Barc124* is not recommended for MAS (SUN *et al.* 2009). *Gdm35* is also particularly useful for MAS for *Lr39* when the KS90WGRC10/TAM 107 population is used as the donor of *Lr39* (SINGH *et al.* 2004).

In the study reported herein, to interpret the results for the BC<sub>1</sub>F<sub>2</sub> generation better (Figure 2), a genetic map was constructed using 5 microsatellites. This linkage map was compared with the genetic maps obtained by SOMERS *et al.* (2004) (map A), SINGH *et al.* (2004) (map C), and SUN *et al.* (2009) (map D) (Figure 2). The locations of markers relative to the locus *Lr39* differed among the maps. For the map obtained with the BC<sub>1</sub>F<sub>2</sub> generation, the order of the molecular markers on chromosome 2DS was as follows: *Barc124* (9.7 cM to *Lr39*), *Gdm35* (14.8 cM), *Gwm261* (20.6 cM), *Gwm296* (26.0 cM), and *Gwm210* (33.2 cM). According to the map of SINGH *et al.* (2004), the closest marker to the resistance locus was *Gdm35* (1.9 cM to *Lr39*), followed by *Barc124* (5.8 cM) and *Gwm210* (7.9 cM). For the map constructed by SUN *et al.* (2009), the closest marker to *Lr39* was *Barc124* (1.0 cM to *Lr39*), followed by *Gwm210* (1.5 cM), *Gdm35* (2.8 cM), and *Cfd36* (4.1 cM). The differences in the locations of the molecular markers among the different maps suggest that the resistance gene *Lr39* might sometimes be difficult or problematic to detect in breeding material. In MAS, selection for the presence of the *Lr39* gene using only three microsatellites might not be sufficient for plant breeding. As a consequence, to increase the efficiency of selection of the *Lr39* gene in this study, 5 microsatellite markers were

used: *Gdm35*, *Barc124*, *Gwm210*, *Gwm261*, and *Gwm296*. On the basis of our results, it is not possible to recommend one universal marker that is suitable for detection of the *Lr39* gene in large-scale breeding programs.

In this study, a total of 3 molecular markers were used to detect *Pm21* in breeding material, *SCAR*<sub>1250'</sub>, *SCAR*<sub>1400'</sub>, and co-dominant PCR marker *NAU/xibao*, all of which appeared to be suitable for




Figure 2. Partial map of wheat chromosome 2DS (B) including resistance gene *Lr39* based on the analysis of the BC<sub>1</sub>F<sub>3</sub> population *Lexus* × *KS90WGRC10*; the order of the markers is compared with reference maps from (A) SOMERS *et al.* 2004, (C) SINGH *et al.* 2004, and (D) SUN *et al.* 2009

MAS. Similar results were reported by LIU *et al.* (1999), YILDIRIM *et al.* (2004), and CAO *et al.* (2006). Other markers were previously used to detect *Pm21* and identify homozygous and heterozygous translocation in breeding materials, these include specific RAPD markers, such as *OPH17-1900* and *OPH17-1000* (QI *et al.* 1996). In addition, LIU *et al.* (2000) used the RFLP marker *Psr113* to select *Pm21*. The most frequently encountered obstacles for MAS are the high cost and time requirements of RFLP technologies, and the low repeatability of RAPD markers. As a consequence, these types of marker are no longer used commonly for MAS.

One of the benefits of using molecular markers in breeding programs is the potential to estimate the proportion of alleles from the parental lines in the genomes of the offspring. Such information is particularly important in backcross breeding programs. Compared with conventional backcrossing, selection based on DNA markers (such as microsatellites and AFLPs) can speed up recovery of the recurrent parent (RP) genome (RIBAUT & HOISINGTON 1998; SERVIN & HOSPITAL 2002). This strategy is highly profitable after the first backcrosses in BC<sub>1</sub>F<sub>1</sub> generations, when the widest variation for the RP genome content is observed and the individuals with the highest RP genome content can be selected, as was demonstrated in this study. Microsatellite and AFLP markers are used commonly to estimate genetic similarity, and they are effective tools for backcross breeding programs. In the present study, a set of 5 AFLP primer combinations was used to estimate genetic similarity among the backcross lines and it was estimated to be in the range of 32.1–70.8%. CHEN *et al.* (2001) estimated genetic similarity using a set of 10 AFLP primer combinations, and found the recovery percentages for the recipient genome to be approximately 92.25% (BC<sub>1</sub>F<sub>1</sub>), 97.67% (BC<sub>2</sub>F<sub>1</sub>), and 98.8% (BC<sub>3</sub>F<sub>1</sub>). In a similar study, which was based on 35 microsatellites and 12 AFLP primer combinations, the range of % RP in BC<sub>1</sub>F<sub>3</sub> was found to be 94.3–97% (GUPTA *et al.* 2008).

In summary, gene pyramiding is a breeding strategy that enables multiple resistance genes to be introduced into a single genotype, which results in cultivars that are resistant to one or more diseases. Using foreground and background selection to introduce resistance genes considerably reduces the time required to breed new cultivars. The use of molecular selection in conjunction with resistance tests confers many benefits for MAS,

especially when the foreground selection is difficult to interpret. In this study, we obtained new homozygous lines that were resistant to leaf rust and powdery mildew. These lines could be used as a source of resistance to these pathogens in wheat breeding programs.

**Acknowledgements.** Supported by the UE BIOEXPLOIT project (FOOD-CT-2005-513959) and Polish Ministry of Science and Higher Education (Grant No. NN310 147035).

## References

- CAO A.Z., WANG X.E., CHEN Y.P., ZOU X.W., CHEN P.D. (2006): A sequence-specific PCR marker linked with *Pm21* distinguishes chromosomes 6AS, 6BS, 6DS of *Triticum aestivum* and 6VS of *Haynaldia villosa*. *Plant Breeding*, **125**: 201–205.
- CHEN S., XU C.G., LIN X.H., ZHANG Q. (2001): Improving bacterial blight resistance of 6078, an elite restorer line of hybrid rice, by molecular marker-assisted selection. *Plant Breeding*, **120**: 133–137.
- GUPTA P.K., BALYAN H.S., KUMA R.J., KULWAL P.L., KUMAR N., MIR R.R., KUMAR A., MOHAN A., PRABHU K.V. (2008): QTL analysis and marker assisted selection for improvement in grain protein content and pre-harvest sprouting tolerance in bread wheat. In: Proc. 11<sup>th</sup> Int. Wheat Genetics Symp. August 24–29, Brisbane.
- HIEBERT C.W., THOMAS J.B., MCCALLUM B.D., SOMERS D.J. (2010): Genetic mapping of the wheat leaf rust resistance gene *Lr60* (LrW2). *Crop Science*, **48**: 1020–1026.
- HIGGINS C.M., HALL R.M., CAMPBELL P.R., DIETZGEN R.G. (2000): PCR rescue and analysis of transgene sequences directly from crude extracts of transgenic embryos and plants. *Plant Molecular Biology Reporter*, **18**: 285a–285g.
- KOSAMBI D.D. (1943): The estimation of map distance from recombination values. *Annals of Human Genetics*, **12**: 172–175.
- LANDER E.S., GREEN P., ABRAHAMSON J., BARLOW A., DALY M.J., LINCOLN S.T., NEWBURG L. (1987): Mapmaker: An interactive computer package for construction primary genetic linkage maps of experimental and natural populations. *Genomics*, **1**: 174–181.
- LEVINE M.N., CHEREWICK W.J. (1952): Studies on Dwarf Leaf Rust on Barley. USDA Technical Bulletin 1056, USDA, Washington D.C., 1–17.
- LIU J., LIU D., TAO W., LI W., WANG S., CHEN P., CHENG S., GAO D. (2000): Molecular marker-facilitated pyramiding of different genes for powdery mildew resistance in wheat. *Plant Breeding*, **119**: 21–24.

- LIU Z., SUN Q., NI Z., YANG T. (1999): Development of SCAR markers linked to the *Pm21* gene conferring resistance to powdery mildew in common wheat. *Plant Breeding*, **118**: 215–219.
- MAINS E.B., DIETZ S.M. (1930): Physiologic forms of barley mildew, *Erysiphe graminis hordei* Marchal. *Phytopathology*, **20**: 229–239.
- PESTSOVA E., GANAL M.W., RÖDER M.S. (2000): Isolation and mapping of microsatellite markers specific for the D genome of bread wheat. *Genome*, **43**: 689–697.
- QI L., CAO M., CHEN P., LI W., LIU D. (1996): Identification, mapping, and application of polymorphic DNA associated with resistance gene *Pm21* of wheat. *Genome*, **39**: 191–197.
- RIBAUT J.M., HOISINGTON D. (1998): Marker-assisted selection: new tools and strategies. *Trends in Plant Science*, **3**: 236–239.
- RÖDER M.S., KORZUN V., WENDEHAKKE K., PLASCHKE J., TIXIER M.H., LEROY P., GANAL M.W. (1998): A microsatellite map of wheat. *Genetics*, **149**: 2007–2023.
- SERWIN B., HOSPITAL F. (2002): Optimal positioning of markers to control genetic background in Marker Assisted Backcrossing. *The Journal of Heredity*, **93**: 214–217.
- SINGH S., FRANKS C.D., HUANG L., BROWN-GUEDIRA G.L., MARSHALL D.S., GILL B.S. (2004): *Lr41*, *Lr39*, and a leaf rust resistance gene from *Aegilops cylindrica* may be allelic and are located on wheat chromosome 2DS. *Theoretical and Applied Genetics*, **108**: 586–591.
- SOMERS D.J., ISAAC P., EDWARDS K. (2004): A high-density microsatellite consensus map for bread wheat (*Triticum aestivum* L.). *Theoretical and Applied Genetics*, **109**: 1105–1114.
- SONG Q.J., SHI J.R., SINGH S., FICKUS E.W., COSTA J.M., LEWIS J., GILL B.S., WARD R., CREGAN P.B. (2005): Development and mapping of microsatellite (SSR) markers in wheat. *Theoretical and Applied Genetics*, **110**: 550–560.
- SUN X., BAI G., CARVER B.F. (2009): Molecular markers for wheat leaf rust resistance gene *Lr41*. *Molecular Breeding*, **23**: 311–321.
- VOS P., HOGERS R., BLEEKER M., REIJANS M., VAN DE LEE T., HORNES M., FRITERS A., POT J., PALEMEN J., KUIPER M., ZABEAU M. (1995): AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research*, **23**: 4407–4414.
- YILDIRIM A., SAKIN M., KARADAG Y., GOKMEN S., KANDEMIR N., AKKAYA M.S., YILDIRIM F. (2004): Genetic marker mediated transfer of an alien gene, *Pm21*, into wheat conferring resistance to powdery mildew. *Biotechnology & Biotechnological Equipment*, **18**: 15–19.

Received for publication July 30, 2012  
Accepted after corrections May 27, 2013

---

*Corresponding author:*

DR. ALEKSANDRA PIETRUSIŃSKA, National Research Institute at Radzików, Plant Breeding and Acclimatization Institute, Department of Plant Breeding and Genetics, 05-870 Błonie, Poland  
e-mail: a.pietrusinska@ihar.edu.pl

---