

Conception rate after sex determination and cryopreservation of D7 bovine embryos

M. LOPATAROVA¹, S. CECH¹, P. KRONTORAD², L. HOLY¹, H. LALOVA¹, R. DOLEZEL¹

¹University of Veterinary and Pharmaceutical Sciences, Brno, Czech Republic

²BOVET Company, Sloupnice, Czech Republic

ABSTRACT: The aim of this study was to evaluate conception rate after sex determination and subsequent freezing of Day 7 (D₇) bovine embryos and to compare it with transfer of fresh female embryos. High quality embryos obtained from superovulated donors were biopsied and the embryonic cells (5–10) were analysed by PCR using specific primers for the Y-chromosome determinant. Fresh embryos were blade biopsied ($n = 172$) and subsequently transferred (ET) ipsilaterally to synchronized recipients. Selected embryos ($n = 112$) were biopsied by cell aspiration and cryopreserved for later transfer. Sex determination was successfully completed in 91.3% of fresh group embryos (44% female) and in 87.5% of freeze-thawed group embryos (45.9% female). The achieved pregnancy rate was 56.5% after fresh transfer of sexed (female) embryos whereas intact embryos (control 1) implanted in 61.9% recipients. 95.6% (43/45) of embryos survived biopsy and cryopreservation. After transfer of freeze-thawed sexed (female) embryos, pregnancy was established in 48.8% of recipients. A similar conception rate was found in intact embryos (50.7%, control 2). The results clearly demonstrate that the microsurgical technique used and subsequent PCR sex analysis allow the rapid commercial exchange of genetic resources on the basis of fresh or frozen sex-desired embryos in embryo transfer programs.

Keywords: cattle; superovulation; sexing; polymerase chain reaction; cryopreservation; embryo transfer

Embryo transfer (ET) technology is of high importance in modern cattle breeding programs. Improvements in embryo micromanipulation techniques have led to the use of direct genetic analysis of preimplantation bovine embryos by biopsy in commercial embryo transfer programs. Implementation of embryo sex determination techniques holds great potential for genetic improvement in cattle herds and market demand satisfaction.

The techniques for sex determination of bovine embryos have evolved from karyotyping of older preimplantation embryos (Hare et al., 1976), to the current variety of widely used polymerase chain reaction (PCR) protocols. The first serious attempt to sex embryos was made by Edwards and

Gardner (1967) with rabbit blastocysts according to the presence of Barr bodies. Time consuming methods were used later in farm animals with high variability of results. Employed methods include cytogenetic analysis (Singh and Hare, 1980; Rall and Leibo, 1987), immunological assays (White et al., 1982; Booman et al., 1989), detection of metabolic differences between male and female embryos (Williams, 1986; Monk and Handyside, 1988), analysis of chromatin with Y-specific DNA probes (Leonhard et al., 1987; Bondioli et al., 1989; Kobayashi et al., 1998) and analysis on the base of differences in cleavage rates (Avery et al., 1989). Better results were published later using PCR amplification of specific DNA sequences to determine embryonic sex in cattle (Herr et al., 1990;

Supported by the Ministry of Education, Youth and Sports of the Czech Republic (Grant No. MSM 6215712403).

Schroder et al., 1990; Peura et al., 1991), pigs (Pomp et al., 1995), horses (Peippo et al., 1995), humans (Handyside et al., 1990) and mice (Han et al., 1993). Recently published results support the suitability of the PCR method for sex determination in cattle on the basis of its high accuracy and quick results (Thibier and Nibart, 1995; Lopes et al., 2001; Ekici et al., 2006; Yu et al., 2007).

Sperm sexing and separation into X- and Y-bearing fractions for purposes of insemination or *in vitro* production of embryos (IVP) seems to be a promising procedure in sex determination. However, recent results show that the capacitation of sorted freeze-thawed fractions of sperm and the developmental capacity of IVP embryos with desired sex is very low (Zhang et al., 2005; Katska-Ksiazkiewicz et al., 2006; Blondin et al., 2009). The pregnancy rates with low doses of sexed sperm drop about 10–40% compared to conventional doses of unsexed sperm (Andersson et al., 2006; Schenk et al., 2006, 2009; Seidel Jr., 2007; DeJarnette et al., 2009). In superovulation programs, the numbers of recovered transferable embryos were significantly higher when donors were inseminated with the unsexed sperm than with the sexed inseminates (Sartori et al., 2004; Schenk et al., 2006; Hayakawa et al., 2009).

The aim of this study is to estimate the validity of a protocol comprising biopsy, to isolate embryonic cells for sex determination of high quality embryos, freeze-thawing of sex-desired embryos and subsequent ipsilateral transfer into recipients. The implementation of freeze-thawed high quality embryos of desired sex in commercial ET programs is discussed.

MATERIAL AND METHODS

Animals and treatment

Selected donors ($n = 56$) were stimulated according to the protocol described by Holy et al. (1990). Briefly, Holstein-Friesian cows were superovulated between days 8–12 of the oestrous cycle with eight decreasing doses of 500 IU of FSHp and 500 IU of LHp *pro toto* (Pluset[®], Calier, Spain) administered at 12-h intervals. Oestrus was induced by double prostaglandin $F_{2\alpha}$ ($PGF_{2\alpha}$) treatment (cloprostenol, Oestrophan[®], Bioveta, Czech Republic) together with the fifth and sixth FSH injections. Three artificial inseminations were performed at 48, 60 and 72 h after the first application of $PGF_{2\alpha}$.

Embryo collection and scoring

Embryos were flushed with PBS (Dulbecco's Phosphate Buffered Saline, Live Technologies, Ltd., U.K.) + 1% FCS (foetal calf serum, Biochrom AG, Germany) on Day 7 after the first insemination (D_0). Flushing, as well as embryo isolation, was carried out as described by Holy et al. (1990). Obtained embryos were washed with Vigro[™] Holding Plus medium (AB Technology, Inc., Pullman, WA) and classified according to their developmental stage and quality with respect to internationally accepted criteria (Robertson and Nelson, 1998). Only high quality compacted morulae, early blastocysts and blastocysts ($n = 284$) were selected for microsurgical procedures. The designation of high quality




Figure 1. Preparation of pipettes and fixation of the compacted morula before biopsy by aspiration




Figure 2. Aspiration of a few surface embryonic cells from the compacted morula

was assigned to morphologically excellent (perfect embryo for its age) or good (trivial imperfections such as oval *zona pellucida*, a few small excluded cells, or slightly asymmetrical shape) embryos.

Embryo biopsy

Microsurgical intervention was carried out with the Labovert micromanipulator (Leitz, Austria). From the embryos specified for fresh transfer, a small portion of the embryo was cut off by transzonal incision using a microsurgical blade as described (Lopatarova et al., 2007). No holding pipette was used during cutting, only PBS without proteins was employed for fixation of embryos. The embryos selected for subsequent cryopreservation were transferred into fresh holding medium and immobilized using the holding pipette (outer diameter 80 μm , inner diameter 15 μm , Microtech, Czech Republic). From compacted morulae (Figures 1 and 2), a few surface embryonic cells (5–10) were removed by careful aspiration with a micro-injection pipette (outer diameter 7 μm , inner diameter 5 μm , bevel angle 50°). Care was taken to ensure that penetration of the *zona pellucida* was not accompanied by large damage. In the blastocysts, the trophoblast cells were aspirated (5–10). The isolated embryonic cells (Figure 3) were analysed immediately. During the analysis (3.0–3.5 h), the embryos were cultured in 4-wells dishes containing 1 ml holding medium per well at ambient temperature (20–25°C).

Sex determination

The harvested cells were immediately transferred to microtubes containing Cresol red (o-Cresolsulfonaphthalein). Embryo sexing was performed with a commercial polymerase chain reaction (PCR) kit using primers specific for the Y-chromosome determinant (YCD) according to the manufacturer's instructions (Herr et al., 1995). The PCR product was detected by UV light in agarose gel with ethidium bromide (10 mg/ml in distilled water) and the embryos were scored as Y-chromosome determinant positive (male) or Y-chromosome determinant negative (female), respectively. All instruments represent a transferable unit that can be used in field conditions (AB Technology, Inc., Pullman, WA).

Following completion of the procedure (about 3.0–3.5 h after cell sample isolation), the female embryos were fresh transferred or frozen. The results of the PCR test and sex ratio in blade biopsied embryos were compared with those that were aspiration biopsied.

Embryo freezing and thawing

The female embryos ($n = 45$) intended for cryopreservation were frozen using a conventional freezing method in 1.5M ethylene glycol with 0.1M sucrose (Vigro™ Ethylene Glycol Freeze Plus, AB Technology, Inc., Pullman, WA) as described by Voelkel and Hu (1992) using the HAAKE F 4Q freezer unit. The rapid thawing procedure was

Table 1. Sex determination in D₇ high quality bovine embryos before fresh and freeze-thawed ET

Mode of embryo transfer	Method of biopsy	Embryos analyzed (n)	Successful sex determination (n/%)	Female sex confirmed (n/%)
Fresh	blade	172	157/91.3 ^a	69/44 ^c
Frozen/thawed	aspiration	112	98/87.5 ^b	45/45.9 ^d

a : b, c : d = $P > 0.05$

done according to Voelkel and Hu (1992) and the embryos were held for a short time in holding medium before transfer. Post-thawing embryo viability was evaluated.

Embryo transfer

After sex determination, the fresh female ($n = 69$) and freeze-thawed viable embryos ($n = 43$) were transferred ipsilaterally into synchronized recipient heifers (weight 340–370 kg). In the frame of the routine ET schedule, the intact high quality fresh ($n = 105$) and freeze-thawed embryos ($n = 75$) formed control groups (control 1 and 2) under identical conditions (during the same period on same dairy farm). Pregnancy diagnosis was performed ultrasonographically on Day 21 after transfer and confirmed by rectal palpation 7–21 days later.

Statistical analysis

The obtained data were analysed by the χ^2 test (2×2 contingency tables). The results of the analyses are summarized in Tables 1–3.

RESULTS

Table 1 summarizes the efficiency of sex determination with respect to the method of embryonic cell biopsy. A total of 172 blade biopsied embryos were selected for fresh transfer and sex analysis was successfully carried out in 157 of them (91.3%). Female sex was confirmed in 69 embryos (44%). Similarly, the analysis of 112 embryos biopsied by aspiration and selected for freeze-thaw transfer was successfully completed in 98 instances (87.5%) and 45 of them (45.9%) were female.

Pregnancy rates of the sexed fresh embryos and the intact fresh embryos (control 1) are compared in Table 2. After the transfer of 69 female embryos, 39 animals became pregnant (56.5%). The conception rate of 61.9% after ET of intact control embryos was not significantly different (65/105). Embryo biopsy, sexing and cryopreservation did not significantly influence conception rate in comparison with the intact cryopreserved embryos (control 2, Table 3). After the thawing of 45 female embryos, 43 of them were eligible for ET (95.6%) and 48.8% of them (21) became pregnant. After ET of the control embryos (control 2), 50.7% of recipients became pregnant (38/75).



Figure 3. The clump of cells isolated during biopsy and subjected to sexing

Table 2. Pregnancy rate achieved after fresh ET of D₇ bovine sexed (female) versus unsexed (control 1) embryos

Embryo biopsy and sexing	Embryos transferred (<i>n</i>)	Pregnant animals (<i>n</i> %)
Yes	69	39/56.5 ^a
Not (control 1)	105	65/61.9 ^b

a : b = $P > 0.05$

Table 3. Pregnancy rate achieved after freeze-thawed ET of D₇ bovine sexed (female) versus unsexed (control 2) embryos

Embryo biopsy and sexing	Freeze-thawed embryos (<i>n</i>)	Transferred of surviving embryos (<i>n</i> %)	Pregnant animals (<i>n</i> %)
Yes	45	43/95.6 ^a	21/48.8 ^c
Not (control 2)	75	75/100 ^b	38/50.7 ^d

a : b, c : d = $P > 0.05$

These data indicate that ET of sexed-fresh and sexed-cryopreserved embryos represents a very efficient approach for achieving a high number of calves with desired sex (Figure 4).

DISCUSSION

The method of sex determination in bovine embryos using embryonic cell biopsy and DNA analysis represents a useful tool for influencing the sex of calves. The technique expands to include other DNA markers associated with valuable production traits. Nowadays it can be used for multiplex

genotyping of bovine embryos (Peippo et al., 2007; Hirayama et al., 2008; Alonso et al., 2009) and for genetic testing for hereditary diseases (Hirayama et al., 2004; Ipate et al., 2007).

In our previous experiments (Lopatarova et al., 2007, 2008) for isolation of embryonic cells we used the blade biopsy for fresh embryo transfer. In this study, we have introduced the aspiration procedure to the biopsy of embryonic cells which can retain embryos with high developmental competence and minimize the damage to the *zona pellucida* before the freeze-thaw procedure. A glass holding pipette stabilizes the embryo by negative pressure while the other pipette performs the biopsy. An embryo-




Figure 4. Pregnancy rate after sexing and cryopreservation of bovine D₇ embryos

onic cell aspiration procedure for *in vitro*-produced (IVP) embryos was applied by Lopes et al. (2001) using two micromanipulator units. Shea (1999) used three methods for biopsy of whole embryos but no differences were found between microsurgical blade, aspiration and fine glass needle in completion of sex determination (86–92%) as well as in pregnancy rates (49–60%), achieved after transfers of fresh embryos. However, frozen blade biopsied embryos, usually *zona*-free embryos, show lower (23%) pregnancy rates after transfer in comparison with needle biopsy (41%; Shea, 1999). In contrast, the present experiments demonstrated better pregnancy rates with the aspiration method and freeze-thawing of embryos (48.8%). In IVP embryos biopsied by aspiration, the mortality rate was 10.3% and the pregnancy rate was 31.7% after ET of fresh embryos (Alonso et al., 2009).

In this study, the gender was determined in 91.3% of embryos for fresh transfer and in 87.5% of embryos intended for freezing (Table 1), which accords with our previous work with whole, intact (88.7%; Lopatarova et al., 2007) and splitted embryos (89.4–92%; Lopatarova et al., 2008). Our results are similar to those published by other authors (85–95%) using embryos from superovulated donors (Thibier and Nibart, 1995; Shea, 1999; Li et al., 2007; Yu et al., 2007) as well as IVP embryos (Lopes et al., 2001; Hasler et al., 2002; Tominaga and Hamada, 2004). However, the successful sex detection was inversely related to the number of isolated embryonic cells (≤ 3 cells 85.5%, 4–6 cells 97.4%, ≥ 7 cells 100%; Lacaze et al., 2008).

In this study we confirmed female sex in 44 and 45.9% of embryos (Table 1). Shea (1999) and Lacaze et al. (2008) reported a similar rate (44–48%) in all stages of D₇–D_{7.5} embryos. Hasler et al. (2002) even showed that the sex ratio of expanded blastocysts was significantly skewed in favour of female embryos (60.3%). They further proved that the overall percentage of male IVP embryos was significantly higher (53%) than the percentage of male *in vivo* produced embryos (49.2%). King et al. (1992) and Gutierrez-Adan et al. (2001) found that male embryos were more resistant to manipulation and cultivation. Nedambale et al. (2004) recovered IVP embryos from culture and then after vitrification. After warming, the embryos were cultured for 48 h and sexed by PCR. The embryos that reached the blastocyst stage earlier were predominantly males and these male embryos had better morphology and subsequently hatched at a greater rate than female embryos. PCR analysis for sexing of *in vivo*

produced D₇ embryos indicated that male embryos developed faster and were graded higher than female embryos (Tominaga, 2004). The sex ratio was significantly shifted to the male IVP embryos also in more recent publications (Kimura et al., 2008; Alonso et al., 2009).

Comparison of ethylene glycol or glycerol as cryoprotectants did not show any influence on the pregnancy rate (Shea, 1999; Lopes et al., 2001). We used only 1.5M ethylene glycol for a rapid and effective thawing procedure. Hasler et al. (2002) first froze embryos in ethylene glycol or glycerol and biopsies were carried out after thawing. The freeze-thawing-biopsy protocol rendered again, with both cryoprotectants, satisfactory pregnancy rates (about 50%). Also the ultra-rapid vitrification, using the gel-loading tip (Tominaga, 2004), is acceptable for embryos that underwent the biopsy and method is available to practical use in the field.

The experiments with IVP blastocysts showed a similar post-thaw survival rate for biopsied (92.3%) and intact (96.2%) embryos after slow cooling in 0.7M glycerol plus 0.1M sucrose, respectively (Tominaga et al., 2007). In our study, 95.6% of embryos survived the biopsy, *in vitro* culture during the sex analysis (up to 3.5 hours) and freezing procedures. In the control group (control 2), all intact embryos were viable after thawing and could be transferred to the recipients (Table 3).

However, the IVP bovine embryos are more sensitive to these procedures than those derived *in vivo*, probably due to accumulation of lipids during culture (Pereira et al., 2008). These authors proved that the resistance of IVP embryos to biopsy and following vitrification was significantly improved by supplementing bovine serum-containing media with trans-10, cis-12 conjugated linoleic acid (CLA) that decreases embryo cytoplasmic lipid deposition during *in vitro* culture. Embryos cultivated in CLA (CLA embryos) show higher post-warming survival rate (95%) and expansion rate (64.6% after 24 h of culture) than embryos cultured in conventional media (62.5 and 27.5%, respectively).

The viability of fresh biopsied embryos after ET varies from 49 to 62% (Lopes et al., 2001; Hasler et al., 2002; Li et al., 2007) using the various mode of biopsy. Our result in pregnancy rate (Table 2) was comparable with published data and does not differ between the pregnancy rates after ET of fresh intact (control 1, 61.9%) and sexed embryos (56.5%).

We obtained a very similar pregnancy rate (Table 3) comparing experimental (biopsied and

frozen) embryos (48.8%) with control 2 (intact and frozen) embryos (50.7%). In contrast, Hasler et al. (2002) reported that transfer of the frozen embryos in ethylene glycol produced a lower pregnancy rate after biopsy (49.8%) than did intact embryos (65.6%). In the study of Lacaze et al. (2008), the pregnancy rate after transfer of freeze-thawed biopsied female embryos averaged 47.2% (corresponding to our study) and the result was mainly influenced by the parity of recipients (heifers 51.7% vs. cows 26.3%).

In our study, no significant differences were found between the pregnancy rates of biopsied fresh (56.5%) and freeze-thawed (48.8%) embryos. However, Shea (1999) reported a great difference between pregnancy rates after the transfer of biopsied non-frozen (58%) and in glycerol (37%), or in ethylene glycol (37%), frozen embryos. Li et al. (2007) observed reduced pregnancy rates in a large number of animals (41.8%) with embryos used for biopsy and subsequently exposed to freezing and thawing procedures, whereas the implantation of fresh embryos after biopsy did not affect pregnancy rate (49.6%) compared to non-biopsied fresh and freeze-thawed embryo groups (52.9 and 46.6%, respectively).

Comparing the efficiency of embryo sexing versus sperm sexing method, recent publications report worse results with sexed sperm compared to unsexed sperm, when X chromosome-bearing fractions of sorted sperm were used for IVF (Blondin et al., 2009), for insemination (Schenk et al., 2009) and in multiple ovulation and embryo transfer (MOET) programs (Hayakawa et al., 2009; Peippo et al., 2009). The use of low-dose X-sorted spermatozoa for the insemination of superovulated embryo donors can improve the proportion of transferable female embryos produced but this potential may be not realised in commercial bovine practice, due to reduced fertilization rates using X-sorted spermatozoa (Peippo et al., 2009). The lower effectiveness may be due to procedures associated with sorting, handling of sperm, bull selection, instrumentation, mode of insemination, management level, etc. (Garner and Seidel Jr., 2008).

In conclusion, our results document that the well established gentle isolation of the lowest number of embryonic cells by microsurgical biopsy of embryos, an efficient and accurate PCR method, and careful cryopreservation provides for a high number of reliably sexed and freeze-thawed viable embryos that successfully implant after ET (Figure 4). The

described method allows, under field conditions, the production of a high number of calves of desired gender and improves the genetic potential of cattle herds in a short time interval.

REFERENCES

- Alonso RV, Hellu JAA, Perri SHV, Visintin JA, Garcia JF (2009): Factors affecting commercial sexing program and multiple genetic analysis perspectives of in vitro-produced bovine embryos. *Reproduction, Fertility and Development*, 21, 227.
- Andersson M, Taponen J, Kommeri M, Dahlbom M (2006): Pregnancy rates in lactating Hostein-Friesian cows after artificial insemination with sexed sperm. *Reproduction in Domestic Animals*, 41, 95–97.
- Avery B, Bak A, Schmidt M (1989): Different cleavage rates and sex determination in bovine embryos. *Theriogenology*, 32, 139–147.
- Blondin P, Beaulieu M, Fournier V, Morin N, Crawford L, Madan P, King WA (2009): Analysis of bovine sexed sperm for IVF from sorting to the embryo. *Theriogenology*, 71, 30–38.
- Bondioli KR, Ellis SB, Pryor JH, Williams MW, Harpold MM (1989): The use of male-specific chromosomal DNA fragments to determine the sex of bovine pre-implantation embryos. *Theriogenology*, 31, 95–104.
- Booman P, Kruijt L, Veerhuis R, Hengst AM, Tiemann M, Ruch FE (1989): Sexing bovine embryos with monoclonal antibodies against the H-Y antigen. *Livestock Production Science (Niederlande)*, 23, 1–16.
- DeJarnette JM, Nebel RL, Marshall CE (2009): Evaluating the success of sex-sorted semen in US dairy herds from on farm records. *Theriogenology*, 71, 49–58.
- Edwards RG, Gardner RL (1967): Sexing of live rabbit blastocyst. *Nature*, 214, 576–577.
- Ekici H, Turan N, Sontas BH, Helps CR, Senunver A, Yilmaz H (2006): Sex determination of bovine embryos using polymerase chain reaction (PCR). *Revue de Medicine Veterinaire*, 157, 441–444.
- Garner DL, Seidel GE Jr. (2008): History of commercializing sexed semen for cattle. *Theriogenology*, 69, 886–895.
- Gutierrez-Adan A, Lonergan P, Rizos D, Ward FA, Boland MP, Pintado B, de la Fuente J (2001): Effect of the in vitro culture system on the kinetics of blastocyst development and sex ratio of bovine embryos. *Theriogenology*, 55, 1117–1126.
- Han Y, Yoo O, Lee K (1993): Sex determination in single mouse blastomeres by polymerase chain reaction. *Journal of Assisted Reproduction and Genetics*, 10, 151.

- Handyside AH, Kontogiann EH, Hardy K, Winston RML (1990): Pregnancies from biopsied human preimplantation embryos sexed by Y-specific DNA amplification. *Nature*, 344, 768–770.
- Hare WCD, Mitchell D, Betteridge KJ, Eaglesome MD, Randalu GCB (1976): Sexing two week-old bovine embryos by chromosomal analysis prior to surgical transfer: preliminary methods and results. *Theriogenology*, 5, 243–253.
- Hasler JE, Cardey E, Stokes JE., Bredbacka P (2002): Nonelectrophoretic PCR-sexing of bovine embryos in a commercial environment. *Theriogenology*, 58, 1457–1469.
- Hayakawa H, Hirai T, Takimoto A, Ideta A, Aoyagi Y (2009): Superovulation and embryo transfer in Holstein cattle using sexed sperm. *Theriogenology*, 71, 68–73.
- Herr CM, Holt NA, Matthaai KI, Reed KC (1990): Sex of progeny from bovine embryos sexed with a rapid Y-chromosome-detection assay. *Theriogenology*, 33, 247.
- Herr CM, Steel T, Lascelles A, Reed KC (1995): Instruction Manual: Splitting, Biopsy & Rapid Sexing of Cattle Embryos. AB Technology, Inc., Pullmann, WA. 58 pp.
- Hirayama H, Kageyama S, Moriyasu S, Sawai K, Onoe S, Takahashi Y, Katagiri S, Toen K, Watanabe K, Notomi T, Yamashina H, Matzuzaki S, Minamihashi A (2004): Rapid sexing of bovine preimplantation embryo using loop-mediated isothermal amplification. *Theriogenology*, 62, 887–896.
- Hirayama H, Fujikawa A, Kageyama S, Moriyasu S, Sawai K, Onoe S, Minamihashi A (2008): Multiple genotyping in bovine pre-implantation embryo with whole genome amplification. *Animal Science Journal*, 79, 554–560.
- Holy L, Lopatarova M, Krontorad P, Holy J (1990): Systematic use of embryo transfer in cattle under condition of high animal concentration (in German). *Monatsheft Veterinaermedizin*, 45, 409–413.
- Ipate I, Bogdan AT, Toba LG, Toba GF, Ivana S (2007): Research regarding the embryo biopsy for sexing and genetic testing the bovines embryos. *Revista Romana de Medicina Veterinara*, 17, 213–218.
- Katska-Ksiazkiewicz L, Rynska B, Bochenek M, Opiela J, Jurkiewicz J (2006): In vitro production of bovine embryos using flow-cytometrically sexed sperm. *Archiv fur Tierzucht*, 49, 133–140.
- Kimura K, Iwata H, Thompson JG (2008): The effect of glucosamine concentration on the development and sex ratio of bovine embryo. *Animal Reproduction Science*, 103, 228–238.
- King WA, Picard L, Bosquet D, Goff AK (1992): Sex-dependent loss of bisected bovine morulae after culture and freezing. *Journal of Reproduction and Fertility*, 96, 453–459.
- Kobayashi J, Sekimoto A, Uchida H, Wada T, Sasaki K, Sasada H, Umezumi M, Sato E (1998): Rapid detection of male-specific DNA sequence in bovine embryos using fluorescence in situ hybridization. *Molecular Reproduction and Development*, 51, 390–394.
- Lacaze S, Humblot P, Ponsart C (2008): Sexing and direct transfer of bovine biopsied frozen-thawed embryos under on-farm conditions in a commercial program. *Recherches sur les Ruminants*, 15^{eme} Recontre, Paris, les 3 et 4 decembre 2008, 387–390.
- Leonhard M, Korszenbaum M, Cotinot C, Chesne P, Heyman Y, Stinnakre MG, Bishop C, Delouis C, Vaiman M, Fellous M (1987): Sexing bovine embryos using Y chromosome specific DNA probe. *Theriogenology*, 27, 248.
- Li S, Yu W, Fu J, Bai Y, Jin F, Shangguan B (2007): Factors influencing pregnancy rates following transfer of bovine in vivo embryos biopsied for sex determination. *Reproduction, Fertility and Development*, 19, 297.
- Lopatarova M, Cech S, Krontorad P, Holy L, Hlavicova J, Dolezel R (2007): Lower quality bovine embryos may be successfully used for sex determination. *Veterinarni Medicina*, 52, 540–546.
- Lopatarova M, Cech S, Krontorad P, Holy L, Hlavicova J, Dolezel R (2008): Sex determination in bisected bovine embryos and conception rate after the transfer of female demi-embryos. *Veterinarni Medicina*, 53, 595–603.
- Lopes RFF, Forell F, Oliveira ATD, Rodrigues JL (2001): Splitting and biopsy for bovine embryo sexing under field conditions. *Theriogenology*, 56, 1383–1392.
- Monk M, Handyside AH (1988): Sexing of preimplantation mouse embryos by measurement of X-linked gene dosage in a single blastomere. *Journal of Reproduction and Fertility*, 82, 365–368.
- Nedambale TL, Dinnyes A, Yank XZ, Tian XC (2004): Bovine blastocysts development in vitro: timing, sex, and viability following vitrification. *Biology of Reproduction*, 71, 1671–1676.
- Peippo J, Huhtinen M, Kotilainen T (1995): Sex diagnosis of equine preimplantation embryos using the polymerase chain reaction. *Theriogenology*, 44, 619–627.
- Peippo J, Viitala S, Virta J, Raty M, Tammiranta N, Lamminen T, Aro J, Myllymaki H, Vilkki J (2007): Birth of correctly genotyped calves after multiplex marker detection from bovine embryo microblade biopsy. *Molecular Reproduction and Development*, 74, 1373–1378.
- Peippo J, Vartia K, Kananen-Anttila K, Raty M, Korhonen K, Hurme T, Myllymaki H, Sairanen A, Maki-

- Tanila A (2009): Embryo production from superovulated Holstein-Friesian dairy heifers and cows after insemination with frozen-thawed sex-sorted X spermatozoa or unsorted semen. *Animal Reproduction Science*, 111, 80–92.
- Pereira RM, Carvalhais I, Pimenta J, Baptista MC, Vasques MI, Horta AEM, Santos IC, Marques MR, Reis A, Silva Pereira M, Marques CC (2008): Biopsied and vitrified bovine embryos viability is improved by trans10, cis12 conjugated linoleic acid supplementation during in vitro embryo culture. *Animal Reproduction Science*, 106, 322–332.
- Peura T, Hyttinen JM, Turunen M, Janne J (1991): A reliable sex determination assay for bovine preimplantation embryos using the polymerase chain reaction. *Theriogenology*, 35, 547–555.
- Pomp P, Good BA, Geisert RD, Corbin CJ, Conley AJ (1995): Sex identification in mammals with polymerase chain reaction and its use to examine sex effect on diameter of day-10 or -11 pig embryos. *Journal of Animal Science*, 73, 1408–1415.
- Rall WE, Leibo SP (1987): Production of sexed bovine pregnancies by cytogenetic analysis of cultured demi-embryos. *Theriogenology*, 27, 269.
- Robertson I, Nelson R (1998): Certification and identification of the embryo. 103–134. In: Strinkfellow D.A., Seidel S.M. (eds.): *Manual of the International Embryo Transfer Society*. IETS, Illinois. 171 pp.
- Sartori R, Souza AH, Guenther JN, Caraviello DZ, Geiger LN, Schenk JL (2004): Fertilization rate and embryo quality in superovulated Holstein heifers artificially inseminated with X-sorted or unsorted sperm. *Animal Reproduction*, 1, 86–90.
- Schenk JL, Suh TK, Seidel GE Jr. (2006): Embryo production from superovulated cattle following insemination of sexed sperm. *Theriogenology*, 65, 299–307.
- Schenk JL, Cran DG, Everett RW, Seidel GE Jr. (2009): Pregnancy rates in heifers and cows with cryopreserved sexed sperm: Effects of sperm numbers per inseminate, sorting pressure and sperm storage before sorting. *Theriogenology*, 71, 717–728.
- Schroder A, Roschlau D, Giehm D, Schwerin M, Thomson PD (1990): Sex determination of bovine embryos using Y-specific primers in the polymerase chain reaction. *Archiv fur Tierzucht*, 33, 293–299.
- Seidel GE Jr. (2007): Overview of sexing sperm. *Theriogenology*, 68, 443–446.
- Shea BF (1999): Determining the sex of bovine embryos using polymerase chain reaction results: a six-year retrospective study. *Theriogenology*, 51, 841–854.
- Singh EL, Hare WCD (1980): The feasibility of sexing bovine morula stage embryos prior to embryo transfer. *Theriogenology*, 14, 421–427.
- Thibier M, Nibart M (1995): The sexing of bovine embryos in the field. *Theriogenology*, 43, 71–80.
- Tominaga K (2004): Cryopreservation and sexing of in vivo- and in vitro-produced bovine embryos for their practical use. *Journal of Reproduction and Development*, 50, 29–38.
- Tominaga K, Hamada Y (2004): Efficient production of sex-identified and cryosurvived bovine in-vitro produced blastocysts. *Theriogenology*, 61, 1181–1191.
- Tominaga K, Iwaki F, Hochi S (2007): Conventional freezing of in vitro-produced and biopsied bovine blastocysts in the presence of a low concentration of glycerol and sucrose. *Journal of Reproduction and Development*, 53, 443–447.
- Voelkel SA, Hu YX (1992): Direct transfer of frozen-thawed bovine embryos. *Theriogenology*, 37, 23–37.
- White KL, Lindner GM, Anderson GB, BonDurant RH (1982): Survival after transfer of “sexed” mouse embryos exposed to H-Y antisera. *Theriogenology*, 18, 655–662.
- Williams TJ (1986): A technique for sexing mouse embryos by a visual colorimetric assay of the X-linked enzyme, glucose 6-phosphate dehydrogenase. *Theriogenology*, 25, 733–739.
- Yu W, Li S, Fang J, Sun X, Cui L, Fu J, Bai Y, Fang Y, Shanguan B (2007): Field studies on the effectiveness of the YCD embryo sexing technique in bovine. *Reproduction, Fertility and Development*, 19, 299–300.
- Zhang M, Lu KH, Seidel GE Jr. (2005): Blastocyst development of male and female bovine embryos produced by IVF with flow cytometrically sorted sperm. *Reproduction, Fertility and Development*, 17, 306–307.

Received: 2009–11–02

Accepted after corrections: 2010–01–29

Corresponding Author:

Prof. Assoc. MVDr. Miloslava Lopatarova, CSc., University of Veterinary and Pharmaceutical Sciences, Faculty of Veterinary Medicine, Clinic of Ruminant Diseases, Palackeho 1–3, 612 42 Brno, Czech Republic
Tel. +420 541 562 323, Fax +420 541 562 332, E-mail: lopatarovam@vfu.cz