

## Lactobacilli Isolated from Lump Sheep's Cheeses and their Antimicrobial Properties

JANA SMETANKOVÁ, ZUZANA HLADÍKOVÁ, MICHAELA ZIMANOVÁ,  
GABRIEL GREIF and MÁRIA GREIFOVÁ

Department of Food Science and Technology, Faculty of Chemical and Food Technology,  
Slovak University of Technology, Bratislava, Slovak Republic

### Abstract

SMETANKOVÁ J., HLADÍKOVÁ Z., ZIMANOVÁ M., GREIF G., GREIFOVÁ M. (2014): **Lactobacilli isolated from lump sheep's cheeses and their antimicrobial properties.** Czech J. Food Sci., 32: 152–157.

A total of 34 strains of lactobacilli were isolated from the lump sheep's cheeses produced from raw sheep milk. The strains were identified by MALDI-TOF MS, and 20 of them demonstrating the best fermentation and sensoric properties in milk were chosen and tested for their antimicrobial activity. All selected strains were active against the indicator bacteria and moulds. The highest inhibitory effect was observed with the strains *Lactobacillus paracasei* 314, *L. paracasei* 316, *L. plantarum* K816, *L. plantarum* L718, and *L. plantarum* 2L2. The subsequent research was focused on the metabolites causing this inhibition. The production of lactic and acetic acids was studied under different cultivation conditions (0, 2, 4, and 6.5% NaCl addition; cultivation at 15, 30, 37, and 45°C; and pH value of the broth before sterilisation 5 and 9). *L. plantarum* L718 produced the highest concentration of lactic and acetic acids under most of the cultivation conditions. Antimicrobial substances such as phenyllactic acid (62.54–101.62 mg/dm<sup>3</sup>), H<sub>2</sub>O<sub>2</sub> (0.78–2.30 µg/cm<sup>3</sup>), and diacetyl (produced by *L. plantarum* K816 and L718) were studied as well.

**Keywords:** *Lactobacillus plantarum*; *Lactobacillus paracasei*; antimicrobial potential

It is known that the natural microflora, present in the traditional biotechnological processes during manufacturing, participates in the production of the characteristic aroma, taste, and texture of different cheeses. At first, only artisanal starters were used for the dairy although fermentations. Although they are still used in certain cases, their microbiological instability promoted the evolution of more precisely defined mixtures of lactic acid bacteria (LAB). As cheesemaking became more industrialised, pasteurised milk and standardised bacterial and fungal inocula were introduced with the aim of obtaining a more stable acidifying activity and consistent quality of the products (CARMINATI *et al.* 2010). In spite of this trend, the production of raw milk cheeses is still significant, as consumers seek organic foods or traditional sensory characteristics. However, the diversity of microbial populations that have devel-

oped in the original cheesemaking environments is largely undefined and is a valuable source of new strains for commercial uses (LAURENČÍK *et al.* 2008). For example, non-starter LABs with the potential to inhibit the undesirable microflora are used in the biopreservation of foods (RODRÍGUEZ *et al.* 2000).

As far as the technology is concerned, several characteristic properties of the bacterial strains of LAB (acid production in different media and at different temperatures, proteinase and peptidase activities, autolysis, production of volatile compounds, resistance to bacteriophages, production of inhibitory compounds) are important for their use as starters or adjuncts and have been evaluated in the screening and characterisation studies (PIRAINO *et al.* 2008).

The main task of this work was to observe the antimicrobial activity of 20 newly isolated strains of lactobacilli and to study the technological potential

Supported by the Slovak Research and Development Agency, Grant APVV No. 07/0158, and by the Scientific Grant Agency of the Slovak Republic, Grant No. VEGA 1/0879/12.

of 5 selected strains (2 strains of *L. paracasei* and 3 strains of *L. plantarum*), which showed very good antimicrobial activities.

## MATERIAL AND METHODS

**Microorganisms studied.** 34 strains of lactobacilli were isolated from the lump sheep's cheeses produced from raw sheep milk in different farms of the Slovak Republic. The isolates were identified by MALDI-TOF MS. 20 strains of lactobacilli (10 *L. paracasei* and 10 *L. plantarum*) were chosen for the subsequent testing in view of the fermentative and sensoric properties in milk.

**Identification.** For MALDI-TOF (matrix-assisted laser desorption/ionisation time-of-flight) bacterial analysis, the cells from a single colony of fresh overnight culture were used for each isolate to prepare samples according to the microorganism profiling ethanol-formic acid extraction procedure, as recommended by the manufacturer. Each sample spot was overlaid with 2 µl of the matrix solution (saturated solution of  $\alpha$ -cyano-4-hydroxy-cinnamic acid in 50% acetonitrile with 2.5% trifluoroacetic acid) and air-dried again for 15 min (Bruker Daltonik GmbH, Bremen, Germany). To identify the microorganisms, the raw spectra obtained with each isolate were imported into BioTyper software, Version 2.0 (Bruker Daltonik GmbH, Bremen, Germany), and analysed without any user intervention (KMEŤ & DRUGDOVÁ 2012).

The antimicrobial activity was studied by the dualculture overlay diffusion method (MAGNUS-SON 2003). Lactobacilli were inoculated (0.1 ml of overnight culture) into sterile disks (Rotilabo<sup>®</sup>-test flakes,  $\varnothing$  12 mm; Carl Roth GmbH+Co. KG, Karlsruhe, Germany) on MRS agar plates (Merk, Darmstadt, Germany) and allowed to grow at 37°C for 48 h in aerobic conditions. The plates were then overlaid with soft BHI agar (broth + 0.8% agar; Merk, Darmstadt, Germany) containing the inoculum of the indicator bacteria (inoculum 1%) or with soft Sabouraud agar (broth + 0.8% agar; Imuna, Šarišské Michalany, Slovakia) containing the inoculum of the indicator mould (10<sup>4</sup> per ml mould spores). After 24 h (bacteria) and 48 h (moulds) of aerobic incubation at 37°C (bacteria) and 25°C (moulds), the respective zone of inhibition was measured and % of inhibition of the total area of the Petri dish (PD) was calculated. The inhibition tests were done in duplicates. The antimicrobial activity of lactobacilli was observed against 6 indicator bacteria (*Bacillus cereus* OPT, *Bacillus subtilis* CCM 2216, *Escherichia coli* CCM 3988, *Listeria*

*monocytogenes* NCTC 4886, *Pseudomonas aeruginosa* CCM 3955 and *Staphylococcus aureus* CCM 3953) and 5 indicator moulds (*Alternaria alternata* CCM F-128, *Aspergillus flavus* CCM F-171, *Mucor rouxii* CCM F-220, *Penicillium chrysogenum* CCM F-432 and *Rhizopus oryzae* CCM F-8284) (CCM – Czech Collection of Microorganisms, Czech Republic; OPT – Collection of Department of Food Science and Technology, Slovakia; NCTC – National Collection of Type Cultures, UK).

Subsequently, the lactobacilli with the best antimicrobial activity were investigated for their production of antimicrobial substances.

**Antimicrobial substances.** *Organic acids production.* The ability of lactobacilli to produce lactic and acetic acids was studied in MRS broth (Merk, Darmstadt, Germany) with different additions of NaCl (no addition, 2, 4, and 6.5% addition) at 37°C; in MRS broth at different temperatures (15, 30, 37, and 45°C) and in MRS broth with pH value adjusted before sterilisation (pH 5 and 9) at 37°C. The inoculum was 1%. The samples were analysed using HPLC after 50 h of fermentation.

*Phenyllactic acid production* was observed in MRS broth with the addition of 0.1% of L-phenylalanine (Merk, Darmstadt, Germany). Lactobacilli were incubated at 37°C for 72 h in anaerobic conditions (inoculum 1%). The concentrations of phenyllactic acid were determined with HPLC (GREIF *et al.* 2008).

*Hydrogen peroxide production.* Lactobacilli were grown in MRS broth at 37°C for 16 h, harvested by centrifugation (4000 g, 5°C, 20 min), washed twice in cold physiological saline, and resuspended in sodium phosphate buffer (pH 6.5). Following the incubation (24 h at 5°C), the cells were removed by centrifugation and the supernatant was assayed for H<sub>2</sub>O<sub>2</sub>. 2.5 ml of the cell free supernatant was placed into tubes containing 0.5 ml of 0.1% aqueous solution of peroxidase (Boehringer, Mannheim, Germany) and 0.05 ml of 1% aqueous solution of *o*-dianisidine (SERVA, Heidelberg, Germany). The blank was prepared using 2.5 ml of sodium phosphate buffer instead of the sample supernatant. The tubes were incubated at 37°C, for 10 minutes. The reaction was stopped by adding 0.1 ml of 4N HCl. The absorbance ( $\lambda = 400$  nm) of each sample was read and peroxide content was determined by comparison with the standard curve (VILLEGAS & GILLIALAND 1998).

*Diacetyl production.* The revitalised strains were inoculated in 9 ml of UHT milk and allowed to grow at 30°C for 24 h in aerobic conditions (inoculum 1%). 1 ml of each cell suspension was combined with

0.5 ml of 1% ethanolic solution of  $\alpha$ -naphthol (Merck, Darmstadt, Germany) and 0.5 ml of 16% aqueous solution of KOH (Mikrochem, Pezinok, Slovakia), and incubated at 30°C for 10 minutes. Diacetyl production was indicated by the formation of a red ring at the top of the test tubes (DAL BELLO *et al.* 2012).

## RESULTS AND DISCUSSION

Antimicrobial activity of 20 new-isolated strains of lactobacilli (10 *L. paracasei* and 10 *L. plantarum*) was observed in this study (Table 1 and Figure 1). All monitored strains showed considerable inhibition of the indicator bacteria *B. cereus*, *B. subtilis*, *E. coli*, *L. monocytogenes*, *P. aeruginosa*, and *S. aureus* (the average more than 20% of the total area of the PD).

Antifungal activity of *Lactobacillus* strains varied. The inhibition zones were in the interval of 4.1–14.3% of the total area of the PD. The activities of the strains *L. plantarum* and *L. paracasei* were

comparable. The most inhibited were *P. chrysogenum* and *R. oryzae* (an average of 10.4% and 9.1% of the total area of PD). *A. alternata*, *A. flavus*, and *M. rouxii* were inhibited at an average of 7.2, 7.2, and 8.4% of the total area of PD.

Two strains of *L. paracasei* (marked 314 and 316) and 3 strains of *L. plantarum* (marked K816, 2L2, and L718) were chosen for the subsequent testing due to their best antimicrobial activity of all the lactobacilli studied. Their average antibacterial activity was higher than 25% and their average antifungal activity was higher than 8.5% of the total area of PD.

The ability of *L. paracasei* and *L. plantarum* to possess antimicrobial activity had been previously published as well. The antibacterial activities of 17 strains of *L. plantarum*, isolated from a Tunisian traditional salted meat, were studied by ESSID *et al.* (2009). All strains displayed inhibition zones against *S. aureus*, 94% against *E. coli*, and 88% against *P. aeruginosa*. REN *et al.* (2012) observed the effects

Table 1. Antimicrobial activity (% of inhibition of the total area of the PD)

| Isolated strains    | Indicator bacteria |                    |                |                         |                      |                  | Indicator moulds    |                  |                  |                       |                  |      |
|---------------------|--------------------|--------------------|----------------|-------------------------|----------------------|------------------|---------------------|------------------|------------------|-----------------------|------------------|------|
|                     | <i>B. cereus</i>   | <i>B. subtilis</i> | <i>E. coli</i> | <i>L. monocytogenes</i> | <i>P. aeruginosa</i> | <i>S. aureus</i> | <i>A. alternata</i> | <i>A. flavus</i> | <i>M. rouxii</i> | <i>P. chrysogenum</i> | <i>R. oryzae</i> |      |
| <i>L. paracasei</i> | 113                | 26.5               | 20.2           | 11.9                    | 16.5                 | 34.2             | 24.8                | 8.8              | 6.7              | 2.5                   | 13.3             | 9.3  |
|                     | 119                | 18.3               | 19.2           | 26.5                    | 19.4                 | 27.7             | 38.8                | 7.6              | 5.0              | 6.3                   | 7.5              | 9.8  |
|                     | 211                | 21.9               | 28.4           | 20.2                    | 22.5                 | 19.9             | 20.8                | 6.3              | 11.1             | 6.4                   | 12.1             | 7.1  |
|                     | 314                | 19.5               | 30.1           | 27.6                    | 22.7                 | 29.3             | 34.6                | 8.0              | 9.0              | 6.3                   | 12.7             | 13.7 |
|                     | 316                | 20.8               | 34.0           | 18.3                    | 23.6                 | 32.4             | 32.4                | 7.2              | 7.2              | 14.3                  | 12.9             | 9.2  |
|                     | 11L5               | 13.8               | 33.2           | 14.1                    | 21.9                 | 26.1             | 30.7                | 5.9              | 6.8              | 8.4                   | 12.2             | 11.4 |
|                     | 11L7               | 30.3               | 30.7           | 12.0                    | 13.0                 | 28.8             | 21.2                | 8.8              | 7.6              | 4.9                   | 7.4              | 6.0  |
|                     | 12L1               | 23.1               | 34.5           | 16.5                    | 29.3                 | 35.3             | 21.5                | 6.6              | 7.0              | 5.7                   | 12.3             | 4.1  |
|                     | 12L3               | 15.9               | 30.7           | 10.4                    | 26.5                 | 25.0             | 27.2                | 6.3              | 8.0              | 13.9                  | 12.2             | 6.8  |
|                     | 12L6               | 19.2               | 31.1           | 14.2                    | 29.7                 | 25.0             | 36.2                | 6.6              | 6.8              | 7.0                   | 6.6              | 6.9  |
| <i>L. plantarum</i> | 115                | 16.5               | 24.6           | 25.9                    | 29.1                 | 20.7             | 26.8                | 7.2              | 5.3              | 11.5                  | 12.2             | 7.7  |
|                     | 219                | 19.2               | 22.5           | 13.6                    | 21.0                 | 37.1             | 38.4                | 7.8              | 7.6              | 5.8                   | 7.6              | 7.6  |
|                     | L 717              | 17.7               | 28.6           | 23.6                    | 19.5                 | 12.3             | 22.9                | 8.4              | 10.9             | 5.1                   | 11.4             | 13.5 |
|                     | L 718              | 28.2               | 28.2           | 32.4                    | 29.1                 | 28.8             | 24.8                | 7.8              | 1.3              | 14.3                  | 9.3              | 9.8  |
|                     | 812                | 41.6               | 25.7           | 22.2                    | 25.2                 | 10.3             | 30.7                | 5.0              | 7.4              | 5.0                   | 9.9              | 14.1 |
|                     | K 816              | 16.2               | 19.5           | 33.2                    | 19.5                 | 35.3             | 29.1                | 6.0              | 6.5              | 7.3                   | 14.1             | 13.8 |
|                     | K 817              | 23.9               | 19.9           | 13.3                    | 19.2                 | 30.3             | 37.9                | 12.6             | 7.7              | 7.2                   | 7.5              | 6.8  |
|                     | 1L5                | 19.4               | 28.0           | 11.1                    | 23.9                 | 29.5             | 44.3                | 5.4              | 5.0              | 6.0                   | 8.0              | 7.1  |
|                     | 2L2                | 23.8               | 45.4           | 18.3                    | 17.1                 | 32.0             | 26.1                | 5.4              | 7.7              | 13.3                  | 12.2             | 8.7  |
|                     | 1L10               | 24.3               | 37.1           | 20.5                    | 17.7                 | 23.9             | 19.7                | 6.4              | 9.3              | 16.7                  | 7.4              | 9.2  |




Figure 1. Inhibitory activity of (a) *L. paracasei* 314 against *E. coli* and (b) *L. plantarum* L718 against *M. rouxii*

of the selected strains of lactobacilli (*L. salivarius* subsp. *salicinius* CICC 23174, *L. plantarum* CGMCC 1.557, *L. rhamnosus* ATCC 53103 and *L. acidophilus* ATCC 4356) on the undesirable *S. aureus*. *L. paracasei* susp. *paracasei* that was isolated from breast-fed newborns' faeces, showed anti-listerial activity due to the ability to produce a bacteriocin-like substance (BENDALI *et al.* 2008).

Antifungal activity of *L. paracasei* (isolated from Edam cheese) was observed against *Fusarium proliferatum* M 5689, followed by *Penicillium* sp. DMF 0006 and *Aspergillus niger* DMF 0801 (TŮMA *et al.* 2007). *L. plantarum* 01, *L. paracasei* 05, *L. paracasei* SF1 and *L. plantarum* 2142 (strains from collections) inhibited the growth of *A. flavus* DMF 0802 (HUDÁČEK *et al.* 2007).

LABs have traditionally been used as natural bio-preservatives in food and animal feed, sauerkraut, and silage. Their preserving effect relates mainly to the formation of organic acids and H<sub>2</sub>O<sub>2</sub>, competition for nutrients and production of antimicrobial substances

(VOULGARI *et al.* 2010). The acidic condition and the reduced pH of the fermented dairy products, as well as the antimicrobial activity of the undissociated lactic acid molecules, prevent the growth or survival of many spoilage and pathogenic bacteria and moulds (CARMINATI *et al.* 2010; LEÓN PELÁEZ *et al.* 2012).

The produced concentrations of lactic and acetic acids are given in Table 2. Generally, the growth and organic acids production were supported by the addition of 2% and 4% of NaCl. The temperatures of 30 and 37°C and pH value 5 had positive effects on the lactic acid production. Acetic acid was produced in higher quantities at the temperatures of 37 and 45°C and pH value of 9. Out of all the strains analysed, *L. plantarum* L718 produced the highest concentration of lactic and acetic acids in most of the cultivation conditions.

GEREZ *et al.* (2013) observed very good antifungal activities in 4 strains of *L. plantarum*, isolated from different food sources, and analysed the antifungal substances. After 24 h of cultivation in MRS broth, lactobacilli produced 19.52–24.13 g/dm<sup>3</sup> of lactic acid; 0.77–1.20 g/dm<sup>3</sup> of acetic acid, and 33.23–581.60 mg/dm<sup>3</sup> of phenyllactic acid. The antifungal activity, however, is not related only to lactic acid, phenyllactic acid, or any other single inhibitory substance, but rather to the combination of such substances (WANG *et al.* 2013).

The bactericidal effect of H<sub>2</sub>O<sub>2</sub> has been attributed to its strong oxidising effect on the bacterial cells and to the destruction of basic molecular structures of the cell proteins. ZALÁN *et al.* (2005) showed that *L. plantarum* 2142 produced enough H<sub>2</sub>O<sub>2</sub> (2.45 µg/ml)

Table 2. Concentration of lactic acid and acetic acid (g/dm<sup>3</sup>) after 50 h cultivation in different growth conditions

| Isolated strains   | NaCl addition (%) |        |        |        | Temperature of cultivation (°C) |        |        |       | pH of broth |        |
|--------------------|-------------------|--------|--------|--------|---------------------------------|--------|--------|-------|-------------|--------|
|                    | 0                 | 2      | 4      | 6.5    | 15                              | 30     | 37     | 45    | 5           | 9      |
| <b>Lactic acid</b> |                   |        |        |        |                                 |        |        |       |             |        |
| 314                | 9.061             | 9.350  | 11.932 | 1.593  | 0.247                           | 9.419  | 9.102  | 8.107 | 11.336      | 8.931  |
| 316                | 15.264            | 16.880 | 11.837 | 1.185  | 1.694                           | 14.279 | 14.469 | 0.427 | 21.371      | 8.138  |
| K816               | 9.745             | 10.650 | 16.039 | 4.146  | 1.796                           | 8.210  | 8.835  | 7.156 | 11.207      | 10.158 |
| L718               | 17.373            | 17.347 | 16.474 | 11.289 | 8.286                           | 16.087 | 16.505 | 4.990 | 18.259      | 2.261  |
| 2L2                | 8.147             | 9.730  | 14.329 | 7.707  | 0.859                           | 9.790  | 9.813  | 9.390 | 13.593      | 8.395  |
| <b>Acetic acid</b> |                   |        |        |        |                                 |        |        |       |             |        |
| 314                | 0.805             | 0.697  | 0.572  | 0.250  | nd                              | 0.978  | 0.881  | 0.846 | 1.084       | 1.558  |
| 316                | nd                | 0.138  | 0.208  | 0.088  | 0.091                           | 0.089  | 0.137  | 0.125 | nd          | 0.181  |
| K816               | 1.378             | 1.546  | 1.065  | 0.214  | nd                              | 0.360  | 0.674  | 0.854 | 0.923       | 1.319  |
| L718               | 0.897             | 0.863  | 1.275  | 0.590  | 0.745                           | 1.453  | 1.464  | 0.824 | 0.551       | ND     |
| 2L2                | 0.761             | 1.341  | 1.002  | 0.373  | nd                              | 0.515  | 0.900  | 1.070 | 1.314       | 1.480  |

nd – not detected

Table 3. Production of other important antimicrobial substances

| Isolated strains | Antimicrobial substances                |   |                       |
|------------------|---|---|-----------------------|
|                  | phenyllactic acid (mg/dm <sup>3</sup> ) | H <sub>2</sub> O <sub>2</sub> (µg/cm <sup>3</sup> ) | diacetyl <sup>1</sup> |
| 314              | 85.13                                   | 1.88  | –                     |
| 316              | 62.54                                   | 0.78  | –                     |
| K816             | 101.62                                  | 2.30  | +                     |
| L718             | 65.70                                   | 1.73  | +                     |
| 2L2              | 94.98                                   | 2.27  | –                     |

<sup>1</sup>classification scale: – no production; + production

after 24 h at 5°C in phosphate buffer to inhibit the growth of *L. monocytogenes* and *B. cereus*.

In this study, antimicrobial substances such as phenyllactic acid (62.54–101.62 mg/dm<sup>3</sup>), H<sub>2</sub>O<sub>2</sub> (0.78–2.30 µg/cm<sup>3</sup>), and diacetyl were also studied (Table 3). Diacetyl was produced only by the strains *L. plantarum* K816 and L718, but NIKOLIC *et al.* (2008) demonstrated its production also by the strains of *L. paracasei* subsp. *paracasei* (isolate from Bukuljac cheese). Antimicrobial activity of diacetyl is known; for example, the activity against *E. coli*, *L. monocytogenes*, and *S. aureus* was proved by LANCIOTTI *et al.* (2003).

## CONCLUSIONS

Five lactobacilli strains were selected due to their marked antimicrobial activities, and their antimicrobial metabolites were analysed. The inhibitory activity of lactobacilli is dependent on the producer, cultivated medium, and also on the strains inhibited, especially in the case of moulds. The determined concentrations of lactic and acetic acids confirmed the ability to inhibit the undesirable microflora. Phenyllactic acid, H<sub>2</sub>O<sub>2</sub> and diacetyl were also involved in the synergic inhibitory effect. Therefore these lactobacilli could be used as adjunct cultures. However, in the first place other studies on their safe application in foods are needed.

**Acknowledgements.** We thank Prof DVM VLADIMÍR KMEŇ, DSc., for the identification of the strains tested.

## References

- BENDALI F., GAILLARD-MARTINIE B., HEBRAUD M., SADOUN D. (2008): Kinetic of production and mode of action of the *Lactobacillus paracasei* subsp. *paracasei* anti-listerial bacteriocin, an Algerian isolate. *LWT-Food Science and Technology*, **41**: 1784–1792.
- CARMINATI D., GIRAFFA G., QUIBERONI A., BINETTI A., SUÁREZ V., REINHEIMER J. (2010): Advances and trends in starter cultures for dairy fermentations. In: MOZZI F., RAYA R.R., VIGNOLO G.M. (eds): *Biotechnology of Lactic Acid Bacteria: Novel Applications*. Wiley-Blackwell, Ames-Singapore: 177–285.
- DAL BELLO B., COCOLIN L., ZEPPA G., FIELD D., COTTER P.D., HILL C. (2012): Technological characterization of bacteriocin producing *Lactococcus lactis* strains employed to control *Listeria monocytogenes* in Cottage cheese. *International Journal of Food Microbiology*, **153**: 58–65.
- ESSID I., MEDINI M., HASSOUNA M. (2009): Technological and safety properties of *Lactobacillus plantarum* strains isolated from a Tunisian traditional salted meat. *Meat Science*, **81**: 203–208.
- GEREZ C.L., TORRES M.J., FONT DE VALDEZ G., ROLLÁN G. (2013): Control of spoilage fungi by lactic acid bacteria. *Biological Control*, **64**: 231–237.
- GREIF G., GREIFOVÁ M., KAROVIČOVÁ J., KOHAJDOVÁ Z., TOMAŠKA M. (2008): Využitie IEX-HPLC na stanovenie metabolitov produkovaných baktériami mliečnej fermentácie. In: 17<sup>th</sup> International Conference “Analytical and Human Health”. Nový Smokovec, October 20–23. (CD ROM)
- HUDÁČEK J., ZSOLT Z., CHUMCHALOVÁ J., HALÁSZ A. (2007): Antifungálny účinok laktobacilov na plesne rodu *Fusarium* a *Aspergillus*. *Chemické listy*, **101**: 730–737.
- KMEŇ V., DRUGDOVÁ Z. (2012): Antimicrobial susceptibility of microflora from ovine cheese. *Folia Microbiologica*, **57**: 291–293.
- LANCIOTTI R., RATRIGNANI F., BAGNOLINI F., GUERZONI M.E., GARDINI F. (2003): Evaluation of diacetyl antimicrobial activity against *Escherichia coli*, *Listeria monocytogenes* and *Staphylococcus aureus*. *Food Microbiology*, **20**: 537–543.
- LAURENČÍK M., SULO P., SLÁVIKOVÁ E., PIECKOVÁ E., SEMAN M., EBRINGER L. (2008): The diversity of eukaryotic microbiota in the traditional Slovak sheep cheese – Bryndza. *International Journal of Food Microbiology*, **127**: 176–179.
- LEÓN PELÁEZ A.M., SERNA CATAÑO C.A., QUINTERO YEPES E.A., GAMBA VILLARROEL R.R., DE ANTONI G.L., GIANNUZZI L. (2012): Inhibitory activity of lactic and acetic acid on *Aspergillus flavus* growth for food preservation. *Food Control*, **24**: 177–183.
- MAGNUSSON J., STRÖM K., ROOS S., SJÖGREN J., SCHNÜRER J. (2003): Broad and complex antifungal activity among environmental isolates of lactic acid bacteria. *FEMS Microbiology Letters*, **219**: 129–135.
- NIKOLIC M., TERZIC-VIDOJEVIC A., JOVICIC B., BEGOVIC J., GOLIC N., TOPISIROVIC L. (2008): Characterization of lactic acid bacteria isolated from Bukuljac, a homemade goat's milk cheese. *International Journal of Food Microbiology*, **122**: 162–170.
- PIRAINO P., ZOTTA T., RICCIARDI A., MCSWEENEY P.L.H., PARENTE E. (2008): Acid production, proteolysis, autolytic and inhibitory properties of lactic acid bacteria isolated from pasta filata cheeses: A multivariate screening study. *International Dairy Journal*, **18**: 81–92.

- REN D., LI CH., QIN Y., YIN R., LI X., TIAN M., DU S., GUO H., LIU C., ZHU N., SUN D., LI Y., JIN N. (2012): Inhibition of *Staphylococcus aureus* adherence to Caco-2 cells by lactobacilli and cell surface properties that influence attachment. *Anaerobe*, **18**: 508–515.
- RODRÍGUEZ E., GONZÁLEZ B., GAYA P., NUÑEZ M., MEDINA M. (2000): Diversity of bacteriocins produced by lactic acid bacteria isolated from raw milk. *International Dairy Journal*, **10**: 7–15.
- TŮMA Š., VOGENSEN F.K., PLOCKOVÁ M., CHUMCHALOVÁ J. (2007): Isolation of antifungally active lactobacilli from Edam cheese. *Acta Alimentaria*, **36**: 405–414.
- VILLEGAS E., GILLILAND S.E. (1998): Hydrogen peroxide production by *Lactobacillus delbrueckii* subsp. *lactis* I at 5°C. *Journal of Food Science*, **63**: 1070–1704.
- VOULGARI K., HATZIKAMARI M., DELEPOGLOU A., GEORGAKOPOULOS P., LITOPOULOU-TZANETAKI E., TZANETAKIS N. (2010): Antifungal activity of non-starter lactic acid bacteria isolates from dairy products. *Food Control*, **21**: 136–142.
- WANG H.K., SUN Y., CHEN CH., SUN Z., ZHOU Y.CH., SHEN F.D., ZHANG H.P., DAI Y.J. (2013): Genome shuffling of *Lactobacillus plantarum* for improving antifungal. *Food Control*, **32**: 341–347.
- ZALÁN Z., NÉMETH E., BARÁTH Á., HALÁSZ A. (2005): Influence of growth medium on hydrogen peroxide and bacteriocin production of *Lactobacillus* strains. *Food Technology and Biotechnology*, **43**: 219–225.

Received for publication April 3, 2013  
Accepted after corrections July 17, 2013

---

*Corresponding author:*

Ing. JANA SMETANKOVÁ, Slovenská technická univerzita v Bratislavě, Fakulta chemickej a potravinárskej technológie, Ústav biotechnológie a potravinárstva, Radlinského 9, 812 37 Bratislava, Slovenská republika; E-mail: jana.smetankova@gmail.com

---