

Rhizobium japonicum as a Biocontrol Agent of Soybean Root Rot Disease Caused by *Fusarium solani* and *Macrophomina phaseolina*

RAKIB A. AL-ANI¹, MUSTAFA A. ADHAB¹, MAJDA H. MAHDI² and HADI M. ABOOD³

¹Plant Protection Department, College of Agriculture and ²Department of Biology, College of Science, University of Baghdad, Baghdad, Iraq; ³Ministry of Sciences and Technology, Baghdad, Iraq

Abstract

AL-ANI R.A., ADHAB M.A., MAHDI M.H., ABOOD H.M. (2012): *Rhizobium japonicum* as a biocontrol agent of soybean root rot disease caused by *Fusarium solani* and *Macrophomina phaseolina*. Plant Protect. Sci., 48: 149–155.

The activity of *Rhizobium japonicum* against the soil-borne pathogens *Fusarium solani* and *Macrophomina phaseolina* as causative agents of soybean root rot disease in both culture medium and soil was evaluated. Rhizobial culture filtrate caused an inhibition of the fungal radial growth of *Fusarium solani* and *Macrophomina phaseolina* on potato dextrose agar medium amended with the filtrate compared with control. The addition of rhizobial culture suspension to the soil contaminated by the two pathogens, *Fusarium solani* and *Macrophomina phaseolina* and their interaction, in pots, improved seed germination percentages and reduced the root rot disease index significantly. The sowing of rhizobial coated seeds in soil contaminated by *Fusarium solani* and *Macrophomina phaseolina* separately and in combination, in the field, increased seed germination significantly and induced a high reduction in disease severity for the same previous combination under field conditions. These results indicate that rhizobia could be an important element in root rot disease management.

Keywords: biological control, *Rhizobium* sp.; soybean; *Fusarium*; *Macrophomina*

Soybean (*Glycine max* L. Merrill) is one of the most important legume crops in the world representing a major component of the diet of food-producing animals and humans (HAPGOOD 1987; FRIEDMAN & BRANDON 2001). Soybean oil was found to induce an increase in bird's weight when added to broiler diet (SCAIFE *et al.* 1994; VIEIRA *et al.* 2002; LARA *et al.* 2003).

It has been reported that soybean plants are subject to the infection by several soil-borne pathogens, inducing root rot disease, which is considered among the most important limiting factors affecting plant growth and yield (COOK *et al.* 1995; YANG & FENG 2001); *Fusarium solani* and *Macrophomina phaseolina* are among the most important of them (ZEMANKOVÁ & LEBEDA 2001; VEVERKA *et al.* 2008). Due to great harms caused by chemicals used to

control soil-borne pathogens, to environment and health, biological control using non-pathogenic microorganisms was adopted as an alternative to chemicals for combating these pathogens.

Saprophytic rhizosphere bacteria are present in large numbers on the plant root surface using root exudate and lysate as nutrients for growth. Certain of these bacteria can stimulate plant growth through releasing secondary metabolites which facilitate the uptake of certain nutrients from the root environment, so they are referred to as plant growth promoting rhizobacteria (PGPR) as well as suppressing soil-borne pathogens in the rhizosphere (BAKKER *et al.* 2007).

The colonization of plant roots by plant growth promoting rhizobacteria and other factors and treatments can elicit the host defence against

various pathogens as indicated by a reduction of disease incidence and severity which constitute a state of induced systemic resistance in the plants to subsequent pathogen attack (HAMMERSCHMIDT 1999; KLOEPFER *et al.* 2004; VAN LOON 2004; AL-ANI *et al.* 2010, 2011a,b,c; DIWAN *et al.* 2011). PGRR isolated from the rhizosphere soil have been shown to inhibit plant pathogens through competition for nutrients, competition for iron by siderophores, antibiosis or secretion of lytic enzymes as well as inducing systemic resistance (ISR) (JOGAIAH 2010). Of these PGRR, rhizobia are reported as effective biocontrol agent for the inhibition of soil-borne plant pathogens. Many species of rhizobia were found to promote plant growth and also to inhibit the growth of various soil-borne pathogens including *Macrophomina phaseolina*, *Rhizoctonia solani*, *Fusarium* sp. in both leguminous and non-leguminous plants (EHTESHAMUL-HAQUE & GHAFFAR 1993; SHARIF *et al.* 2003; EL-BATANONY *et al.* 2007).

The present study was conducted to evaluate the antagonistic activity of *Rhizobium japonicum* against *Fusarium solani* and *Macrophomina phaseolina* causative agents of root rot disease in soybean.

MATERIAL AND METHODS

Fungal isolates. Small parts of soybean roots from plants showing yellowing and wilting symptoms were surface sterilised in 5% sodium hypochlorite for 3 minutes. The sterilised fragments were washed several times with sterile distilled water and dried on filter papers. The fragments were assigned to two groups. Fragments of the first group were placed on potato sucrose agar (PSA), while fragments of the second group were placed on Modified Nash and Synders Medium (MNSM) (CHO *et al.* 2001) in Petri dishes of 9 cm diameter. The plates were incubated at $25 \pm 2^\circ\text{C}$ for 5 days. *Macrophomina phaseolina* isolates were identified according to HOLLIDAY and PUNITHALINGAT (1970), and *Fusarium* isolates according to BOOTH (1981) and stored at 4°C .

Rhizobial isolates. Isolates of *Rhizobium japonicum*, maintained on peat moss, were obtained from Rhizobial Research Laboratory, General State of Applied Research, Ministry of Agriculture, Iraq. The isolates were grown on yeast mannitol broth (YMB) containing 0.1% yeast extract, 1% mannitol, 0.05% potassium phosphate, 0.02% magnesium sul-

phate, 0.02% NaCl, pH 7.6, in 250 ml Erlenmeyer flasks, in shaking incubator at 28°C for 28 hours. The number of cells/ml in the rhizobial culture was determined by colony count method (BLACK 1965) on yeast mannitol agar (YMA) and used at 10^8 cells/ml in laboratory and greenhouse experiments.

Rhizobial culture filtrate. Rhizobial filtrate was obtained by centrifugation of rhizobial culture at 6000 rpm for 20 min and the supernatant were passed through $0.45 \mu\text{m}$ bacterial filter.

Activity of rhizobial culture filtrate against *F. solani* and *M. phaseolina* on culture media. The test was carried out using an agar well diffusion method on PDA culture medium. Four wells of 0.5 cm diameters were equidistantly made on PDA in Petri dishes of 9 cm diameter. One ml of rhizobial filtrate at 0, 25, 50, and 100% was placed separately in each well. A disc of 0.5 cm diameter of fungal growth, from fresh culture on PDA medium, was placed at the centre of the PDA plate. The plates were incubated at $28 \pm 2^\circ\text{C}$ for 5 days and the radial growth of fungi was assessed in triplicate Petri dishes. Percentage growth inhibition was calculated according to the following formula: % growth inhibition = (control – treatment/control) \times 100.

The efficiency of rhizobial filtrate on the radial growth of the pathogenic fungi was also determined by mixing the filtrate with PDA before solidification at the same above concentrations in Petri dishes. The medium was inoculated by the fungi and the percentage of growth reduction was calculated as before.

Rhizobium antagonism against *F. solani* and *M. phaseolina* in pots. Sterile clay-loamy to peat moss (3:1) soil in pots (2 kg/pot) was contaminated separately by *F. solani*, *M. phaseolina*, and both of them grown on bran and groats at 2 g/kg soil. Each pot was drenched with 10 ml of *R. japonicum* suspension at 1×10^8 cell/ml after 2 days of fungal contamination. Pots contaminated by pathogens only and non-treated ones (without fungal and bacterial inoculant) were considered as control. The pots were arranged in a greenhouse ($28 \pm 2^\circ\text{C}$) in a randomised block design in three replicates, cultivated with surface sterilised soybean seeds (10 seed/pot) and watered as needed. The treatments were *F. solani*, *M. phaseolina*, *F. solani* + *M. phaseolina*, *F. solani* + rhizobia, rhizobia only, and control.

The percentage of germination was determined one week after sowing. Woltz and Arther index (WOLTZ & ARTHUR 1973) was used to determine disease severity. The disease severity was calculated according to the following equation:

$$\text{Disease severity (\%)} = \frac{\sum(\text{disease rate} \times \text{number of plant with this rate})}{\text{total number of plants} \times \text{maximum value of disease scale}} \times 100$$

Field trial. Field experiment was conducted in the 2010 growing season on 4/5/2010 at state of Applied Research Station, Baghdad, Abu-Ghraib, in a randomised block design with plots of $3 \times 3 \text{ m}^2$ and 3 replicates. Seeds of soybean were dipped in Rhizobium culture suspension at 10^8 cells/ml, containing 2% Arabic gum as sticker, for one hour. Non-treated seed were dipped in distilled water as control. Rhizobium-treated and non-treated seeds were sown in rows 3 m long and 30 cm in spacing.

The fungal inoculum, grown on a mixture of bran and groats, was distributed along the rows at 6 g/m long, 24 h before seed sowing. The treatments were as follows:

- (1) Rhizobium treated seeds in *F. solani* contaminated soil;
- (2) Rhizobium treated seeds in *M. phaseolina* contaminated soil;
- (3) Rhizobium treated seeds in *F. solani* and *M. phaseolina* contaminated soil;
- (4) Non-treated seeds in *F. solani* contaminated soil;
- (5) Non-treated seeds in *M. phaseolina* contaminated soil;
- (6) Non-treated seeds in *F. solani* + *M. phaseolina* contaminated soil;
- (7) Non-treated seeds in non-contaminated soil;
- (8) Rhizobium treated seeds in non-contaminated soil.

Germination percentage and root rot index were determined as previously after 19 weeks.

RESULTS

Activity of rhizobial culture filtrate against *Fusarium solani* and *Macrophomina phaseolina* on culture medium

Soybean rhizobial culture filtrate showed a significant reduction in the radial growth of *F. solani* and *M. phaseolina* on PDA culture medium. The inhibitory effect of the filtrate on the two pathogens increased with increasing concentration of filtrate. The reduction percentages of the radial growth for the two pathogenic fungi were found to be 33.84, 46.46, 59.27, 65.58% and 39.61, 47.12, 57.06, 64.04% at 25, 50, 75, 100% of rhizobial filtrate, respectively, as determined by the agar well

diffusion method (Table 1). On the other side, the reduction percentage of fungal growth by the filtrate was found to be 54.11, 63.33, 67.11, 72.66% for *F. solani*, 46.33, 52.66, 59.66, 73.33% for *M. phaseolina* at the same concentration of rhizobial filtrate, respectively, when determined by a culture poisoning technique.

Effect of Rhizobium seed treatment on seed germination and root rot disease severity under field conditions

Results in Table 2 indicate that the pathogenic fungi, *F. solani*, *M. phaseolina* separately and in combination, affected highly seed germination, 28.5, 35.83, 27.50%, respectively, compared to 60.0% in the control. The coating of soybean seeds with *R. japonicum* increased the germination percentage significantly, 50.93, 56.5, and 50.13% in soil contaminated by *F. solani*, *M. phaseolina* and both, respectively. *R. japonicum* induced a high reduction in disease severity, 36.1, 27.7, 41.6% compared to 83.3, 74.9, and 86.0% on roots for the same previous combination, respectively.

Effect of rhizobial suspension on seed germination and root rot disease development in pots

Results of the effect of rhizobial suspension on seed germination and soybean root rot disease caused by *F. solani* and *M. phaseolina* separately or in combination, in contaminated soil are shown in Table 3. It was found that the two pathogenic fungi caused a high reduction in seed germination. The seed germination percentages were 40.0, 43.3, 46.6% in soil contaminated by *F. solani*, *M. phaseolina* and both, respectively, compared to 76.6% in the control.

The addition of *R. japonicum* suspension to the soil contaminated by the two pathogens improved seed germination and significantly reduced root rot disease severity. The germination percentage was 66.6, 73.3, 63.3% in rhizobial treated soil compared to *F. solani*, *M. phaseolina* and both, respectively.

The reduction of root rot disease severity caused by *F. solani* and *M. phaseolina* in the contaminated

Table 1. Effect of *Rhizobium japonicum* filtrate on the radial growth of *F. solani* and *M. phaseolina* estimated by agar well diffusion method on a culture medium

Filtrate concentration	<i>Fusarium</i> radial growth	Inhibition percentage of <i>Fusarium</i>	<i>Macrophomina</i> radial growth	Inhibition percentage of <i>Macrophomina</i>
100	1.8	65.58	2.06	64.04
75	2.3	59.27	2.24	57.06
50	2.8	46.46	3.03	47.12
25	3.46	33.84	3.46	39.61
Control	5.23	0	5.73	0
LSD ($P = 0.05$)	0.355		0.340	

Each value is a mean of 3 replicates

Table 2. Effect of *Rhizobium japonicum* on seed germination and root rot disease severity of soybean

Treatments	Germination (%)	Disease index (%)	
		on foliage	on roots
<i>F. solani</i>	28.50	79.96	83.3
<i>M. phaseolina</i>	35.83	68.83	74.9
<i>F. solani</i> + <i>M. phaseolina</i>	27.50	75.53	86.0
<i>R. japonicum</i> + <i>F. solani</i>	50.93	46.63	36.1
<i>R. japonicum</i> + <i>M. phaseolina</i>	56.50	43.23	27.7
<i>R. japonicum</i> + <i>F. solani</i> + <i>M. phaseolina</i>	50.13	51.00	41.6
<i>R. japonicum</i>	65.13	30.00	30.5
Control	60.00	30.00	33.3
LSD ($P = 0.05$)	5.06	9.42	16.26

Each value is a mean of 3 replicates

Table 3. Antagonistic activity of *R. japonicum* against *F. solani* and *M. phaseolina*

Treatments	Germination (%)	Disease severity	
		on foliage	on roots
Control	76.6	18.80	9.40
<i>F. solani</i>	40.0	82.20	77.76
<i>M. phaseolina</i>	43.3	50.00	69.40
<i>F. solani</i> + <i>M. phaseolina</i>	46.6	68.96	74.96
<i>R. japonicum</i> + <i>F. solani</i>	66.6	37.76	44.40
<i>R. japonicum</i> + <i>M. phaseolina</i>	73.3	34.43	36.10
<i>R. japonicum</i> + <i>F. solani</i> + <i>M. phaseolina</i>	63.3	55.53	38.83
<i>R. japonicum</i>	83.3	19.96	27.73
LSD ($P = 0.05$)	12.89	12.49	12.85

Each value is a mean of 3 replicates

soil was highly significant at $P = 0.05$, 37.76, 34.43, 55.53% on foliage with rhizobia, compared to 77.76, 69.4, 74.96% without rhizobia for the two pathogens separately and in combination respec-

tively. On the root system the disease severities were 44.4, 36.1, 38.83% with rhizobia compared to 77.76, 69.4, and 74.96% without rhizobia for the three treatments, respectively.

DISCUSSION

The results of this study revealed that *F. solani* and *M. phaseolina* are among the most important soil borne-pathogens infecting roots and causing up to 40% of plant mortality. The control of soil-borne pathogens is difficult since they produce resting structures such as chlamyospores and sclerotia resistant to adverse environmental conditions. The misuse of chemicals to control these pathogens caused enormous problems to ecosystem and human's health as well as has led to development of resistant races of pathogens (EL-BRAMAWY & ABDUL-WAHID 2006; EL-BATANONY *et al.* 2007).

The present study aims to protect soybean plants against soil-borne pathogens (*F. solani* and *M. phaseolina*) and improve growth and yields by using *Rhizobium japonicum*, an environmentally friendly alternative to fungicides. Results obtained showed that *Rhizobium japonicum* induced a high reduction in fungal radial growth on a culture medium as well as promoted plant growth under greenhouse and field conditions. On the other hand, root rot severity caused by these pathogens was also reduced by the addition of rhizobia to the contaminated soil.

The growth promotion induced by rhizobia may be directly through nitrogen fixation and production of plant growth regulators, i.e. substances that stimulate plant growth. Several researchers reported that rhizobia produced plant growth regulators such as indole acetic acid, auxins, cytokinins, gibberellin-like substances and rhizopine that stimulated and enhanced plant growth (TRIPLETT *et al.* 1981; ATZORN *et al.* 1988; HUSSAIN *et al.* 1990; SHENG 1993; MURPHY *et al.* 1995; NOEL *et al.* 1996; BODDEY & HUNGRIA 1997; DESHWAL *et al.* 2003; SHARIF *et al.* 2003). It was also reported that rhizobia increase P-availability to plants (DE FREITAS *et al.* 1997). In addition, the promotion of plant growth may indirectly suppress soil-borne pathogens by rhizobia metabolites. Several previous studies reported that rhizobia increase seed germination significantly and improve plant growth and yields through a reduction of soil-borne pathogens (SHEIKH *et al.* 2006; MAZEN *et al.* 2008).

The mechanism of rhizobia activity against *F. solani* and *M. phaseolina* may be due to the colonization of plant roots using root exudate for growth and synthesising metabolites protecting the roots against pathogens through antibiosis, degrading pathogenicity and inhibiting spore ger-

mination as well as induction of plant defence mechanisms. It was reported that rhizobia in the rhizosphere of plants may prevent the control of pathogenic fungi by covering hyphal tips of the fungus and produce antibiotics leading to the lysis of fungal hyphae (SHARIF *et al.* 2003). Several bacterial strains including rhizobia were isolated and used to control soil-borne pathogens, while they showed high efficiency under greenhouse and field conditions (NELSON 2004; SIDDIQUI 2006; AKHTAR & SIDDIQUI 2009). Rhizobia were reported to inhibit the growth of pathogenic fungi including *F. solani* and *M. phaseolina* in both leguminous and non-leguminous crops (EHTISHAMUL-HAQUE & GAFFAR 1993; OMAR & ABD-ALLA 1998).

Besides nitrogen fixation and promotion of plant growth, *Rhizobium japonicum* was found to exhibit high activity against *F. solani* and *M. phaseolina* causative agent of soybean root rot, and could be used as an important element in management of root rot disease.

References

- AKHTAR M.S., SIDDIQUI Z.A. (2009): Use of plant growth-promoting rhizobacteria for the biocontrol of root-rot disease complex of chickpea. *Australian Plant Pathology*, **38** (Suppl. 1): 44–50.
- AL-ANI R.A., DIWAN S.N.H., ADHAB M.A. (2010): Efficiency of *Thuja orientalis* and *Artemisia campestris* extracts to control of *Potato leaf roll virus* (PLRV) in potato plants. *Agriculture and Biology Journal of North America*, **1**: 579–583.
- AL-ANI R.A., ADHAB M.A., EL-MUADHIDI M.A., AL-FAHAD M.A. (2011a): Induced systemic resistance and promotion of wheat and barley plants growth by biotic and non-biotic agents against *Barley yellow dwarf virus*. *African Journal of Biotechnology* **10**: 12079–12084.
- AL-ANI R.A., ADHAB M.A., HAMAD S.A.H., DIWAN S.H.H. (2011b): *Tomato yellow leaf curl virus* (TYLCV), identification, virus vector relationship, strains characterization and control in Iraq. *African Journal of Agricultural Research*, **6**: 5149–5155.
- AL-ANI R.A., ADHAB M.A., HASSAN K.A. (2011c): Antiviral activity of Vit-org and 2-nitromethyl phenol and Thuja extract against *Eggplant blister mottled virus* (EBMV). *African Journal of Microbiology Research* **5**: 3555–3558.
- ATZORN R., CROZIER A., WHEELER C.T., SANDBERG G. (1988): Production of gibberellins and indole-3-acetic acid by *Rhizobium phaseoli* in relation to nodulation of *Phaseolus vulgaris* roots. *Planta*, **175**, 532–538.

- BAKKER P.A.H.M., PIETERSE C.M.J., VAN LOON L.C. (2007): Induced systemic resistance by fluorescent *Pseudomonas* spp. *Phytopathology*, **97** (Suppl. 2): 239–243.
- BLACK C.A. (1965): *Methods of Soil Analysis*. Part 1, 2. American Soil Agronomy, Inc., Madison.
- BODDEY L.H., HUNGARIA M. (1997): Phenotypic grouping of Brazilian *Bradyrhizobium* strains which nodulate soybean. *Biology and Fertility of Soils*, **25**: 407–415.
- BOOTH C. (1981): Perfect states (telemorphs) of *Fusarium* species. In: NELSON P.E., TOUSSOUN T.A., COOK R.J. (eds): *Fusarium Disease, Biology and Taxonomy*. Pennsylvania State University Press. University Park: 446–452.
- CHO J.H., RUPE J.C., CUMMINGS M.S., GBUR JR. E.E., (2001): Isolation and identification of *Fusarium solani* f.sp. *glycines* from soil on modified Nash and Snyder's medium. *Plant Disease*, **85**: 256–260.
- COOK R.J., GABRIEL C.J., KELMAN A., TOLIN S., VIDAVER A.K. (1995): Research on plant disease and pest management is essential to sustainable agriculture. *BioScience*, **45**: 354–357.
- DE FREITAS J.R., BANERJEE M.R., GERMIDA J.I. (1997): Phosphate solubilizing rhizobacteria enhance the growth and yield but not phosphorus uptake of canola (*Brassica napus* L.). *Biology and Fertility of Soils*, **24**: 358–364.
- DESHWAL V.K., DUBEY R.C., MAHESHWARI D.K. (2003): Isolation of plant growth promoting strains of *Bradyrhizobium* (*Arachis*) sp. with biocontrol potential against *Macrophomina phaseolina* causing charcoal rot of peanut. *Current Science*, **84**: 443–448.
- DIWAN S.N.H., AL-ANI R.A., ADHAB M.A. (2011): Systemic resistance induced in potato plants against *Potato virus Y* common strain (PVY^O) by plant extracts in Iraq. *Advances in Environmental Biology*, **5**: 209–215.
- EHTESHAMUL-HAQUE S., GHAFFAR A. (1993): Use of rhizobia in the control of root rot diseases of sunflower, okra, soybean and mungbean. *Journal of Phytopathology*, **138**: 157–163.
- EL-BATANONY N.H., MASSOUD O.N., MAZEN M.M., ABD EL-MONIUM M.M. (2007): The inhibitory effects of culture filtrates of some wild *Rhizobium* spp. on some faba bean root rot pathogens and their antimicrobial synergistic effect when combined with arbuscular mycorrhiza. *World Journal of Agricultural Sciences*, **3**: 721–730.
- EL-BRAMAWY M.A.S., ABDUL-WAHID O.A. (2006): Field resistance of crosses of sesame (*Sesamum indicum* L.) to charcoal root rot caused by *Macrophomina phaseolina* (Tassi.) Goid. *Plant Protection Science*, **42**: 66–72.
- FRIEDMAN M., BRANDON D. (2001): Nutritional and health benefits of soy proteins. *Journal of Agricultural and Food Chemistry*, **49**: 1069–1086.
- GHOSH A.C., BASU P.S. (1997): Culture growth and IAA production by microbial diazotrophic symbiont of stem nodules of the legume *Aeschynomene aspera*. *Folia Microbiologica*, **42**: 595–600.
- HAMMERSCHMIDT R. (1999): Induced disease resistance: how do induced plants stop pathogens? *Physiological and Molecular Plant Pathology*, **55**: 77–84.
- HAPGOOD F. (1987): The prodigious soybean. *National Geographic*, **172**: 66–91.
- HOLLIDAY P., PUNITHALINGAM E. (1970): *Macrophomina phaseolina*. C.M. Description of Pathogenic Fungi and Bacteria. No. 275. Commonwealth Mycological Institute. Kew.
- HUSSAIN S., GHAFFAR A., ASLAM M. (1990): Biological control of *Macrophomina phaseolina* charcoal rot of sunflower and mung bean. *Journal of Phytopathology*, **130**: 157–160.
- JOGAIAH S., SHIVANNA R.K., GNANAPRAKASH P.H., HUNTHRIKE S.S. (2010): Evaluation of plant growth-promoting Rhizobacteria for their efficiency to promote growth and induce systemic resistance in pearl millet against downy mildew disease. *Archives of Phytopathology and Plant Protection*, **43**: 368–378.
- KLOEPPER J.W., RYU C.-M., ZHANG S. (2004): Induced systemic resistance and promotion of plant growth by *Bacillus* species. *Phytopathology*, **94**: 1259–1266.
- LARA L.J.C., BAIÃO N.C., LÓPEZ C.A.A., MOURA B.H.S., RIBEIRO B.R.C. (2003): Fuentes de aceite en la ración de pollos de carne. In: XVIII Congreso Latinoamericano de Avicultura, Santa Cruz De La Sierra, Bolivia. (CD-ROM).
- MAZEN M.M., EL-BATANONY N.H., ABD EL-MONIUM M.M., MASSOUD O.N. (2008): Cultural filtrate of *Rhizobium* spp. and arbuscular mycorrhiza are potential biological control agents against root rot fungal diseases of faba bean. *Global Journal of Biotechnology and Biochemistry*, **3**: 32–41.
- MURPHY P.J., WEXLER W., GRZEMSKI W., RAO J.P., GORDON D. (1995): Rhizopines their role in symbiosis and competition. *Soil Biochemistry*, **27**: 525–529.
- NELSON L.M. (2004): Plant growth promoting rhizobacteria (PGPR): Prospects for new inoculants. [Online.] *Crop Management*. doi: 10.1094/CM-2004-0301-05-RV.
- NOEL T.C., SHENG C., YOST C.K., PHARIS R.P., HYNES M.F. (1996): *Rhizobium leguminosarum* as a plant growth promoting rhizobacterium: direct growth promotion of Canola and Lettuce. *Canadian Journal of Microbiology*, **42**: 279–283.
- OMAR S.A., ABD-ALLA M.H. (1998): Biocontrol of fungal root rot diseases of crop plants by the use of rhizobia and bradyrhizobia. *Folia Microbiologica*, **43**: 431–437.
- SCAIFE J.R., MOYO J., GALBRAITH H., MICHIE W., CAMPBELL V. (1994): Effect of different dietary supplemental fats and oils on the tissue fatty acid composition and growth of female broilers. *British Poultry Science*, **35**: 107–118.

- SHARIF T., KHALIL S., AHMAD S. (2003): Effect of *Rhizobium* sp., on growth of pathogenic fungi under *in vitro* conditions. Pakistan Journal of Biological Sciences, **6**: 1597–1599.
- SHEIKH L.I., DAWAR S., ZAKI M.J., GHAFFAR A. (2006): Efficacy of *Bacillus thuringiensis* and *Rhizobium meliloti* with nursery fertilisers in the control of root infecting fungi on mung bean and pkra plants. Pakistan Journal of Botany, **38**: 465–473.
- SHENG C. (1993): Hormones and direct effect of plant growth promoting rhizobacteria on higher plants. [Ph.D. Thesis.] University of Calgary, Alta.
- SIDDIQUI Z.A. (2006): PGPR: Prospective biocontrol agents of plant pathogens. In: SIDDIQUI Z.A. (ed.): PGPR: Biocontrol and Biofertilization. Springer, Dordrecht: 111–142.
- TRIPLETT E.W., HEITHOLT J.J., EVENSON K.B., BLAVINS D.G. (1981): Increase in internode length *Phaseolus lunatus* L., caused by inoculation with a nitrate reductase deficient strain of *Rhizobium* sp. Plant Physiology, **67**: 1–4.
- VAN LOON L.C. BAKKER P.A.H.M., PIETERSE C.M.J. (2004): Systemic resistance induced by rhizosphere bacteria. Annual Reviews of Phytopathology **36**, 453–483.
- VEVERKA K., PALICOVÁ J., KŘÍŽKOVÁ I. (2008): The incidence and spreading of *Macrophomina phaseolina* (Tassi) Goidanovich on sunflower in the Czech Republic. Plant Protection Science, **44**: 127–137.
- VIEIRA S.L., RIBEIRO A.M.L., KESSLER A.M., FERNANDES L.M., EBERT A.R., EICHNER G. (2002): Utilização da energia de dietas para frangos de corte formulados com óleo ácido de soja. Revista Brasileira Ciência Avícola, **4**(2): 1–13.
- WOLTZ S.S., ARTHUR W.E. (1973): Fusarium wilt of chrysanthemum, effect of nitrogen source and lime and disease development. Phytopathology, **63**: 155–157.
- YANG X.B., FENG F. (2001): Ranges and diversity of soybean fungal diseases in North America. Phytopathology, **91**: 769–775.
- ZEMÁNKOVÁ M., LEBEDA A. (2001): Soil *Fusarium* species, their taxonomy, variability and significance in plant pathology. Plant Protection Science. **37**: 24–42.

Received for publication March 24, 2012
Accepted after corrections May 3, 2012

Corresponding author:

Dr. MUSTAFA A. ADHAB, University of Baghdad, College of Agriculture, Plant Protection Department, Baghdad, Iraq
tel. + 96 479 041 680 48, e-mail: maa_adhab@hotmail.com
