

Effects of dietary *Enterococcus faecium* on growth performance, carcass characteristics, faecal microbiota, and blood profile in broilers

M. MOHAMMADI GHEISAR, A. HOSSEINDOUST, I.H. KIM

Dankook University, Cheonan, Choongnam, Republic of Korea

ABSTRACT: This research was performed to evaluate the effect of supplementing broiler diets with a probiotic containing *Enterococcus faecium* on growth performance, carcass characteristics, faecal microbiota, and blood profile. A total of 384 one-day-old Ross 308 broiler chicks (mixed gender) with an average initial BW of 39.2 g were used in a 35 days feeding trial. The chicks were allotted to pens with 16 birds per pen and eight replications per treatment with food and water provided *ad libitum*. Treatments were: (1) basal diet, (2) 0.25% probiotic, and (3) 0.5% probiotic. Results indicated that body weight gain (BWG) on Day 7 to 21, Day 21 to 35 and overall (0 to 35) increased ($P < 0.05$) linearly but feed intake (FI) and feed conversion ratio (FCR) were not affected. A linear increase ($P < 0.05$) was observed in the relative weight of breast muscle when comparing the 0 to 0.5% concentration of probiotic, but breast meat colour was not affected by treatments. A significant impact (linear effect, $P < 0.05$) was observed on drip loss on Day 1. Inclusion of probiotic decreased ($P < 0.05$) the count of *Salmonella* linearly but the counts of *E. coli* and *Lactobacillus* were not affected. There was no remarkable influence on blood profile. Thus, it was concluded that inclusion of a probiotic containing *Enterococcus faecium* improved growth performance and altered the intestinal microbial population, without any negative effects on meat colour and blood profile in broiler chickens.

Keywords: broiler; carcass characteristics; *Enterococcus faecium*; faecal microbiota

Since antibiotics as growth promoters are being removed from poultry and swine diets worldwide, there is a pressing requirement to find alternatives. Several natural products, such as organic acids, plant extracts, probiotics, and prebiotics, have been assessed as alternatives to antibiotics as growth promoters (Patterson and Burkholder 2003; Higgins et al. 2008; Markovic et al. 2009; Mountzouris et al. 2010; Vondruskova et al. 2010; Zhang et al. 2013; Zhang and Kim 2013). Probiotics are live microbial feed additives that can beneficially influence the intestinal microflora of the host animal. Various microorganisms such as *Lactobacillus*, *Lactococcus*, *Streptococcus*, *Bacillus*, and *Enterococcus* can be used as probiotics (Fooks and Gibson 2002; Ouwehand et al. 2002; Lodemann et al. 2006; Park and Kim 2015; Zhang et al. 2014). Previous studies have confirmed the positive effects of probiotics in pigs (Meng et al. 2010; Yan and Kim 2013). Several studies have been conducted to investigate the effects of supplement-

ing poultry diets with probiotics and it has been reported that probiotics can exert positive effects on the development and function of immune cells (Huang et al. 2004; Kabir et al. 2004). Recently, some probiotic feed additives have been produced that contain viable cells of *Enterococcus faecium*. These products are currently authorised for use in piglets and calves (Vahjen et al. 2007).

This study was conducted to investigate the effects of supplementing broiler diets with different concentrations of a probiotic containing *Enterococcus faecium* M74 on growth performance, meat quality, relative organ weights, faecal microbiota, and blood profile.

MATERIAL AND METHODS

Animals, diets, and facilities. The use and management of the broiler chickens used in this

doi: 10.17221/8680-VETMED

study were approved by the Animal Care and Use Committee of Dankook University. A total of 384 one-day-old Ross 308 broiler chicks (mixed sexes) with an average initial BW of 39.2 g were used in a 35 day experimental period. The chicks were allotted to pens with 16 birds per pen and eight pens per treatment. Treatments were: (1) basal diet, (2) 0.25% pro-

biotic (3×10^{11} CFU/g), and (3) 0.5% probiotic (3×10^{11} CFU/g). The chicks were weighed and placed randomly in three floor battery cages in an environmentally controlled room (32 to 24 °C and 65% relative humidity). During the entire experimental period, the chickens were provided access to feed and water *ad libitum*. All diets were formulated to meet or exceed NRC recommendations for nutrient recommendations (1994). Feed ingredients and the chemical composition are presented in Table 1. Relative weight of breast meat, abdominal fat and organs were described as a percentage of live weight. The probiotic used in this experiment contained 3×10^{11} CFU of *Enterococcus faecium* M74 per gram (Lactiferm[®], Chr. Hansen, Germany).

Sampling and measurements. All the chickens and the remaining feed were weighed on Days 0, 7, 21, and 35 to allow calculations of body weight gain (BWG), feed intake (FI), and feed conversion ratio (FCR). On the last day of the trial period, 30 chickens were selected randomly and blood samples were taken from the wing vein (six chickens per treatment). Blood samples were collected into K₃EDTA vacuum tubes (Becton Dickinson Vacutainer Systems, Franklin Lake, NJ). The samples were centrifuged (3000 × g, 15 min) to recover blood plasma. Whole blood cell counts (white blood cells (WBC), red blood cells (RBC), and lymphocytes) were analysed using an automatic blood analyser (ADVIA 120, Bayer, NY). After blood sample collection, the same chickens were weighed individually and sacrificed. Breast meat, abdominal fat, gizzard, liver, spleen, bursa of Fabricius, and heart were excised, blotted to remove excess moisture, and weighed by trained personnel. Hunter *L** (lightness), *a** (redness), and *b** (yellowness) of breast meat were determined using a Minolta CR410 chromameter (Konica Minolta Sensing Inc., Osaka, Japan). Drip loss percentage was measured on Days 1, 3, 5, and 7 using approximately 2 g of breast meat sample according to the plastic bag method, described by Honikel (1998). Faecal samples from the cloacae were collected into micro-tubes and were analysed for counts of *Lactobacillus*, *E. coli*, and *Salmonella* using agar media. Viable bacteria in excreta samples were determined by plating 10-fold serial dilutions (in 1% peptone solution) onto MacConkey agar plates (Difco Laboratories, Detroit, MI) and *Lactobacilli* medium agar plates (Medium 638, DSMZ, Braunschweig, Germany) to isolate the *E. coli* and *Lactobacillus*, respectively.

Table 1. Composition of diets (as-fed basis)

Item (%)	Phase 1 Day 0–7	Phase 2 Day 7–21	Phase 3 Day 21–35
Corn	45.48	36.69	40.80
Wheat	10.00	20.00	20.00
Soybean meal (CP 48%)	34.25	33.62	25.48
Corn gluten meal (CP 60%)	2.00	–	–
Rape seed meal	–	–	3.50
Tallow	1.90	5.54	6.01
Soybean oil	1.50	–	–
Limestone	1.06	1.12	1.17
Dicalcium phosphate	2.23	1.90	1.84
Salt	0.35	0.32	0.29
DL-Methionine	0.46	0.39	0.41
L-Lysine-HCl	0.42	0.15	0.20
Threonine	0.17	0.09	0.12
Vitamin mix ¹	0.03	0.03	0.03
Vitamin E (10%)	0.04	–	–
Mineral mix ²	0.10	0.10	0.10
CuSO ₄ ·5 H ₂ O	0.01	0.05	0.05
Total	100	100	100
Calculated nutritional content			
ME (MJ/kg)	12.62	13.03	13.31
Analysed nutritional content (%)			
CP	22.12	20.43	18.55
Lysine	1.45	1.22	1.10
Met + Cys	1.06	0.95	0.93
Ca	1.05	1.00	1.00
Available P	0.53	0.50	0.50
Crude fat	5.55	7.27	7.96
Crude fiber	3.24	3.29	3.25

¹ Provided per kg of diet: 15 000 IU of vitamin A, 3750 IU of vitamin D₃, 37.5 mg of vitamin E, 2.55 mg of vitamin K₃, 3 mg of B₁, 7.5 mg of B₂, 4.5 mg of vitamin B₆, 24 µg of vitamin B₁₂, 51 mg of niacin, 1.5 mg of folic acid, 126 mg of biotin and 13.5 mg of pantothenic acid

² Provided per kg of complete diet: 37.5 mg of Zn, 137.5 mg of Mn, 37.5 mg of Fe, 0.83 mg of I, and 0.23 mg of Se, and 1.408 mg of choline

The *Lactobacilli* medium agar plates were then incubated for 48 h at 39 °C under anaerobic conditions. The MacConkey agar plates were incubated for 24 h at 37 °C. The *E. coli* and *Lactobacillus* colonies were counted immediately after removal from the incubator. For *Salmonella*, the serially diluted peptone broth tubes were incubated overnight at 37 °C, after which 1 ml was transferred to 9 ml of tetrastinate broth (Neogen Corporation, Lansing, MI) followed by incubation for 48 h at 42 °C. From these tubes, 1 ml was used to inoculate 9 ml of Rappaport-Vassiliadis *Salmonella* Enrichment broth (Neogen Corporation, Lansing, MI) followed by incubation for 48 h at 42 °C. The Rappaport was used to inoculate XLT4 plates for *Salmonella* isolation, and *Salmonella* was identified using LIS (VIDAS Listeria) and TSI (Triple Sugar Iron) agar tubes (Difco Laboratories, Detroit, MI).

Statistical analysis. Data were analysed as a completely randomised design using the mixed procedures of SAS (SAS Institute 1996). Mean values and standard errors (SE) are reported. Linear and quadratic polynomial contrasts were performed to determine the effects of inclusion of 0, 0.25, and 0.5% *Enterococcus faecium* M74-containing probiotic in the diets.

RESULTS

Growth performance

The data presented in Table 2 indicate that from Day 1 to 7, there were no significant effects on body weight gain (BW), feed intake (FI), and feed conversion ratio (FCR). From Day 7 to 21, BW increased ($P = 0.038$) linearly in response to supplementation with 0 to 0.5% of the probiotic containing *Enterococcus faecium*, but FI and FCR were not affected. From Day 21 to 35 a linear increase ($P = 0.001$) was observed on BW, without any significant effect on FI and FCR. During the entire experimental period, BW increased ($P = 0.005$) and there was a trend for linearly increasing FI ($P = 0.061$), but FCR was not affected by experimental treatments.

Carcass characteristics

Supplementation with 0 to 0.5% probiotic containing 3×10^{11} CFU/g *Enterococcus faecium* resulted in a linear ($P = 0.01$) increase in relative weight breast muscle. Supplementing the diets with 0 to 0.5% dietary *Enterococcus faecium* had no re-

Table 2. Effect of dietary supplementation of probiotic containing *Enterococcus faecium* on growth performance in broilers

	Probiotic (%)			SE ¹	P-value	
	0	0.25	0.5		linear	quadratic
Day 1 to 7						
BWG (g) ²	99.3	105.6	102.3	1.9	0.269	0.056
FI (g) ³	129.3	130.9	132.2	3	0.704	0.582
F:G	1.30	1.25	1.28	0.03	0.734	0.352
Day 7 to 21						
BWG (g)	579.3	616.2	623.5	11.5	0.038	0.088
FI (g)	886.1	897.7	910.9	13.3	0.208	0.960
F:G	1.53	1.44	1.48	0.03	0.265	0.083
Day 21 to 35						
BWG (g)	1020.9	1075.2	1113.5	22.2	0.001	0.77
FI (g)	1753.4	1778.7	1800.9	21.8	0.144	0.957
F:G	1.72	1.66	1.62	0.04	0.105	0.753
Overall						
BWG (g)	1699.5	1804.3	1832.1	28.7	0.005	0.291
FI (g)	2768.8	2808.4	2842.7	25.8	0.061	0.933
F:G	1.63	1.56	1.55	0.03	0.071	0.302

¹standard error, ²gain in BW per bird, ³feed intake per bird, number of observations per mean: 6

doi: 10.17221/8680-VETMED

Table 3. Effect of dietary supplementation of probiotic containing *Enterococcus faecium* on relative breast meat, abdominal fat and organ weights in broilers

Body part (%)	Probiotic (%)			SE ¹	P-value	
	0	0.25	0.5		linear	quadratic
Breast muscle	7.08	8.09	8.80	0.42	0.01	0.76
Abdominal fat	1.56	1.14	1.27	0.15	0.19	0.14
Gizzard	1.17	1.16	1.17	0.05	0.96	0.83
Heart	0.49	0.42	0.43	0.03	0.11	0.25
Liver	2.05	2.13	2.14	0.12	0.60	0.81
Spleen	0.10	0.10	0.09	0.02	0.79	0.50
Bursa of Fabricius	0.12	0.13	0.14	0.02	0.55	0.86

¹standard error, number of observations per mean: 6

markable effect on relative weights of organs and abdominal fat (Table 3).

Faecal microbiota

The data presented in Table 4 shows that increasing concentrations (0 to 0.5%) of dietary *Enterococcus faecium* decreased ($P = 0.008$) the count of faecal *Salmonella* linearly. Supplementing diets with 0 to 0.5% of probiotic containing *Enterococcus faecium* resulted in a linear decrease ($P = 0.008$) in faecal *Salmonella* counts. The inclusion of the probiotic had no marked influence on faecal *E. coli* or *Lactobacillus* counts.

Blood characteristics

Whole blood cells and haptoglobin concentrations were not affected by dietary treatments (Table 5).

DISCUSSION

Various probiotics have been assessed in pigs and poultry, and most studies have reported that supplementing diets with probiotics represents a viable alternative to antibiotics for improved growth performance without adverse effects on mortality in poultry or pigs (Fairchild et al. 2001; Hooge

Table 4. Effect of dietary supplementation of probiotic containing *Enterococcus faecium* on faecal microbiota in broilers

Bacterial counts (log ₁₀ cfu/g)	Probiotic (%)			SE ¹	P-value	
	0	0.25	0.5		linear	quadratic
<i>Lactobacillus</i>	7.65	7.78	7.80	0.08	0.22	0.54
<i>E. coli</i>	6.56	6.43	6.46	0.06	0.25	0.35
<i>Salmonella</i>	2.72	2.59	2.57	0.04	0.01	0.11

¹standard error, number of observations per mean: 6

Table 5. Effect of dietary supplementation of probiotic containing *Enterococcus faecium* on blood constituents in broilers

Constituents	Probiotic (%)			SE ¹	P-value	
	0	0.25	0.5		linear	quadratic
WBC (10 ³ /μl)	458.50	497.90	635.20	98.8	0.22	0.69
RBC (10 ⁶ /μl)	2.68	2.65	2.71	0.07	0.76	0.63
Lymphocyte (%)	81.78	71.78	71.32	11.9	0.54	0.75
Haptoglobin (mg/l)	161.6	173.3	163.3	1.4	0.93	0.54

¹standard error, number of observations per mean: 6

et al. 2004). The results of the current study were consistent with previous studies and demonstrated that supplementing the diet with 0.5% probiotic improved the growth performance over the whole experimental period. Other studies have also shown that supplementing diets with probiotics can improve the growth performance of broilers, and that the inclusion of probiotics may enhance the activity of digestive enzymes, such as proteases, lipases, and amylases, resulting in better nutrient utilisation and consequently improved growth performance (Fuller 2001).

Zamanzad-Ghavidel et al. (2011) reported that relative breast meat in chickens fed a diet containing *Lactobacillus*-based probiotic was higher than in those that did not receive the probiotic, consistent with the findings of the current study. Also Zheng et al. (2015) reported that feeding broiler chickens diets containing *Enterococcus faecium* led to a significant improvement in breast muscle yield. They suggested that the main effects of *Enterococcus faecium* occur in the intestine through modulation of the intestinal microbiota in favour of the host animal and through improved mucosa ultrastructure, enhanced nutrient absorption and reduced energy consumption.

Probiotics can be considered as modulators of the gut environment as they increase the population of beneficial micro-organisms and inhibit the proliferation of pathogens in the intestinal microbiota; consequently, they can improve growth performance (Patterson and Burkholder 2003; Anjum et al. 2005; Higgins et al. 2008). Several studies have confirmed the stimulatory effects of probiotics on the intestinal microbiota (Roth et al. 1992; Depta et al. 1998; Mathew et al. 1998; Jadamus et al. 2001; Scharek et al. 2005; Reiter et al. 2006; Scharek et al. 2007; Lodemann et al. 2008). Pajarillo et al. (2015) reported that supplementation of *Enterococcus faecium* NCIMB 11181 to a swine diet significantly increased faecal *Lactobacilli* counts and reduced *E. coli* counts. In agreement with our findings, Chen et al. (2005) reported that supplementing growing pig diets with a probiotic did not affect whole blood cell counts.

In conclusion, the current study demonstrates that inclusion of a probiotic containing *Enterococcus faecium* improves the growth performance and favourably alters the intestinal microbiota by increasing *Lactobacilli* and decreasing *E. coli* and *Salmonella* populations in broilers. Importantly,

inclusion of the probiotic in the diet does not exert any negative effects on breast muscle colour, relative weights of organs and blood profile.

REFERENCES

- Anjum MI, Khan AG, Azim A, Afzal M (2005): Effect of dietary supplementation of multi-strain probiotic on broiler growth performance. *Pakistan Veterinary Journal* 25, 25–29.
- Chen YJ, Son KS, Min BJ, Cho JH, Kwon OS, Kim IH (2005): Effects of dietary probiotic on growth performance, nutrients digestibility, blood characteristics and fecal noxious gas content in growing pigs. *Asian Australian Journal of Animal Science* 18, 1464–1468.
- Depta A, Rychlik R, Nieradka R, Rotkiewicz T, Kujawa K, Bomba A, Grabowska-Swiecicka G (1998): The influence of alimentary tract colonization with *Lactobacillus* sp. strains on chosen metabolic profile indices in piglets. *Polish Journal of Veterinary Science* 1, 3–7.
- Fairchild AS, Grimes JL, Jones FT, Wineland MJ, Edens FW, Sefton AE (2001): Effects of hen age, Bio-Mos and flavomycin on poult susceptibility to oral *Escherichia coli* challenge. *Poultry Science* 80, 562–571.
- Fooks LJ, Gibson GR (2002): Probiotics as modulators of the gut flora. *British Journal of Nutrition* 88, 39–49.
- Fuller R (2001): The chicken gut microflora and probiotic supplements. *Poultry Science* 38, 189–196.
- Higgins SE, Higgins JP, Wolfenden AD, Henderson SN, Torres-Rodriguez A, Tellez G, Hargis B (2008): Evaluation of a lactobacillus-based probiotic culture for the reduction of *Salmonella enteritidis* in neonatal broiler chicks. *Poultry Research* 87, 27–31.
- Honikel KO (1998): Reference methods for the assessment of physical characteristics of meat. *Meat Science* 49, 447–457.
- Hooge DM, Ishimaru H, Sims MD (2004): Influence of dietary *Bacillus subtilis* C-3102 spores on live performance of broiler chickens in four controlled pen trials. *Journal of Applied Poultry Research* 13, 222–228.
- Huang MK, Choi JL, Houde R, Lee JW, Lee B, Zhao X (2004): Effects of lactobacilli and an acidophilic fungus on the production in Nigeria. *World Journal of Microbiology and Biotechnology* 20, 51–56.
- Jadamus A, Vahjen W, Simon O (2001): Growth behavior of a spore forming probiotic strain in the gastrointestinal tract of broiler chicken and piglets. *Archives of Animal Breeding* 54, 1–17.
- Kabir SML, Rahman MM, Rahman MB, Rahman MM, Ahmen SU (2004): The dynamics of probiotics on growth

doi: 10.17221/8680-VETMED

- performance and immune response in broilers. *International Journal of Poultry Science* 3, 361–364.
- Lodemann U, Hubener K, Jansen N, Martens H (2006): Effects of *Enterococcus faecium* NCIMB 10415 as probiotic supplement on intestinal transport and barrier function of piglets. *Archives of Animal Nutrition* 1, 35–48.
- Lodemann U, Lorenz BM, Weyrauch KD, Martens H (2008): Effects of *Bacillus cereus* var. *toyoi* as probiotic feed supplement on intestinal transport and barrier function in piglets. *Archives of Animal Nutrition* 62, 87–106.
- Markovic R, Sefer D, Krstic M, Petrujkic B (2009): Effect of different growth promoters on broiler performance and gut morphology. *Archivos de Medicina Veterinaria* 41, 163–169.
- Mathew AG, Chattin SE, Robbins CM, Golden DA (1998): Effects of a direct-fed yeast culture on enteric microbial populations, fermentations acids, and performance of weanling pigs. *Journal of Animal Science* 76, 2138–2145.
- Meng QW, Yan L, Ao X, Zhou TX, Wang JP, Wang J, Lee H, Kim IH (2010): Influence of probiotic in different energy and nutrient density diets on growth performance, nutrient digestibility, meat quality, and blood characteristics in growing finishing pig. *Journal of Animal Science* 88, 3320–3326.
- Mountzouris KC, Tsitsrikos P, Palamidi I, Arvaniti A, Mohnl M, Schatzmayr G, Fegeros K (2010): Effect of probiotic inclusion levels in broiler nutrition on growth performance, nutrient digestibility, plasma immunoglobulins, and cecal microflora composition. *Poultry Science* 89, 58–67.
- NRC – National Research Council (1994): Nutrient requirements of poultry. 9th rev. ed. National Academies Press, Washington, DC.
- Ouwehand AC, Salminen S, Isolauri E (2002): Probiotics: an overview of beneficial effects. *Antonie van Leeuwenhoek* 82, 279–289.
- Pajarillo EAB, Chae JP, Balolong MP, Kim HB, Park CS, Kang DK (2015): Effects of probiotic *Enterococcus faecium* NCIMB 11181 administration on swine fecal microbiota diversity and composition using barcoded pyrosequencing. *Animal Feed Science and Technology* 201, 80–88.
- Park JH, Kim IH (2015): The effects of the supplementation of *Bacillus subtilis* RX7 and B2A probiotics on the performance, blood profiles, intestinal *Salmonella* microflora, noxious gas emission, organ weight, and breast meat quality of broiler challenged with *Salmonella typhimurium*. *Journal of Animal Physiology and Animal Nutrition*, 99, 326–334.
- Patterson JA, Burkholder KM (2003): Application of prebiotics and probiotics in poultry production. *Poultry Science* 82, 627–631.
- Reiter K, Eggebrecht S, Drewes B, Riess M, Weyrauch KD (2006): Effect of *Enterococcus faecium* and *Bacillus cereus* var. *toyoi* on the morphology of the intestinal mucous membrane in piglets. *Biologia Bratislava* 61, 1–7.
- Roth FX, Kirchgessner M, Eidelsburger U, Gedek B (1992): Nutritive efficacy of *Bacillus cereus* as a probiotic in the fattening of calves. 1. Influence on fattening performance, carcass criteria and microbial activity in the small intestine. *Agribiological Research-Zeitschrift für Agrarbiologie Agrikulturchemie Ökologie* 45, 294–302.
- SAS Institute (1996): SAS User's Guide: Statistics. Version 7.0. SAS Institute Inc., Cary, NC.
- Scharek L, Guth J, Reiter K, Weyrauch KD, Tara D, Schwerk P, Schierack P, Schmidt MF, Wieler LH, Tedin K (2005): Influence of a probiotic *Enterococcus faecium* strain on development of the immune system of sows and piglets. *Veterinary Immunology and Immunopathology* 105, 151–161.
- Scharek L, Altherr BJ, Tolke C, Schidt MFG (2007): Influence of the probiotic *Bacillus cereus* var. *toyoi* on the intestinal immunity of piglets. *Veterinary Immunology and Immunopathology* 120, 136–147.
- Vahjen W, Taras T, Simon O (2007): Effect of the probiotic *Enterococcus faecium* NCIMB10415 on cell numbers of total *Enterococcus* spp., *E. faecium* and *E. faecalis* in the Intestine of piglets. *Current Issues of Intestinal Microbiology* 8, 1–8.
- Vondruskova H, Slamova R, Trckova M, Zraly Z, Pavlik I (2010): Alternatives to antibiotic growth promoters in prevention of diarrhoea in weaned piglets: a review. *Veterinarni Medicina* 55, 199–224.
- Yan L, Kim IH. (2013): Effects of probiotics supplementation in diets with different nutrient densities on growth performance, nutrient digestibility, blood characteristics, faecal microbial population and faecal noxious gas content in growing finishing pigs. *Journal of Applied Animal Research* 41, 23–28.
- Zamanzad-Ghavidel S, Nazer Adl K, Maheri Sis N, Aharizad S, Mirzaei Aghsaghali A, Mohammadian M, Siadati SA. (2011): Effects of lactobacillus-based probiotic on growth performance, mortality rate and carcass yield in broiler chickens. *Annals of Biological Research* 2, 325–331.
- Zhang ZF, Kim IH (2013): Effects of probiotic supplementation in different energy and nutrient density diets on performance, egg quality, excreta microflora, excreta noxious gas emission, and serum cholesterol concentrations in laying hens. *Journal of Animal Science* 91, 4781–4787.
- Zhang ZF, Cho JH, Kim IH (2013): Effects of *Bacillus subtilis* UBT-MO₂ on growth performance, relative immune organ weight, gas concentration in excreta, and intestinal microbial shedding in broiler chickens. *Livestock Science* 155, 343–347.

doi: 10.17221/8680-VETMED

Zhang ZF, Lee MJ, Kim IH (2014): Effects of *Enterococcus faecium* DSM 7134 on weanling pigs were influenced by dietary energy and crude protein density. *Livestock Science* 169, 106–111.

Zheng A, Lou J, Meng K, Li J, Zhang S, Li K, Liu G, Cai H, Bryden WL, Yao B (2015) Proteome changes underpin

improved meat quality and yield of chickens (*Gallus gallus*) fed the probiotic *Enterococcus faecium*. *BMC Genomics* 15, 1167.

Received: 2014–12–05

Accepted after corrections: 2015–11–30

Corresponding Author:

I.H. Kim, Dankook University, Department of Animal Resource and Science, Cheonan, Choongnam, 330-714 Republic of Korea

E-mail: inhokim@dankook.ac.kr
