

## Effects of lactulose on growth, carcass characteristics, faecal microbiota, and blood constituents in broilers

M. MOHAMMADI GHEISAR<sup>1</sup>, C.M. NYACHOTI<sup>2</sup>, J.D. HANCOCK<sup>3</sup>, I.H. KIM<sup>1</sup>

<sup>1</sup>Dankook University, Cheonan, Choongnam, Republic of Korea

<sup>2</sup>University of Manitoba, Winnipeg, Manitoba, Canada

<sup>3</sup>Kansas State University, Manhattan, Kansas, USA

**ABSTRACT:** This study was conducted to determine the effect of supplementing diets with lactulose on growth performance, carcass characteristics, faecal microbiota, and blood constituents. A total of 324 one-day-old Ross 308 mixed-sex broiler chicks with an average initial body weight of 38 g were used in a 35-day growth assay. There were 18 birds/pen and six pens/treatment with food and water available *ad libitum*. Treatments consisted of a corn-soybean-meal-based diet with 0, 0.25 and 0.5% of lactulose. The results indicated that body weight gain (BWG) was improved (linear effect,  $P < 0.05$ ) by increasing the concentration of lactulose in the diet from zero to 0.5% while the feed conversion ratio (FCR) decreased (linear effect,  $P < 0.05$ ) for Days 8 to 21, 21 to 35, and overall (Day 0 to 35). Chickens fed the diet supplemented with 0.5% lactulose showed a higher relative weight of breast meat compared to other groups. Inclusion of lactulose decreased the count of *Salmonella* and *E. coli* in excreta of chickens fed diets containing 0.25 or 0.5% lactulose ( $P < 0.05$ ), but the count of *Lactobacillus* was not affected. Drip loss percentage was decreased ( $P < 0.05$ ) on Day 1 by addition of 0.5% lactulose, but there was no effect on meat colour. Blood characteristics were not influenced. Thus, it was concluded that inclusion of lactulose improves growth performance and alters excreta microbial populations with no adverse effect on broilers.

**Keywords:** broiler; carcass characteristics; excreta microbiota; growth performance; lactulose

Concern about antibiotics as growth promoters has led to a desire to identify alternatives. Prebiotics, probiotics, and synbiotics are being increasingly adopted as promoters of growth and gut health in poultry and swine (Kleesen et al. 2001; Patterson and Burkholder 2003; Higgins et al. 2008; Markovic et al. 2009; Zhang et al. 2013; Zhang and Kim 2013; Zhao et al. 2013). Huang et al. (2004) reported that supplementing diets with a probiotic (*Lactobacillus acidophilus* and *Lactobacillus casei*) may enhance the development and function of immune cells in calves.

Prebiotics are feed additives which may stimulate the growth and activity of beneficial microorganisms such as *Bifidobacteria* and *Lactobacillus* in the gut (Cummings and MacFarlane 2002). Steiner (2006) reported that prebiotics may improve performance and nutrient digestibility by creating suitable conditions for beneficial microorganisms. Kermanshahi and Rostami (2006) reported that using dried whey

as a substrate for acidophilic flora (e.g. *Lactobacilli*) as prebiotics or fermentable sugars improved the useful microbial population of the gastrointestinal tract. The results of other studies have demonstrated that supplementing diets with probiotics can improve the broiler growth performance and may enhance the activity of digestive enzymes such as proteases, lipases, and amylases resulting in better nutrient utilisation and consequently, improved growth performance (Fuller 2001).

Several carbohydrates including NSPs, resistant starch and non-digestible oligosaccharides that may be fermented by intestinal microorganisms can be classified as prebiotics (Bauer et al. 2006). Lactulose (4-*O*- $\beta$ -*D*-galactopyranosyl-*D*-fructose) is a synthetic disaccharide that is classified as a non-digestible carbohydrate (Bird et al. 1990). However, it can be metabolised in the colon by saccharolytic microbiota. A benefit of using non-

doi: 10.17221/8722-VETMED

digestible carbohydrates is stimulation of growth of *Lactobacilli* in the gut and eventual fermentation by *Lactobacilli*, *Bifidobacteria*, and other bacteria species in the hind-gut (Mitsuoka et al. 1987; Fleige et al. 2007). Experiments with pigs indicate that lactulose can indeed not be digested and absorbed in the small intestine and that it passes to the large intestine where microorganisms use it to produce acetic and lactic acid. Consequently, lactulose can stimulate the growth of *Lactobacillus* and *Bifidobacterium* and reduce the activity of proteolytic bacteria (Gibson 2004; Marinho et al. 2007). The ingestion of lactulose has been reported to exert beneficial effects by increasing probiotic bacteria and putrefactive bacteria and significantly reducing potential pathogens and consequently reducing the activity of pro-carcinogenic enzymes (e.g. azoreductase, 7- $\alpha$ -dehydroxylase) in humans, mice, rats, sows and pigs (Ballongue et al. 1997; Bianchi et al. 1997; Krueger et al. 2002). It was hypothesised that through its prebiotic effects, dietary lactulose will improve the performance of broiler chickens. Thus, the objective of the current study was to determine the effect of supplementing broiler diets with lactulose on growth performance, carcass characteristics, excreta microbiota, and blood constituents.

## MATERIAL AND METHODS

**Animals, diets, and facilities.** Use and management of the broiler chickens used in this experiment were approved by the Animal Care and Use Committee of Dankook University. In this study 324 one-day-old Ross 308 broiler chicks (mixed gender) with an average initial BW of 37.8 g were used in a 35 days growth assay. Chicks were sorted into pens with 18 birds/pen and six pens/treatment. Treatments consisted of a corn-soybean-meal-based diet with 0, 0.25 or 0.5% of lactulose. Diets (Table 1) were fed in three phases with ME 3015, 3114, and 3180 Kcal/kg, Met + Cys 1.06, 0.95, and 0.93%, and P 0.53, 0.50, and 0.50% for phase 1 (Day 0 to 7), phase 2 (Day 7 to 21), and phase 3 (Day 21 to 35), respectively. All other nutrients met or exceeded nutrient concentrations recommended by the National Research Council (NRC 1994). Birds were housed in battery cages (1.75 × 1.55 m/pen) in an environmentally controlled room (temperature started at 32 °C and was reduced by

Table 1. Composition of diets in percent, (as-fed basis)

Item (%)	Phase 1	Phase 2	Phase 3
	Day 0–7	Day 7–21	Day 21–35
Corn	45.49	36.74	40.85
Wheat	10.00	20.00	20.00
Soybean meal (CP 48%)	34.25	33.62	25.48
Corn gluten meal (CP 60%)	2.00	–	–
Rapeseed meal	–	–	3.50
Tallow	1.90	5.54	6.01
Soybean oil	1.50	–	–
Limestone	1.06	1.12	1.17
Dicalcium phosphate	2.23	1.90	1.84
Salt	0.35	0.32	0.29
DL-Methionine	0.46	0.39	0.41
L-Lysine-HCl	0.42	0.15	0.20
Threonine	0.17	0.09	0.12
Vitamin mix <sup>1</sup>	0.03	0.03	0.03
Vitamin E (10%)	0.04	–	–
Mineral mix <sup>2</sup>	0.10	0.10	0.10
Total	100	100	100
<b>Calculated nutritional content</b>			
ME (kcal/kg)	3015	3114	3180
<b>Analysed nutritional content (%)</b>			
CP	22.12	20.43	18.55
Lysine	1.45	1.22	1.10
Met + Cys	1.06	0.95	0.93
Ca	1.05	1.00	1.00
Available P	0.53	0.50	0.50
Crude Fat	5.55	7.27	7.96
Crude fibre	3.24	3.29	3.25

<sup>1</sup>Provided per kg of diet: 15 000 IU of vitamin A, 3750 IU of vitamin D<sub>3</sub>, 37.5 mg of vitamin E, 2.55 mg of vitamin K<sub>3</sub>, 3 mg of B<sub>1</sub>, 7.5 mg of B<sub>2</sub>, 4.5 mg of B<sub>6</sub>, 24  $\mu$ g of B<sub>12</sub>, 51 mg of niacin, 1.5 mg of folic acid, 126 mg of biotin, and 13.5 mg of pantothenic acid

<sup>2</sup>Provided per kg of complete diet: 37.5 mg of Zn, 137.5 mg of Mn, 37.5 mg of Fe, 0.83 mg of I, 0.23 mg of Se, and 1408 mg of choline

2 °C every week down to 24 °C, and 65% relative humidity) and were allowed free access to feed and water during the experiment. Relative breast meat, abdominal fat and organ weights were described as a percentage of live body weight.

**Sampling and measurements.** On Day 0, 7, 21, and 35, chickens and feeders were weighed to allow calculations of body weight gain (BWG), feed

intake (FI) and feed conversion ratio (FCR). Upon completion of the growth assay, 36 chickens (two/pen) were selected randomly and blood samples were collected from a wing vein into K<sub>3</sub>EDTA vacuum tubes (Becton Dickinson Vacutainer Systems, Franklin Lake, NJ). The samples were centrifuged at 3000 × *g* for 15 min to recover plasma. Whole blood cell counts [white blood cells (WBC), red blood cells (RBC), and lymphocytes] were determined using an automatic blood analyser (ADVIA 120, Bayer, NY).

The chickens used for blood samples were weighed and slaughtered so that the breast meat, abdominal fat, gizzard, liver, spleen, bursa of Fabricius, and heart could be removed by trained personnel. All samples were patted dry to remove excess moisture and were weighed. Hunter *L\** (lightness), *a\** (redness), and *b\** (yellowness) of breast meat were measured using a Minolta CR410 chromameter (Konica Minolta Sensing Inc., Osaka, Japan). Drip loss percentage was determined on Day 1, 3, 5, and 7 by using the procedure described by Honikel (1998). Before slaughter six birds from each treatment group (one bird per pen) were chosen randomly and by massaging their abdominal area, excreta samples were collected from the cloacae into microtubes. One gram of the excreta sample

from each pen was diluted with 9 ml of 1% peptone broth (Becton, Dickinson and Co., Franklin Lakes, NJ) and then homogenised. Counts of viable bacteria in excreta samples were determined by plating 10-fold serial dilutions (in 1% peptone solution) onto MacConkey agar plates (Difco Laboratories, Detroit, MI) and *Lactobacilli* medium agar plates (Medium 638, DSMZ, Braunschweig, Germany) to isolate *E. coli* and *Lactobacillus*, respectively. The *Lactobacilli* medium agar plates were then incubated for 48 h at 39 °C under anaerobic conditions. The MacConkey agar plates were incubated for 24 h at 37 °C. The *E. coli* and *Lactobacillus* colonies were counted immediately after removal from the incubator. For *Salmonella*, the serially diluted peptone broth tubes were incubated overnight at 37 °C, after which 1 ml was transferred to 9 ml of tetrastinate broth (Neogen Corporation, Lansing, MI) which was then incubated for 48 h at 42 °C. From these tubes, 1 ml was used to inoculate 9 ml of Rappaport-Vassiliadis *Salmonella* Enrichment broth (Neogen Corporation, Lansing, MI) which was then incubated for 48 h at 42 °C. The Rappaport was used to inoculate XLT4 plates for *Salmonella* isolation, and *Salmonella* were identified using LIS (VIDAS Listeria) and TSI (Triple Sugar Iron) agar tubes (Difco Laboratories, Detroit, MI).

Table 2. Effect of dietary lactulose supplementation on growth performance in broiler chickens

Item	Lactulose (%)			SE	<i>P</i> -value	
	0	0.25	0.5		linear	quadratic
<b>Day 1 to 7</b>						
BWG (g)	101	102	105	3	0.392	0.795
FI (g)	131	133	135	3	0.350	0.936
F : G	1.30	1.30	1.29	0.03	0.968	0.869
<b>Day 7 to 21</b>						
BWG (g)	586	617	638	14	0.018	0.777
FI (g)	894	899	911	15	0.443	0.848
F : G	1.53	1.46	1.43	0.03	0.016	0.515
<b>Day 21 to 35</b>						
BWG (g)	1,028	1,046	1,100	18	0.012	0.439
FI (g)	1,764	1,788	1,800	17	0.164	0.552
F : G	1.72	1.69	1.64	0.03	0.046	0.622
<b>Overall</b>						
BWG (g)	1,715	1,765	1,843	25	0.002	0.664
FI (g)	2,789	2,820	2,846	25	0.134	0.589
F : G	1.63	1.58	1.55	0.02	0.003	0.903

SE = standard error, BWG = gain in BW per bird, FI = feed intake per bird

doi: 10.17221/8722-VETMED

Table 3. Effect of dietary lactulose supplementation on relative weight of breast meat, abdominal fat, and organs in broilers

Item <sup>1</sup> (% of live body weight)	Lactulose (%)			SE	P-value	
	0	0.25	0.5		linear	quadratic
Breast muscle	14.52	17	17.28	0.49	0.001	0.095
Abdominal fat	1.49	1.18	1.21	0.11	0.079	0.217
Gizzard	1.19	1.17	1.17	0.06	0.853	0.881
Heart	0.48	0.49	0.48	0.03	0.967	0.869
Liver	2.09	2.04	2.04	0.17	0.818	0.900
Spleen	0.09	0.07	0.09	0.01	0.922	0.115
Bursa of Fabricius	0.12	0.13	0.13	0.02	0.868	0.924

SE = standard error

**Statistical analysis.** Data were analysed by ANOVA using the GLM procedure of SAS (SAS Institute 1996), with the pen being defined as the experimental unit. The linear and quadratic effect of lactulose among treatments was analysed by using a polynomial regression to describe the shape of the response to increasing concentrations of lactulose in the diet.

response to increasing concentrations of lactulose. From Day 21 to Day 35 there was a linear increase ( $P = 0.012$ ) in BWG and decrease ( $P = 0.046$ ) in FCR, and overall BWG increased ( $P = 0.002$ ) and FCR decreased ( $P = 0.003$ ) linearly in response to increasing concentrations of lactulose in the diets. FI was not affected from Day 0 to 7, Day 7 to 21, Day 21 to 35, or for the overall experimental period.

## RESULTS

### Growth performance

The results of growth performance presented in Table 2 indicate that from Day 1 to Day 7, gain in body weight (BWG), feed intake (FI), and feed conversion ratio (FCR) were not affected by treatments. From Day 7 to Day 21, BWG increased ( $P = 0.018$ ) and FCR decreased ( $P = 0.016$ ) linearly in

### Carcass characteristics

There was a linear ( $P = 0.001$ ) increase in breast muscle weight and a trend for a linear ( $P = 0.079$ ) decrease in abdominal fat by in response to increasing dietary levels of lactulose (Table 3). Relative weights of organs were not affected by treatments. Supplementation of diets with 0.5% lactulose linearly decreased ( $P = 0.002$ ) drip loss percentage on Day 1 and tended (linear,  $P < 0.10$ ) to reduce

Table 4. Effect of dietary lactulose supplementation on colour and drip loss of breast meat

Item	Lactulose (%)			SE	P-value	
	0	0.25	0.5		linear	quadratic
<b>Breast meat colour</b>						
Lightness ( $L^*$ )	57.14	55.40	55.54	1.22	0.370	0.544
Redness ( $a^*$ )	14.84	15.45	15.58	0.53	0.341	0.721
Yellowness ( $b^*$ )	11.28	10.40	11.81	1.01	0.716	0.368
<b>Drip loss (%)</b>						
Day 1	3.98	2.94	2.35	0.30	0.002	0.559
Day 3	6.32	5.59	5.21	0.40	0.070	0.717
Day 5	9.35	8.81	8.10	0.47	0.080	0.873
Day 7	11.08	11.08	10.41	0.51	0.375	0.604

SE = standard error

Table 5. Effect of dietary lactulose supplementation on excreta microbiota in broiler chickens

Item (log <sub>10</sub> cfu/g)	Lactulose (%)			SE	P-value	
	0	0.25	0.5		linear	quadratic
<i>Lactobacillus</i>	7.65	7.77	7.83	0.11	0.263	0.829
<i>E. coli</i>	6.57	6.39	6.40	0.05	0.027	0.143
<i>Salmonella</i>	2.73	2.50	2.47	0.04	0.001	0.106

SE = standard error

drip loss on Day 3. These results demonstrate that addition of lactulose to the diets did not have any significant impact on breast meat colour (Table 4).

### Excreta microbiota

Data presented in Table 5 indicate that there was a linear decrease in the count of excreta *E. coli* ( $P = 0.027$ ) and *Salmonella* ( $P = 0.001$ ). The count of *Lactobacillus* was not affected by experimental diets.

### Blood constituents

Counts of blood cells and haptoglobin concentrations in blood were not affected by treatments but RBC counts increased (linear effect,  $P = 0.164$ ) with increasing lactulose concentrations in the diet.

## DISCUSSION

Several researchers have reported that supplementing diets with prebiotics may improve growth performance and possibly decrease mortality in chickens, pigs, and calves (Fairchild et al. 2001; Hooge et al. 2004; Fleige et al. 2007; Cho and Kim 2014). Our results are in agreement with these previous studies and indicate that supplementing diets with 0.5% lactulose improves the growth performance of broiler chickens.

Previous research demonstrated that prebiotics can modulate the gut environment by increasing the number of beneficial microorganisms and inhibiting the proliferation of pathogens in the intestine (Patterson and Burkholder 2003; Higgins et al. 2008). Prebiotics in the diet reduced the population of *Salmonella* in the intestine of chickens (Bailey et al. 1991; Pascual et al. 1999; Stern et al. 2001) and prebiotics supported competitive exclusion and immune modulation (Jin et al. 1997; Simon et al. 2001). Our observations that feeding broiler chickens diets supplemented with lactulose increased excreta *Lactobacilli* whereas the counts of *E. coli* and *Salmonella* were reduced support the argument that lactulose acts as a prebiotic in chickens and this may partly explain the improved growth performance observed for the birds fed treatment diets.

Heckert et al. (2002) suggested measuring the weight of immune organs as a method for estimating the immune status of chickens. The bursa of Fabricius and spleen are the main lymphoid organs in broilers but these authors concluded that inclusion of prebiotics did not increase the relative weight of these organs. In the current study the lymphocyte counts were not influenced by dietary treatment, which is consistent with the lack of an effect on the relative weights of the spleen and bursa of Fabricius. These findings are consistent with the results reported by Mohebbifar et al. (2013).

In conclusion, supplementing diets with lactulose improved the growth performance and altered the

Table 6. Effect of dietary lactulose supplementation on blood constituents in broilers

Item	Lactulose (%)			SE	P-value	
	0	0.25	0.5		linear	quadratic
WBC (10 <sup>3</sup> /μl)	451.7	410.5	536.3	82.5	0.479	0.421
RBC (10 <sup>6</sup> /μl)	2.29	2.73	2.82	0.3	0.164	0.572
Lymphocyte (%)	77.4	57.3	68.5	13.8	0.656	0.371
Haptoglobin (mg/dl)	16	17	18	0.9	0.094	0.565

SE = standard error, WBC = white blood cells, RBC = red blood cells

doi: 10.17221/8722-VETMED

excreta microbiota by increasing *Lactobacilli* and decreasing *E. coli* and *Salmonella* counts. However, breast meat colour, relative weights of organs and blood constituents were not affected.

## REFERENCES

- Bailey JS, Blankenship LC, Cox NA (1991): Effect of fructooligosaccharide on *Salmonella* colonization of the chicken intestine. *Poultry Science* 70, 2433–2438.
- Ballongue J, Schumann C, Quignon P (1997): Effects of lactulose and lactitol on colonic microflora and enzymatic activity. *Scandinavian Journal of Gastroenterology* 32, 41–44.
- Bauer E, Williams BA, Verstegen MWA, Mosenthin R (2006): Fermentable carbohydrates: potential dietary modulators of intestinal physiology, microbiology and immunity in pigs. In: Mosenthin R, Zentek J, Zebroska T (eds.): *Biology of Growing Animals Series: Biology of Nutrition in Growing Animals*. Elsevier Limited, Edinburgh, United Kingdom. 4, 33–63.
- Bianchi G, Ronchi M, Marchedini G (1997): Effects of lactulose on carbohydrate metabolism and diabetes mellitus. *Scandinavian Journal of Gastroenterology* 32, 62–64.
- Bird SP, Hewitt D, Ratcliffe B, Gurr MI (1990): Effects of lactulose and lactitol on protein digestion and metabolism in conventional and germ free animal models: relevance of the results to their use in the treatment of portosystemic encephalopathy. *Gut* 31, 1403–1406.
- Cho JH, Kim IH (2014): Effects of lactulose supplementation on performance, blood profiles, excreta microbial shedding of *Lactobacillus* and *Escherichia coli*, relative organ weight and excreta noxious gas contents in broilers. *Journal of Animal Physiology and Animal Nutrition* 98, 424–430.
- Cummings JH, McFarlane GT (2002): Gastrointestinal effects of prebiotics. *British Journal of Nutrition* 87, 145–151.
- Fairchild AS, Grimes JL, Jones FT, Wineland MJ, Edens FW, Sefton AE (2001): Effects of hen age, Bio-Mos and flavomycin on poult susceptibility to oral *Escherichia coli* challenge. *Poultry Science* 80, 562–571.
- Fleige S, Preibinger W, Meyer HHD, Pfaffl MW (2007): Effect of lactulose on growth performance and intestinal morphology of pre-ruminant calves using a milk replacer containing *Enterococcus faecium*. *Animal* 1, 367–373.
- Fuller R (2001): The chicken gut microflora and probiotic supplements. *Journal of Poultry Science* 38, 189–196.
- Gibson GR (2004): From probiotics to prebiotics and a healthy digestive system. *Journal of Food Science* 69, 141–143.
- Heckert RA, Estevez I, Russek-Cohen E, Pettie-Riley R (2002): Effects of density and perch availability on the immune status of broilers. *Poultry Science* 81, 451–457.
- Higgins SE, Higgins JP, Wolfenden AD, Henderson SN, Torres-Rodriguez A, Tellez Hargis GB (2008): Evaluation of a lactobacillus-based probiotic culture for the reduction of *Salmonella enteritidis* in neonatal broiler chicks. *Poultry Research* 87, 27–31.
- Honikel KO (1998): Reference methods for the assessment of physical characteristics of meat. *Meat Science* 49, 447–457.
- Hooge DM, Ishimaru H, Sims MD (2004): Influence of dietary *Bacillus subtilis* C-3102 spores on live performance of broiler chickens in four controlled pen trials. *Journal of Applied Poultry Research* 13, 222–228.
- Huang MK, Choi JL, Houde R, Lee JW, Lee B, Zhao X (2004): Effects of *Lactobacilli* and an acidophilic fungus on the production performance and immune responses in broiler chickens. *Poultry Science* 83, 788–795.
- Jin LZ, Ho YW, Abdullah N, Jalaludin S (1997): Probiotics in poultry: Modes of action. *World's Poultry Science Journal* 53, 351–368.
- Kermanshahi H, Rostami H (2006): Influence of supplemental dried whey on broiler performance and cecal flora. *International Journal of Poultry Science* 5, 538–543.
- Kleesen B, Hartmann C, Blaut N (2001): Oligofructose and long-chain inulin: influence on the gut microbial ecology of rats associated with human faecal flora. *British Journal of Nutrition* 80, 1–11.
- Krueger M, Schroedl W, Isik K, Lange W, Hagemann L (2002): Effects of lactulose on the intestinal microflora of periparturient sows and their piglets. *European Journal of Nutrition* 41, 26–31.
- Marinho MC, Lordelo MM, Cunha LF, Freire JPB (2007): Microbial activity in the gut of piglets: I. Effect of prebiotic and probiotic supplementation. *Livestock Science* 108, 236–239.
- Markovic R, Sefera D, Krstic M, Petrujkic B (2009): Effect of different growth promoters on broiler performance and gut morphology. *Archivos de Medicina Veterinaria* 41, 163–169.
- Mitsuoka T, Hidaka H, Eida T (1987): Effect of fructo-oligosaccharides on intestinal microflora. *Nahrung* 31, 426–436.
- Mohebbifar A, Kashani H, Afsari M, Torki M (2013): Effects of commercial prebiotics and probiotics on performance of laying hens, egg traits and some blood parameters. *Annual Review and Research in Biology* 3, 921–934.
- NCR – National Research Council (1994): *Nutrient Requirements of Poultry*. 9<sup>th</sup> rev. ed. Natl. Acad. Press, Washington, DC.
- Pascual M, Hugas M, Badiola JI, Monfort JM, Garriga M (1999): *Lactobacillus salivarius* CTC2197 prevents *Salmonella enteritidis* colonization in chickens. *Applied Environmental Microbiology* 65, 4981–4986.

- Patterson JA, Burkholder KM (2003): Application of prebiotics and probiotics in poultry production. *Poultry Science* 82, 627–631.
- SAS Institute (1996): SAS User's Guide: Statistics. Version 7.0. SAS Institute Inc., Cary, NC.
- Simon O, Jadamus A, Vahjen W (2001): Probiotic feed additives – effectiveness and expected modes of action. *Journal of Animal Feed Science* 10, 51–67.
- Steiner T (2006): Managing gut health: Natural growth promoters as a key to animal performance. Nottingham University Press, Nottingham, UK.
- Stern NJ, Cox NA, Bailey JS, Berrang ME, Musgrove MT (2001): Comparison of mucosal competitive exclusion and competitive exclusion treatment to reduce *Salmonella* and *Campylobacter* spp. colonization in broiler chickens. *Poultry Science* 80, 156–160.
- Zhang ZF, Kim IH (2013): Effects of probiotic supplementation in different energy and nutrient density diets on performance, egg quality, excreta microflora, excreta noxious gas emission, and serum cholesterol concentrations in laying hens. *Journal of Animal Science* 91, 4781–4787.
- Zhang, ZF, Cho JH, Kim IH (2013): Effects of *Bacillus subtilis* UBT-MO<sub>2</sub> on growth performance, relative immune organ weight, gas concentration in excreta, and intestinal microbial shedding in broiler chickens. *Livestock Science* 155, 343–347.
- Zhao PY, Wang JP, Kim IH (2013): Evaluation of dietary fructan supplementation on growth performance, nutrient digestibility, meat quality, fecal microbial flora, and fecal noxious gas emission in finishing pigs. *Journal of Animal Science* 91, 5280–5286.

Received: 2014–11–18

Accepted after corrections: 2016–01–14

---

**Corresponding Author:**

In Ho Kim, Dankook University, Department of Animal Resource and Science, Cheonan, Choongnam, 330-714, Republic of Korea  
E-mail: inhokim@dankook.ac.kr

---