

<https://doi.org/10.17221/181/2017-CJGPB>

Genetic effects of F₁ pollen sterility genes *S-b*, *S-d* and *S-e* in rice (*Oryza sativa* L.)

MINGSONG JIANG, JIANDI XU, FENG CHEN, WENYIN ZHU*

Rice Research Institute, Jinan, Shandong, P.R. China

*Corresponding author: wyzhu0321@163.com

Citation: Jiang M., Xu J., Chen F., Zhu W. (2019): Genetic effects of F₁ pollen sterility genes *S-b*, *S-d* and *S-e* in rice (*Oryza sativa* L.). Czech J. Genet. Plant Breed., 55: 55–60.

Abstract: An experimental population commonly used in genetic analyses of gene or quantitative trait loci (QTLs) in rice is chromosome segment substitution lines (CSSLs). In the present study, with the typical *indica* variety Guangluai 4 as a donor and *japonica* variety Taichung 65 as a recipient, seven CSSLs carrying F₁ pollen sterility genes *S-b*, *S-d*, *S-e*, *S-b/S-d*, *S-b/S-e*, *S-d/S-e*, and *S-b/S-d/S-e* were obtained by specific selection for the target genes, non-specific selection for the genome of the recurrent parents in four backcross populations (BC₁F₂, BC₂F₂, BC₃F₂ and BC₃F₃). We evaluated the genetic effect of the F₁ pollen sterility genes using 35 F₁ hybrid individuals in crosses derived from CSSLs and Taichung 65. Pollen fertility of F₁ hybrid plants was observed and the results indicated that the single genes *S-b*, *S-d* and *S-e* can cause 67.7%, 14.6% and 53.2% of pollen sterility, respectively. Multiple genes *S-b/S-d*, *S-b/S-e*, *S-d/S-e*, and *S-b/S-d/S-e* can cause 76.6%, 85%, 68.7%, and 93% of pollen sterility, respectively.

Keywords: chromosome segment substitution lines; donor; marker-assisted selection; recipient

Rice (*Oryza sativa* L.) is one of the most important crops feeding about half of the world's population. At present, with a reduction of the global cultivated rice land area, breeding more productive rice varieties becomes an urgent task for rice breeders (LI *et al.* 2006; ALI *et al.* 2010). F₁ hybrids between *indica* and *japonica* subspecies usually demonstrate very strong hybrid vigour or heterosis, which has attracted considerable research interest with the hope of using such heterosis in hybrid rice production (QIU *et al.* 2005). However, varying degrees of hybrid sterility are commonly seen in crosses between *indica* and *japonica* subspecies, and the partial or complete sterility of the hybrids is a serious constraint for utilizing heterosis in hybrid rice breeding (OKA 1957; LIU *et al.* 1997; LONG *et al.* 2008).

Apart from female gamete abortion, male gamete abortion (F₁ pollen sterility) is considered to play a key role in *indica-japonica* hybrid sterility (OKA 1974; JING *et al.* 2007). ZHANG and LU (1989, 1993) put forward the "Specific Compatibility Hypothesis".

According to the hypothesis, F₁ pollen sterility was the main form of hybrid sterility and there were at least six genes, namely *S-a*, *S-b*, *S-c*, *S-d*, *S-e* and *S-f*, controlling the pollen sterility of F₁ hybrids. The degree of pollen sterility caused by the allelic interaction at the loci varied with the particular gene and the gene number: the higher the level of heterozygosity at the six genes, the higher pollen and spikelet sterility is observed. Up to now, five loci, *S-a*, *S-b*, *S-c*, *S-d* and *S-e*, have been located in the genetic map of rice (ZHANG & ZHANG 2001; LI *et al.* 2002; ZHUANG *et al.* 2002; YANG *et al.* 2004; LI *et al.* 2006; ZHU *et al.* 2008). Among them, three loci, *S-b*, *S-c* and *S-d*, have already been located in the physical map, and one locus, *S-a*, has been cloned by positional cloning strategy (LONG *et al.* 2008).

Chromosome segment substitution lines (CSSLs) are important genetic stocks for investigating the function and regulation of single genes and useful for gene cloning. In rice, several CSSLs have been produced (ZHU *et al.* 2009; RAMOS *et al.* 2016). The

goals of the present study were (1) construction of CSSLs containing F_1 pollen sterility genes *S-b*, *S-d* and *S-e*, (2) validation of these genes using the CSSLs, and (3) estimation of the gene effect on the hybrid sterility trait.

MATERIAL AND METHODS

Plant materials and growth conditions. Guangluai 4 used as a donor in this study is a typical *indica* variety in South China's Guangdong province. Taichung 65 is a typical *japonica* variety, which was used as a recipient. First, the cross between Guangluai 4 and Taichung 65 was carried out, then a series of backcrosses with Taichung 65 as the recurrent parent was performed. A total of 240 BC_3F_3 individual plants as well as the parents and 35 F_1 individuals (CSSLs/Taichung 65) were planted in the experimental farm of Nanyanghu, Jining city, Shandong province of China.

DNA preparation and PCR protocol. DNA was extracted from fresh young leaves using the CTAB method (MURRAY & THOMPSON 1980). The PCR was conducted with minor modifications by PANAUD *et al.* (1996). The protocol for PCR was as follows: the template DNA was subjected to denaturation at 94°C for 5 min, followed by 35 cycles (denaturation at 94°C for 1 min, primer annealing at 55°C for 1 min, and primer extension at 72°C for 1 min), and a final extension at 72°C for 5 min. The PCR products were separated through electrophoresis on a 6% non-denaturing polyacrylamide gel. Bands were revealed using a silver staining procedure.

Genetic background detection and molecular marker linked with *S-b*, *S-d*, and *S-e*. For genetic

background detection, 180 SSR markers selected from dense rice microsatellite maps (McCOUCH *et al.* 2002) were used to survey the polymorphism between Guangluai 4 and Taichung 65. Polymorphic markers were used for the genetic background detection of CSSLs. For selection of CSSLs carrying F_1 pollen sterility genes, six markers were used for *S-b*, *S-d*, and *S-e* (Table 1). Bulk segregant analysis (BSA) was employed to evaluate the uniformity of the genetic background in the CSSLs.

Pollen fertility analysis. The identification of pollen fertility was assessed as described by ZHANG and LU (1989). Ten florets per panicle were collected from the upper one-third portion of the panicle and fixed in a FAA solution of 6% (v/v) formaldehyde, 89% (v/v) alcohol, and 5% (v/v) acetic acid. The pollen phenotypes of rice were observed in a 1% I₂-KI (potassium iodide) solution under a microscope. Pollen was classified as fertile (round and full dark), empty abortive (irregular and yellow) and staining abortive (round and partial dark) based on their shape and stainability. Pollen fertility of hybrids was observed in F_1 hybrid plants and different types of individuals from F_2 populations.

RESULTS

Construction of CSSLs. The procedure for the development of CSSLs is summarized in Figure 1. The F_1 plants derived from Guangluai 4/Taichung 65 were selfed to produce an F_2 generation. In the F_2 population, individuals similar to Taichung 65 in plant height and heading date were selected to backcross with Taichung 65 to produce BC_1F_2 population con-

Table 1. The primers linked with rice F_1 pollen sterility genes *S-b*, *S-d* and *S-e*

F_1 pollen sterility locus	Chromosome	Linked markers	Sequence of marker (5'-3')
<i>S-b</i>	5	PSM60	F: CCCATCCAGGTCACCACCACAAT R: AATCCGAATCGCATCAGAAGCAG
		PSM202	F: CGGAATCAATGGAAGGTTT R: CCCTTGACTTCCCATTCT
<i>S-d</i>	1	PSM43	F: CGTAGTGGTCCATCGGAGGC R: TGAGCTGAGCTGCGGCAAG
		RM84	F: TAAGGGTCCATCCACAAGATG R: TTGCAAATGCAGCTAGAGTAC
<i>S-e</i>	12	PSM597	F: GCCTGATGCTACTCGCCAC R: CAGCCATTGTTAATTCTACGATG
		PSM459	F: CTGCATTGCGCTTGGTGT R: GATCCTGAAGTGTTGTGGC

<https://doi.org/10.17221/181/2017-CJGPB>




Figure 1. The development of seven chromosomal segment substitution lines with Guangluai 4 as donor and Taichung 65 as recipient; marker-assisted selection (MAS) was used for the selection of F_1 pollen sterility genes *S-b*, *S-d*, and *S-e*

taining 240 individuals. For each backcross only individuals carrying the genes *S-b*, *S-d* and *S-e* derived from Guangluai 4 were used. With marker-assisted selection (MAS), 48 BC_1F_2 plants were backcrossed with Taichung 65 to produce 305 BC_2F_2 plants. In the same way, 245 BC_3F_2 individuals derived from 20 BC_2F_2 plants were obtained. MAS with a whole-genome survey of 245 BC_3F_2 enabled the identification of 13 plants homozygous for Taichung 65 in the majority of genomic regions except for one or two heterozygous substituted segments from the donor with sterility genes. These plants were selfed to produce 13 BC_3F_3 lines (650 individuals). Finally, a total of seven CSSLs carrying one or multiple F_1 pollen sterility genes (*S-b*, *S-d*, *S-e*, *S-b/S-d*, *S-b/S-e*, *S-d/S-e*, and *S-b/S-d/S-e*) were obtained.

Detection of residual segments and uniformity of genetic background in the CSSLs. One hundred and

Table 2. Residual segments in the BC_3F_2 population of rice

Chromosome	Marker	Genotype of six BSAs
2	RM109	heterozygote
	RM112	heterozygote
	RM154	heterozygote
3	RM85	heterozygote
	RM218	heterozygote
	RM442	heterozygote
4	RM131	heterozygote
7	RM298	heterozygote

BSA – bulked segregant analysis

twenty-nine SSR markers displayed polymorphism between parents and the rate of polymorphism was 71.67%. To evaluate the uniformity of the genetic background in the seven CSSLs, a round of examination for the genetic background was carried out and Bulk Segregant Analysis (BSA) was employed with polymorphic markers. Six BSAs (245 BC_3F_2 individuals) were tested using 129 polymorphic SSR markers selected in non-target chromosomal regions. From 774 PCR products, residual segments or heterozygous segments in non-target chromosomal regions were detected in 48 PCR products. The percentage of PCR products of residual segments was 6.2% in the BC_3F_2 population (Table 2). All residual segments in the population were treated as non-target segments for further selection in 245 BC_3F_2 individuals. The second round of detection for donor residual segments was done in the BC_3F_3 generation. From 1600 PCR products in 10 candidate CSSLs, 20 PCR products were found to reveal donor alleles. The percentage of PCR products of residual segments was 1.3%. The lines with residual segments were eliminated. These results implied that the homozygosity of Taichung 65 alleles in non-target chromosomal regions or the uniformity of the Taichung 65 genetic background in the seven CSSLs was greater than 98.8%.

Genetic effects of F_1 pollen sterility genes *S-b*, *S-d* and *S-e*. The results indicated that the single genes *S-b*, *S-d* and *S-e* can cause 67.7%, 14.6% and 53.2% of pollen sterility (Figure 2), respectively. While two genes *S-b/S-d*, *S-b/S-e* and *S-d/S-e* can cause 76.6%, 85% and 68.7% of pollen sterility (Figure 3),




Figure 2. Genetic effects of single rice F_1 pollen sterility genes *S-b*, *S-d*, and *S-e* evaluated in three F_1 hybrids derived from CSSL-1/Taichung 65, CSSL-2/Taichung 65 and CSSL-3/Taichung 65




Figure 3. Genetic effects of double rice F₁ pollen sterility genes *S-b/S-d*, *S-b/S-e*, and *S-d/S-e* evaluated in three F₁ hybrids and six different types of F₂ individuals derived from CSSL-4/Taichuang 65, CSSL-5/Taichung 65 and CSSL-6/Taichung 65




Figure 4. Genetic effects of multiple rice F₁ pollen sterility genes *S-b/S-d/S-e* evaluated in F₁ hybrid plants and six different types of F₂ individuals derived from CSSL-7/Taichung 65

respectively, and multiple genes *S-b/S-d/S-e* can cause 93% of pollen sterility (Figure 4).

DISCUSSION

The hybrid sterility between *indica* and *japonica* subspecies of Asian cultivated rice is a complex trait. The genetic basis of F₁ partial sterility has been intensively studied. An allelic interaction model may

explain the mechanism of hybrid sterility (IKEHASHI & ARAKI 1986, 1988). According to the model, there were three alleles at the *S5* gene, *S5ⁱ*, *S5^j* and *S5ⁿ*, representing *indica*, *japonica* and a widely compatible variety, respectively (CHEN *et al.* 2008). The heterozygote of *indica/japonica* (*S5ⁱ/S5^j*) produced semi-sterile panicles, resulting from the partial abortion of female gametes carrying *S5^j*, whereas the heterozygote consisting of the *S5ⁿ* allele with either of the other two alleles, e.g. *S5ⁿ/S5ⁱ* or *S5ⁿ/S5^j*, would be fully fertile. So far, a series of female sterility genes including major QTLs was identified and mapped (WAN *et al.* 1993, 1996; WAN & IKEHASHI 1995; WANG *et al.* 1998; LIU *et al.* 2001; SONG *et al.* 2005).

It seemed that two approaches may be able to overcome F₁ sterility between *indica-japonica* subspecies varieties. The one was to screen widely compatible varieties. A number of these varieties produced fertile F₁ hybrids when crossed to both *indica* and *japonica* varieties and thus they were called widely compatible varieties. IKEHASHI and ARAKI (1984) recognized the significance of this phenomenon and then proposed a genetic basis for F₁ sterility and wide compatibility. The finding of the neutral allele *S5ⁿ* excited rice breeders to screen widely compatible varieties from a large amount of *indica* and *japonica* subspecies through extensive test-crosses and some were gathered. The widely compatible varieties were currently the main tools for breaking the sterility barriers between *indica* and *japonica* subspecies and have made an important contribution to rice breeding. Recently, however, some *indica-japonica* hybrids

<https://doi.org/10.17221/181/2017-CJGPB>

which had the neutral allele $S5^n$ have been tested for their yield potential. Such hybrids showed strong heterosis, but their seed set was unstable in adverse conditions. In fact, besides $S5^n$, there are neutral alleles of other genes. It seemed that more neutral alleles should be pyramided to make varieties widely compatible. The other was breeding and utilization of *indica*-compatible *japonica* lines by introducing the S^i allele into *japonica* varieties through continuous backcrosses. *Indica*-compatible *japonica* lines carried S^i/S^i at the hybrid sterility loci. When crossed with *indica* varieties, the F_1 not only had a strong vigour, but also was fully fertile. Such a kind of breeding program is being undertaken and some excellent *indica*-compatible *japonica* lines have already been cultivated (DING *et al.* 2002; GUO *et al.* 2016). This approach needs to pyramid F_1 pollen sterility genes which carry S^i/S^i to the same *japonica* variety. Hence, searching for F_1 pollen sterility genes and evaluating their genetic effect are very important for developing *indica*-compatible *japonica* lines. The development of the CSSLs of F_1 pollen sterility genes not only enriched and perfected the ‘Specific Compatibility Hypothesis’ but also provided the strong basis for breeding *indica*-compatible *japonica* lines with molecular marker-assisted selection.

References

- Ali M., Paul L., Yu S., Lorieux M., Eizenga G. (2010): Chromosome segment substitution lines: A powerful tool for the introgression of valuable genes from *Oryza* wild species into cultivated rice (*O. sativa*). *Rice*, 3: 218–234.
- Chen J., Ding J., Ouyang Y., Du H., Yang J., Cheng K., Zhan J., Qiu S., Zhang X., Yao J., Liu K., Wang L., Xu C., Li X., Xue Y., Xia M., Lu J., Xi M., Zhang Q. (2008): A triallelic system of $S5$ is a major regulator of the reproductive barrier and compatibility of *indica-japonica* hybrids in rice. *Proceeding of the National Academy of Sciences of the USA*, 105: 11436–11441.
- Ding X., Chen Y., Yang C., Lu Y., Zhang G. (2002): Test on compatibility of *indica*-compatible *japonica* lines in rice. *Chinese Journal of Rice Science*, 16: 366–368. (in Chinese with English abstract)
- Guo J., Xu X., Li W., Zhu W., Zhu H., Liu Z., Luan X., Dai Z., Liu G., Zhang Z., Zeng R., Tang G., Fu X., Wang S., Zhang G. (2016): Overcoming inter-subspecific hybrid sterility in rice by developing *indica*-compatible *japonica* lines. *Scientific Reports*, 6: 26878.
- Ikehashi H., Araki H. (1984): Variety screening of compatibility types revealed in F_1 fertility of distant crosses in rice. *Japanese Journal of Breeding*, 34: 304–313.
- Ikehashi H., Araki H. (1986): Genetics of F_1 sterility in remote cross of rice (*Oryza sativa* L.). In: Banta J.S. (ed.): *Rice Genetics I*, Part 2. IRRI: 119–130.
- Ikehashi H., Araki H. (1988): Multiple alleles controlling F_1 sterility in remote crosses of rice (*Oryza sativa*). *Japanese Journal of Breeding*, 38: 283–291.
- Jing W., Zhang W., Jiang L., Chen L., Zhai H., Wan J. (2007): Two novel loci for pollen sterility in hybrids between the weedy strain Ludao and the *japonica* variety Akihikari of rice (*Oryza sativa* L.). *Theoretical and Applied Genetics*, 114: 915–925.
- Li W., Zeng R., Zhang Z., Zhang G. (2002): Mapping of $S-b$ locus for F_1 pollen sterility in cultivated rice using PCR based markers. *Acta Botanica Sinica*, 44: 463–467.
- Li W., Zeng R., Zhang Z., Ding X., Zhang G. (2006): Fine mapping of locus $S-b$ for F_1 pollen sterility in rice (*Oryza sativa* L.). *Chinese Science Bulletin*, 51: 675–680.
- Liu K., Wang J., Li H., Xu C., Liu A., Li X., Zhang Q. (1997): A genome-wide analysis of wide compatibility in rice and the precise location of the $S5$ locus in the molecular map. *Theoretical and Applied Genetics*, 95: 809–814.
- Liu Y., Zhu L., Sun J., Chen Y. (2001): Mapping QTLs for defective female gametophyte development in an inter-subspecific cross in *Oryza sativa*. *Theoretical and Applied Genetics*, 102: 1243–1251.
- Long Y., Zhao L., Niu B., Su J., Wu H., Chen Y., Zhang Q., Guo J., Zhuang C., Mei M., Xia J., Wang L., Wu H., Liu Y. (2008): Hybrid male sterility in rice controlled by interaction between divergent alleles of two adjacent genes. *Proceeding of the National Academy of Sciences of the USA*, 105: 18871–18876.
- McCouch S., Teytelman L., Xu Y., Lobos K., Clare K., Walton M., Fu B., Maghirang R., Li Z., Xing Y., Zhang Q., Kono I., Yano M., Fjellstrom R., DeClerck G., Schneider D., Cartinhour S., Ware D., Stein L. (2002): Development and mapping of 2240 new SSR markers for rice (*Oryza sativa* L.). *DNA Research*, 9: 199–207.
- Murray M., Thompson W. (1980): Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Research*, 8: 4321–4325.
- Oka H. (1957): Genetic analysis for the sterility of hybrids between distantly related varieties of cultivated rice. *Journal of Genetics*, 53: 397–409.
- Oka H. (1974): Analysis of genes controlling F_1 sterility in rice by the use of isogenic lines. *Genetics*, 77: 521–534.
- Panaud O., Chen X., McCouch S. (1996): Development of microsatellite markers and characterization of simple sequence length polymorphism (SSLP) in rice (*Oryza sativa* L.). *Molecular Genetics and Genomics*, 252: 597–600.
- Qiu S., Liu K., Jiang J., Song X., Xu C., Li X., Zhang Q. (2005): Delimitation of the rice wide compatibility gene

<https://doi.org/10.17221/181/2017-CJGPB>

- S5(n)* to a 40-kb DNA fragment. Theoretical and Applied Genetics, 111: 1080–1086.
- Ramos J.M., Furuta T., Uehara K., Chihiro N., Angeles-Shim R.B., Shim J., Brar D.S., Ashikari M., Jena K.K. (2016): Development of chromosome segment substitution lines (CSSLs) of *Oryza longistaminata* A. Chev. & Röhr in the background of the elite *japonica* rice cultivar, Taichung 65 and their evaluation for yield traits. Euphytica, 210: 151–163.
- Song X., Qiu S., Xu C., Li X., Zhang Q. (2005): Genetic dissection of embryo sac fertility, pollen fertility, and their contributions to spikelet fertility of intersubspecific hybrids in rice. Theoretical and Applied Genetics, 110: 205–211.
- Wan J., Ikehashi H. (1995): Identification of a new locus *S-16* causing hybrid sterility in native rice varieties (*Oryza sativa* L.) from Taihu lake region and Yunnan province. Breeding Science, 45: 461–470.
- Wan J., Yanagihara S., Kato H., Ikehashi H. (1993): Multiple alleles at a new locus causing hybrid sterility between a Korean *indica* variety and a *japonica* variety in rice. Japanese Journal of Breeding, 43: 507–516.
- Wan J., Yamaguchi Y., Kato H., Ikehashi H. (1996): Two new loci for hybrid sterility in cultivated rice (*Oryza sativa* L.). Theoretical and Applied Genetics, 92: 183–190.
- Wang J., Liu K., Xu C., Li X., Zhang Q. (1998): The high level of wide-compatibility of variety 'Dular' has a complex genetic basis. Theoretical and Applied Genetics, 97: 407–412.
- Yang C., Chen Z., Zhuang C., Mei M., Liu Y. (2004): Genetic and physical fine-mapping of the *S-c* locus conferring *indica-japonica* hybrid sterility in rice (*Oryza sativa* L.). Chinese Science Bulletin, 49: 1718–1721.
- Zhang G., Lu Y. (1989): Genetic studies of the hybrid sterility in cultivated rice (*Oryza sativa*). I. Diallel analysis of the hybrid sterility among isogenic F_1 sterile lines. Chinese Journal of Rice Science, 3: 97–101. (in Chinese with English abstract)
- Zhang G., Lu Y. (1993): Genetic studies of the hybrid sterility in cultivated rice (*Oryza sativa*). II. A genic model for F_1 pollen sterility. Acta Genetica Sinica, 20: 222–228. (in Chinese with English abstract)
- Zhang Z., Zhang G. (2001): Fine mapping of the *S-c* locus and marker-assisted selection using PCR markers in rice. Acta Agronomica Sinica, 27: 704–709. (in Chinese with English abstract)
- Zhu W., Li., Ding X., Zhang Z., Zeng R., Zhu H., Zhang G. (2008): Preliminary identification of F_1 pollen sterility gene *S-e* in *Oryza sativa*. Journal of South China Agricultural University, 29: 1–5. (in Chinese with English abstract)
- Zhu W., Lin J., Yang D., Zhao L., Zhang Y., Zhu Z., Chen T., Wang C. (2009): Development of chromosome segment substitution lines derived from backcross between two sequenced rice cultivars, *indica* recipient 93-11 and *japonica* donor Nipponbare. Plant Molecular Biology Reporter, 27: 126–131.
- Zhuang C., Fu Y., Zhang G., Mei M., Lu Y. (2002): Molecular mapping of *S-c*, an F_1 pollen sterility gene in cultivated rice. Euphytica, 127: 133–138.

Received for publication November 21, 2017

Accepted after corrections June 29, 2018

Published online December 19, 2018