

<https://doi.org/10.17221/187/2018-CJAS>

## Antioxidant defence system in protein fractions of common carp (*Cyprinus carpio*) seminal plasma

ANNA SHALIUTINA-KOLEŠOVÁ<sup>1,2\*</sup>, MO XIAN<sup>1</sup>, RUI NIAN<sup>1\*</sup>

<sup>1</sup>CAS Key Laboratory of Biobased Materials, Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao, P.R. China

<sup>2</sup>Research Institute of Fish Culture and Hydrobiology, South Bohemian Research Center of Aquaculture and Biodiversity of Hydrocenoses, Faculty of Fisheries and Protection of Waters, University of South Bohemia in České Budějovice, Vodňany, Czech Republic

\*Corresponding authors: [kolesova@frov.jcu.cz](mailto:kolesova@frov.jcu.cz), [nianrui@qibebt.ac.cn](mailto:nianrui@qibebt.ac.cn)

**Citation:** Shaliutina-Kolešová A., Xian M., Nian R. (2019): Antioxidant defence system in protein fractions of common carp (*Cyprinus carpio*) seminal plasma. Czech J. Anim. Sci., 64, 265–271.

**Abstract:** There are growing evidences that the damage which is caused to the spermatozoa by the reactive oxygen species (ROS) plays a key role in male infertility. The seminal plasma is endowed with many enzymatic and non-enzymatic antioxidants which protect the spermatozoa against oxidative stress. The present study was undertaken to compare the level of superoxide dismutase (SOD), glutathione reductase (GR) and glutathione peroxidase (GPx) in separated protein fractions of whole common carp *Cyprinus carpio* seminal plasma. Seminal plasma from six male common carp was separated into four fractions. The fractions differed from each other in their total seminal plasma protein content and antioxidant status. The highest value of SOD ( $2.81 \pm 0.11$  mU/mg protein) was found in fraction 4, and the lowest ( $1.83 \pm 0.18$  mU/mg protein) in fraction 1. There were significant variations (ANOVA;  $P = 0.05$ ) in GR activity, mainly between fractions 1, 2, and 4. The lowest value of GPx activity ( $14.9 \pm 0.4$  mU/mg protein) was found in fraction 2; however, the GPx activity showed low variability among fractions. We conclude that distinct seminal plasma fractions differ in antioxidant capacity. The inclusion of fractions with high levels of SOD, GR, and GPx in sperm storage medium may enhance protection against oxidative stress and promote sperm survival during the *in vitro* storage.

**Keywords:** fish; seminal plasma fractions; proteins; antioxidant activity

Seminal plasma (SP) is a natural medium containing components that provide an optimal environment for the storage of spermatozoa in the sperm duct (Ciereszko et al. 2011). The role of seminal plasma was traditionally assumed to be primarily a medium for sperm transport; however, its isolated constituents demonstrate direct effects

on sperm physiology and fertility (Maxwell et al. 2007; Ratto et al. 2011), which differ with species.

The use of specific seminal plasma fractions, rather than the entire supernatant, for preservation of sperm *in vitro* has been recently studied with promising results (Saravia et al. 2007). Barrios et al. (2000) demonstrated by scanning electron

Supported by the Ministry of Education, Youth and Sports of the Czech Republic (projects CENAKVA (LM2018099) and Biodiversity (CZ.02.1.01/0.0/0.0/16\_025/0007370)), by the CAS President's International Fellowship for Postdoctoral Researchers (2017PB0060), National Natural Science Foundation of China (21676286), and Primary Research Development Plan of Shandong Province (2016GSF121006).

microscopy that fractions of ram seminal plasma can restore the membrane of spermatozoa damaged by cold shock. The study also showed that *in vitro* incubation of ram sperm in selected fractions of seminal plasma increased the rate of spermatozoon viability. In studies conducted in boar, spermatozoa incubated with SP pre-sperm-rich fraction, sperm-rich fraction, and post-sperm rich fractions, showed the highest percentage motility, with the highest spermatozoon viability recorded after exposure to the sperm-rich fraction (Garcia et al. 2009). Research suggests that seminal plasma fractions show potential as additives for preserving the function of semen; however, their effects have been little studied in fish and need investigation.

Oxidative stress is the result of an imbalance between the production of reactive oxygen species (ROS) and their neutralisation or scavenging by the antioxidant system (Fazeli and Salimi 2016). When the production of ROS by spermatozoon mitochondria is excessive, the gamete's limited defences are rapidly overwhelmed, and oxidative damage can induce lipid peroxidation with a loss of vitality and fertilizing potential (Aitken et al. 1998). Unfortunately, fish spermatozoa possess limited endogenous antioxidant protection and are vulnerable to ROS damage (Koppers et al. 2010; Gazo et al. 2013). Therefore, seminal plasma that is well supplied with an array of antioxidants acting as free radical scavengers can protect the spermatozoa against oxidative stress.

Seminal plasma contains a number of enzymatic antioxidants including superoxide dismutase (SOD), glutathione reductase (GR), catalase (CAT), and glutathione peroxidase (GPx) (Li et al. 2010; Martinez-Paramo et al. 2012). In addition, it possesses a variety of non-enzyme antioxidants that suppress the formation of ROS or counteract their actions (Lahnsteiner et al. 2010; Ciereszko et al. 2011). The antioxidant power of a biological fluid such as seminal plasma can be assessed only through the analysis of individual antioxidants.

The aim of this study was to analyse the antioxidant defence system of common carp *Cyprinus carpio* seminal plasma by measuring SOD, GR, and GPx activity in its separated fractions and to compare the capacity of each fraction to protect spermatozoa from oxidative stress. The common carp was selected, as it is among the most commonly cultured fish species and is a representative model teleost.

## MATERIAL AND METHODS

**Experimental fish, location and rearing conditions.** Procedures were conducted according to the principles of the Ethics Committee for the Protection of Animals in Research of the University of South Bohemia in České Budějovice, Research Institute of Fish Culture and Hydrobiology, Vodňany, based on the EU-harmonised Animal Welfare Act of the Czech Republic.

Male common carp ( $n = 10$ ) weighing 4.0–4.5 kg were obtained from the experimental station of the Faculty of Fisheries and Protection of Waters, University of South Bohemia at Vodňany, Czech Republic. Prior to experimentation, fish were held in 10 000 l outdoor hatchery tanks with constant flowing pond water. Water parameters followed seasonal ambient conditions in the range 20–22°C, pH 7.0–7.5, and 6.5–7.1 mg O<sub>2</sub>/l. Ambient light entered the tanks via netting that prevented fish from escaping.

**Semen collection and preparation of seminal plasma.** Spermiation was induced with an injection of carp pituitary extract at 1 mg/kg. 24 h post-injection, sperm samples were collected by abdominal massage directly into 10 ml plastic syringes, taking care to prevent contamination with blood, faeces, or urine. Samples were stored 30 min on ice (0–4°C) in closed assay tubes until assessment.

Each sperm sample was initially centrifuged at 700 g for 30 min followed by 10 min at 16 000 g at 4°C. The seminal plasma was carefully collected and stored at –80°C until analysis. The bicinchoninic acid assay (BCA) with the Infinite M200 photometer (Tecan, Switzerland) were used to determine protein concentrations.

**Seminal plasma fractionation.** Nanosep<sup>®</sup> 100 kDa, 50 kDa, 30 kDa, and 10 kDa centrifugal devices (Pall Laboratory, USA) were used to separate the seminal plasma into fractions according to the protein content. The separation of each sample usually started with the 100 kDa molecular weight cutoff (MWCO): 500 µl of SP was centrifuged at low speed, 2000 g, for 30 min. The filtrate from the bottom receiver was collected and divided into two aliquots. A 450 µl aliquot was filled to 500 µl with PBS buffer (137 mM NaCl, 2.7 mM KCl, 10 mM Na<sub>2</sub>HPO<sub>4</sub> and 1.8 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4), briefly mixed, centrifuged in a lower MWCO device, and collected as before. The second SP aliquot (20 µl) containing protein fractions with defined molecular weight (MW) was stored at –80°C until analysis.

<https://doi.org/10.17221/187/2018-CJAS>

**SDS-gel electrophoresis (SDS-PAGE).** The seminal plasma protein fractions obtained from each male were analysed by 12% polyacrylamide-bisacrylamide gel electrophoresis. Samples were re-suspended in buffer containing 65 mM Tris 10% (v/v) glycerol, 2% (w/v) sodium dodecyl sulphate (SDS), and 5% (v/v) beta-mercaptoethanol and denatured at 95°C for 3 min prior to loading to gels. Protein bands were visualised by Coomassie Brilliant Blue R-250 staining (Applichem, Germany) and scanned. GelQuant v. 2.7.0 (Bio-Imaging Systems, Israel) was applied to determine the MW of protein bands in each fraction.

**Evaluation of SOD activity in fractionated seminal plasma.** SOD activity was determined by the method of Marklund and Marklund (1974) based on the autoxidation of pyrogallol. Each prepared seminal plasma fraction was centrifuged at 12 000 g at 4°C for 30 min. The SOD activity was assessed spectrophotometrically at 420 nm and expressed as the quantity of enzyme per mg of protein. One unit of SOD activity is defined as the amount of the enzyme necessary to produce 50% dismutation of the superoxide radical per min. Results were expressed as international milliunits (mU) per mg of protein and obtained in triplicate for each sample.

**Evaluation of GR and GPx activity in fractionated seminal plasma.** GR activity was determined spectrophotometrically, measuring NADPH oxidation at 340 nm. GPx activity was assayed based on the rate of NADPH oxidation at 340 nm, by the coupled reaction with GR. The specific activity was determined using the extinction coefficient of 6.22 mM/cm (Lawrence and Burk 1976). One unit of GPx or GR activity was defined as the quantity of the enzyme that consumes 1  $\mu\text{mol}$  of substrate or generates 1  $\mu\text{mol}$  of product per min. The activity was expressed as mU/mg protein. GPx and GR activity was assessed in triplicate for each sample.

**Data analysis.** Prior to comparison of the variables, data were tested by the Kolmogorov test and the Bartlett test. Values of SOD, GR, and GPx were expressed as mean  $\pm$  standard deviation (SD) ( $n = 6$ ) and analysed by factorial analysis of variance (ANOVA). Statistical comparison was made by ANOVA followed by Tukey's HSD test for each analysed variable. All analyses were performed at a significance level of  $P < 0.05$  using STATISTICA (Version 9.0, 2009) software for MS Windows.

## RESULTS

**Protein separation by SDS-PAGE and gel imaging.** The SDS gel electrophoresis results revealed four distinct fractions with differing total protein content (Figure 1). The molecular weight of separated proteins, based on the band pattern on denatured SDS-PAGE, and band quantity of each fraction are summarised in Table 1. Briefly, the first seminal plasma fraction showed five protein bands with MW in the range 19–73 kDa. Fraction 2 comprised six protein bands of MW 18–57 kDa, fraction 3 consisted of three bands ranging from 35 to 58 kDa, and fraction 4 bands of 17 kDa and 18 kDa (Table 1).

**Enzymatic antioxidant activity in seminal plasma fractions.** The antioxidant activity was assessed as total SOD, GR, and GPx level.

**Superoxide dismutase.** The level of SOD varied significantly among seminal plasma fractions (Figure 2A). The highest value of SOD ( $2.81 \pm 0.11$  mU/mg protein) was observed in the fraction separated by a Nanosep 10 kDa device (fraction 4), with the lowest ( $1.83 \pm 0.18$  mU/mg protein) de-




Figure 1. Sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) pattern of common carp (*Cyprinus carpio*) seminal plasma protein fractions

line 1 = fraction separated in a Nanosep 100 kDa device, line 2 = fraction separated in a Nanosep 50 kDa device, line 3 = fraction separated in a Nanosep 30 kDa device, line 4 = fraction separated in a Nanosep 10 kDa device, M = molecular weight marker




Figure 2. Level of (A) superoxide dismutase (SOD), (B) glutathione reductase (GR) and (C) glutathione peroxidase (GPx) activity in common carp (*Cyprinus carpio*) seminal plasma fractions. Data are expressed as mean  $\pm$  SD fraction 1 = samples separated in a Nanosep 100 kDa device, fraction 2 = samples separated in a Nanosep 50 kDa device, fraction 3 = samples separated in a Nanosep 30 kDa device, fraction 4 = samples separated in a Nanosep 10 kDa device <sup>a-c</sup>significant differences between fractions (ANOVA,  $P < 0.05$ )

tected in fraction 1 separated by the Nanosep 100 kDa device. There was no significant difference (ANOVA;  $P > 0.05$ ) in SOD activity between

Table 1. Distribution of protein bands in seminal plasma fractions of common carp (*Cyprinus carpio*)

Seminal plasma fractions	Molecular weight of proteins (kDa) ( $n = 10$ )
1	73.0 $\pm$ 2.67
	57.4 $\pm$ 0.64
	49.4 $\pm$ 0.79
	31.2 $\pm$ 0.44
	19.6 $\pm$ 0.54
2	57.5 $\pm$ 2.12
	55.0 $\pm$ 0.63
	50.3 $\pm$ 0.73
	48.2 $\pm$ 0.75
	30.5 $\pm$ 0.49
3	18.3 $\pm$ 0.64
	58.4 $\pm$ 2.68
	39.3 $\pm$ 0.79
4	35.4 $\pm$ 1.15
	18.0 $\pm$ 0.60
	17.4 $\pm$ 0.43

the first and the second seminal plasma protein fractions (Figure 2A).

**Glutathione reductase.** Significant differences (ANOVA;  $P = 0.05$ ) in GR activity were seen among fractions 1, 2, and 4 separated by Nanosep 100 kDa, 50 kDa, and 10 kDa devices, respectively. The GR activity in seminal plasma protein fractions 2 and 3 ranged from  $23.38 \pm 0.53$  mU/mg protein to  $24.5 \pm 0.45$  mU/mg protein with no significant differences detected between them (Figure 2B). The highest GR level at  $25.15 \pm 0.54$  mU/mg protein was measured in fraction 4.

**Glutathione peroxidase.** Low variability was observed in GPx activity in seminal plasma protein fractions (Figure 2C) with no significant differences (ANOVA;  $P > 0.05$ ) between fractions 1, 2, and 4. The lowest value of GPx activity ( $14.9 \pm 0.4$  mU/mg protein) was found in fraction 2 separated by a Nanosep 50 kDa and the highest ( $16.35 \pm 0.54$  mU/mg protein) in fraction 4.

## DISCUSSION

Antioxidant activity assessed as the total SOD, GR, and GPx level showed a significant variation

<https://doi.org/10.17221/187/2018-CJAS>

among selected fractions, confirming the existence of an integrated antioxidant system in individual fractions of common carp seminal plasma. Nevertheless, the highest levels of SOD, GR, and GPx was recorded in fraction 4 that can make this fraction as a candidate to be used for improving storage protocols for sperm.

A large number of seminal plasma components influence the biological quality of semen (Ciereszko et al. 2011; Chacur 2012). Studies have provided direct evidence that specific components can be adsorbed into the surface of spermatozoa and have an important function in the process of capacitation (Manjunath and Therien 2002), sperm transport (Clark 2011), and fertilisation (Barrios et al. 2000). Seminal plasma fractions have also been associated with protective properties and show the ability to restore the integrity of damaged sperm (Barrios et al. 2000; Perez-Pe et al. 2001). In our study, SDS-PAGE electrophoresis was used to define the proteins contained in an SP fraction obtained by centrifugation in a Nanosep device. Four separated fractions were visualised on the gel that correspond to protein bands with the following MW: from 73 to 19 kDa (fraction 1); from 57 to 18 kDa (fraction 2); from 58 to 35 kDa (fraction 3); from 18 to 17 kDa (fraction 4). According to Dietrich et al. (2014, 2017), we can speculate that the protein band identified as 73 kDa belongs to the family of major proteins of carp seminal plasma, transferrin. Proteins of  $\geq 50$  kDa may be hemopexin-like proteins, whereas proteins with  $MW \geq 30$  kDa could relate to members of the large group of SP apolipoproteins. Proteins of  $MW \leq 19$  kDa may be cofilin-2-like proteins (Dietrich et al. 2017). However, during the separation of protein fractions using SDS-PAGE, proteins of similar molecular weight may overlap. Therefore, further investigation of protein content in fractionated seminal plasma is required.

The protective antioxidant system in semen is made up of both enzymes and non-enzymatic factors along with low molecular weight compounds with antioxidant capacity, all of which closely interact to ensure optimum protection against oxidative damage (Walczak-Jedrejowska et al. 2013). The quantity of enzymes varies among species and is present in both seminal plasma and spermatozoa (Shaliutina-Kolesova et al. 2013). Donkey and stallion semen displays the highest reported seminal plasma antioxidant activity (Cassani et

al. 2005), whereas bovine sperm exhibits a low level of activity (Bilodeau et al. 2000). Compared to mammals, information on the mechanism of action of antioxidant enzymatic systems in fish sperm is limited.

SOD is the most abundant enzyme in biological systems (Halliwell 2006) and is considered the first line of defence against the effects of oxyradicals in the cell through catalysing the dismutation of superoxide radicals to hydrogen peroxide and molecular oxygen (Li et al. 2010). We found considerable variation in SOD activity among fractions, with the highest value recorded in fraction 4, indicating that most SOD activity in seminal plasma of common carp is contained in its low molecular weight protein fraction, which may possess beneficial properties. Garcia et al. (2009) separated boar seminal plasma and showed a positive effect of certain fractions on sperm survival during *in vitro* storage. Barrios et al. (2005) reported that ram seminal plasma fraction containing low MW proteins may protect the sperm and reverse spermatozoon damage caused by cold shock. These observations, together with the findings of this study, suggest that separated fractions containing high concentrations of SOD and low MW proteins would be best equipped to counteract an increase in oxygen radical production.

GR provides intracellular defence of sperm against oxidative stress (Cheema et al. 2009), acting against lipid peroxidation (LP) caused by ROS and playing a major role in maintaining sperm quality (Shaliutina-Kolesova et al. 2014). We confirmed GR activity in all separated fractions, contrary to results of Marti et al. (2007) who found GR activity in only two fractions of ram seminal plasma. The variation of GR between fractions in our study showed a trend similar to that of SOD, with the highest value in fraction 4 and the lowest in fraction 1. The source of the low level of GR in fraction 1 may have been the presence of high MW proteins that inhibit the activity of GR.

Differences in GPx activity among fractions were also observed. GPx, a ROS scavenger, exerts a protective effect on spermatozoon membranes from LP damage (Alvarez and Storey 1989) and can inhibit LP of the cell membrane (Sharma and Agarwal 1996). Jervis and Robaire (2001) demonstrated high GPx activity in fractions of ejaculated ram semen, possibly due to increased protein synthesis, and reported a low level of GPx in the

<https://doi.org/10.17221/187/2018-CJAS>

sperm-rich fraction of ram semen. Three protein bands with peroxidase activity in the sperm-rich fraction of dog semen provided evidence that GPx plays the primary role in the decomposition of hydrogen peroxide (Koziorowska-Gilun and Strzezek 2011). We observed low variation in GPx level in the analysed seminal plasma fractions, hypothetically suggesting similarity in preservation of spermatozoon membrane against ROS. However, GPx activity should not be lower than that of SOD to avoid the production of toxic levels of hydrogen peroxide. GPx and SOD activity in separated fractions should be synergistic. It has been reported that both GPx and SOD play a key role in protecting mammalian sperm against oxygen radical induced damage leading to motility loss (Alvarez and Storey 1989).

## CONCLUSION

We confirmed that common carp seminal plasma can be separated by centrifugal devices into four protein fractions. The fractions contained proteins differing in molecular weight as well as in antioxidant activity. The highest levels of SOD, GR, and GPx was recorded in fraction 4 separated by a Nanosep 10 kDa device. We suggest that this fraction can be used as an additive in fish sperm preservation to provide protection to sperm during *in vitro* storage by reducing oxidative stress induced by ROS. Further work is needed to determine whether proteins present in separated seminal plasma fractions can be characterized, synthesized, and used as supplements in sperm preservation media to improve fish fertility.

**Acknowledgement.** The Lucidus Consultancy, UK is gratefully acknowledged for English correction.

## REFERENCES

- Aitken R.J., Gordon E., Harkiss D., Twigg J.P., Milne P., Jennings Z., Irvine D.S. (1998): Relative impact of oxidative stress on the functional competence and genomic integrity of human spermatozoa. *Biology of Reproduction*, 59, 1037–1046.
- Alvarez J.G., Storey B.T. (1989): Role of glutathione peroxidase in protecting mammalian spermatozoa from loss of motility caused by spontaneous lipid peroxidation. *Gamete Research*, 23, 77–90.
- Barrios B., Perez-Pe R., Gallego M., Tato A., Osada J., Muino-Blanco T., Cebrian-Perez J.A. (2000): Seminal plasma proteins revert the cold-shock damage on ram sperm membrane. *Biology of Reproduction*, 63, 1531–1537.
- Barrios B., Fernandez-Juan M., Muino-Blanco T., Cebrian-Perez J.A. (2005): Immunocytochemical localization and biochemical characterization of two seminal plasma proteins that protect ram spermatozoa against cold shock. *Journal of Andrology*, 26, 40–47.
- Bilodeau J.F., Chatterjee S., Sirard M.A., Gagnon C. (2000): Levels of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Molecular Reproduction and Development*, 55, 282–288.
- Cassani P., Beconi M.T., O’Flaherty C. (2005): Relationship between total superoxide dismutase activity with lipid peroxidation, dynamics and morphological parameters in canine semen. *Animal Reproduction Science*, 86, 163–173.
- Chacur M.G.M. (2012): Seminal plasma proteins as potential markers of relative fertility in Zebu Bulls (*Bos taurus-indicus*). In: Ghowsi K. (ed.): *Electrophoresis*. IntechOpen, 173–193.
- Cheema R.S., Bansal A.K., Bilaspuri G.S. (2009): Manganese provides antioxidant protection for sperm cryopreservation that may offer new consideration for clinical fertility. *Oxidative Medicine and Cellular Longevity*, 2, 152–159.
- Ciereszko A., Glogowski J., Dabrowski K. (2011): Biochemical characteristics of seminal plasma and spermatozoa of freshwater fish and the relation to semen biology, quality and cryopreservation. In: Tiersch T.R. and Green CC (eds.): *Cryopreservation in Aquatic Species*. World Aquaculture Society, Baton Rouge, USA, 46–79.
- Clark G.F. (2011): Molecular models for mouse sperm–oocyte binding. *Glycobiology*, 21, 3–5.
- Dietrich M.A., Arnold G.J., Nynca J., Frohlich T., Otte K., Ciereszko A. (2014): Characterization of carp seminal plasma proteome in relation to blood plasma. *Proteomics*, 98, 218–232.
- Dietrich M.A., Irnazarow I., Ciereszko A. (2017): Proteomic identification of seminal plasma proteins related to the freezability of carp semen. *Proteomics*, 162, 52–61.
- Fazeli F., Salimi S. (2016): Correlation of seminal plasma total antioxidant capacity and malondialdehyde levels with sperm parameters in men with idiopathic infertility. *Avicenna Journal of Medical Biochemistry*, 4, e29736.
- Garcia E.M., Calvete J.J., Sanz L., Roca J., Martinez E.A., Vazquez J.M. (2009): Distinct effects of boar seminal plasma fractions exhibiting different protein profiles on the functionality of highly diluted boar spermatozoa. *Reproduction in Domestic Animals*, 44, 200–205.
- Gazo I., Linhartova P., Shaliutina A., Hulak M. (2013): Influence of environmentally relevant concentrations of vin-

<https://doi.org/10.17221/187/2018-CJAS>

- clozolin on sperm quality, DNA integrity, and antioxidant responses in sterlet *Acipenser ruthenus* spermatozoa. *Chemico-Biological Interactions*, 203, 377–385.
- Halliwell B. (2006): Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life. *Journal of Plant Physiology*, 141, 312–322.
- Jervis K.M., Robaire B. (2001): Dynamic changes in gene expression along the rat epididymis. *Biology of Reproduction*, 65, 696–703.
- Koppers A.J., Garg M.L., Aitken R.J. (2010): Stimulation of mitochondrial reactive oxygen species production by unesterified, unsaturated fatty acids in defective human spermatozoa. *Free Radical Biology and Medicine*, 48, 112–119.
- Koziorowska-Gilun M., Strzerek R. (2011): Molecular forms of selected antioxidant enzymes in dog semen – electrophoretic identification. *Polish Journal of Veterinary Sciences*, 14, 29–33.
- Lahnsteiner F., Mansour N., Plaetzer K. (2010): Antioxidant systems of brown trout (*Salmo trutta f. fario*) semen. *Animal Reproduction Science*, 119, 314–321.
- Lawrence R.A., Burk R.F. (1976): Glutathione peroxidase activity in selenium deficient rat liver. *Biochemical and Biophysical Research Communications*, 71, 952–958.
- Li P., Li Z.H., Dzyuba B., Hulak M., Rodina M., Linhart O. (2010): Evaluating the impacts of osmotic and oxidative stress on common carp (*Cyprinus carpio* L.) sperm caused by cryopreservation techniques. *Biology of Reproduction*, 83, 852–858.
- Manjunath P., Therien I. (2002): Role of seminal plasma phospholipid-binding proteins in sperm membrane lipid modification that occurs during capacitation. *Journal of Reproductive Immunology*, 53, 109–119.
- Marklund S., Marklund G. (1974): Involvement of superoxide anion radical in autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *European Journal of Biochemistry*, 47, 469–474.
- Marti E., Mara L., Marti J.I., Muino-Blanco T., Cebrian-Perez J.A. (2007): Seasonal variations in antioxidant enzyme activity in ram seminal plasma. *Theriogenology*, 67, 1446–1454.
- Martinez-Paramo S., Diogo P., Dinis M.T., Herraes M.P., Sarasquete C., Cabrita E. (2012): Incorporation of ascorbic acid and  $\alpha$ -tocopherol to the extender media to enhance antioxidant system of cryopreserved sea bass sperm. *Theriogenology*, 77, 1129–1136.
- Maxwell W.M.C., de Graaf S.P., El-Hajj Ghaoui R., Evans G. (2007): Seminal plasma effects on sperm handling and female fertility. *Society of Reproduction and Fertility supplement*, 64, 13–38.
- Perez-Pe R., Barrios B., Muino-Blanco T., Cebrian-Perez J.A. (2001): Seasonal differences in ram seminal plasma revealed by partition in an aqueous two-phase system. *Journal of Chromatography B: Biomedical Sciences and Applications*, 760, 113–121.
- Ratto M.H., Delbaere L.T.J., Leduc Y.A., Pierson R.A., Adams G.P. (2011): Biochemical isolation and purification of ovulation-inducing factor (OIF) in seminal plasma of llamas. *Reproductive Biology and Endocrinology*, 9, 1–8.
- Saravia F., Wallgren M., Johannisson A., Rodriguez-Martinez H. (2007): Controlled cooling during semen cryopreservation does not induce capacitation of spermatozoa from two portions of the boar ejaculate. *International Journal of Andrology*, 30, 485–499.
- Shaliutina-Kolesova A., Gazo I., Cosson J., Linhart O. (2013): Comparison of oxidant and antioxidant status of seminal plasma and spermatozoa of several fish species. *Czech Journal of Animal Science*, 58, 313–320.
- Shaliutina-Kolesova A., Gazo I., Cosson J., Linhart O. (2014): Protection of common carp (*Cyprinus carpio* L.) spermatozoa motility under oxidative stress by antioxidants and seminal plasma. *Fish Physiology and Biochemistry*, 40, 1771–1781.
- Sharma R.K., Agarwal A. (1996): Role of reactive oxygen species in male infertility. *Urology*, 48, 835–850.
- Walczak-Jedrzejowska R., Wolski J.K., Slowikowska-Hilczner J. (2013): The role of oxidative stress and antioxidants in male fertility. *Central European Journal of Urology*, 66, 60–67.

Received: 2018–10–10

Accepted: 2019–05–17