

Soil aggregate stability index and particulate organic matter in response to differently afforested lands in the temperate regions of Iran

MASOOMEH SOLEIMANY¹, JAMSHID ESLAMDoust^{2*}, MOSLEM AKBARINIA²,
YAHYA KOOCH²

¹Department of Forestry, Sari Agricultural Sciences and Natural Resources University, Sari, Iran

²Department of Forestry, Faculty of Natural Resources and Marine Science,
Tarbiat Modares University, Tehran, Iran

*Corresponding author: j.eslamdoust@modares.ac.ir

Citation: Soleimany M., Eslamdoust J., Akbarinia M., Kooch Y. (2021): Soil aggregate stability index and particulate organic matter in response to differently afforested lands in the temperate regions of Iran. J. For. Sci., 67: 376–384.

Abstract: Aggregate Stability Index (ASI) and particulate organic matter (POM) are strongly influenced by land use and management. This work illustrates the effects of plantations on ASI and POM-C and POM-N in northern Iran. Three plantations of *P. deltoides* (PD), *T. distichum* (TD), *A. subcordata* (AS), and a fourth site – adjacent abandoned lands (BL, as control) were selected. Soil samples were taken within 16 quadrats of each plantation and BL from the two depths of 0–15 cm and 15–30 cm during the summer. Soil C was significantly higher under TD (2.10%) than under BL (2.02%) > PD (1.61%) > AS (1.30%). Soil N was found in ranked order of AS (8.99%) > TD (7.82%) > PD (5.30%) > BL (3.68%) ($P < 0.019$). The significantly higher ASI was found under TD (57.49) in comparison with PD (53.10), BL (51.23), and AS (36.57). The POM-C was as follows: TD (0.209%) > PD (0.141%) > AS (0.139%) > BL (0.075%) ($P = 0.020$). The highest POM-N was found under TD (0.035), followed by AS (0.0284%), PD (0.0288%), and BL (0.007%). The results indicate the positive effect of afforestation on soil ASI and POM-C and POM-N, especially in the surface layers of soil.

Keywords: afforestation; organic matter; soil properties; tree species; land use

Afforestation is defined as the conversion of non-forest lands to forested land (Berthrong, Finzi 2006) and is one of the ways of restoring degraded lands (Cannell 2003). Perhaps, one of the most important factors provided by forests is soil organic matter (SOM) accumulation and soil fertility maintenance (Boley et al. 2009). Besides the environmental and economic effects, the afforestation has an important impact on the soil physical and chemical properties (Xie et al. 2021). Thus, in addition to the major role of forest soil in creating change and diversity, forests, in turn, play a significant role in changing and developing forest soils (Fisher, Binkley 2000). Therefore, understanding the soil properties is considered

as a basis for appropriate forest management. Many studies have indicated that planted forest trees change soil properties (Antunes et al. 2008; Vesterdal et al. 2008; Ayres et al. 2009; Luan et al. 2010; Mueller et al. 2012; Kooch, Zoghi 2014). The results of most of these studies indicate that afforestation can amend and modify soil physical and chemical properties of afforested lands through adding litter to the soil, decomposition of litter, and also microbiological phenomena such as nitrogen fixation and mycorrhizal activity (Evans 1992).

The soil aggregate stability index (ASI) is related to its ability to maintain particle and pore arrangement when faced with different environmental stresses

This study was carried out with the financial support provided by Tarbiat Modares University.

<https://doi.org/10.17221/20/2021-JFS>

(Angers, Carter 1995), and assessment of soil structure is usually expressed based on aggregate stability (Bronick, Lal 2005). ASI plays a significant role in the development of root systems, carbon and water cycles as well as resistance against erosion (Barthès et al. 2008). ASI can also be controlled by management practices (Oades 1993). Afforested lands impact on soil structure by distributing SOM through litter and recycling of roots and root exudates. Thus, vegetation and soil organic carbon are some of the most important factors determining ASI (Carpenter, Chong 2010; Fattet et al. 2011). However, any decrease of SOM can lead to reduced fertility, soil structure instability, and declined soil productivity potential (Obalum et al. 2017).

SOM is considered as one of the main indicators of soil quality because it strongly influences the soil chemical, physical, and biological processes (Spaccini et al. 2004). SOM consists of two parts, namely humus substances and labile compounds. Particulate organic matter (POM) is a part of the soil labile compounds (Yoosefi et al. 2008). In comparison with the total SOM, POM consisting of organic carbon and nitrogen is very sensitive to changes in management (Haynes 2005).

ASI and POM are two important properties that are strongly influenced by different land covers, but little attention has been paid to these properties

concerning afforestation with different tree species in northern Iran. Due to the high amount of rainfall in the north of Iran and the increasing risk of water erosion, the necessity of afforestation on degraded lands is higher than ever before because trees due to creating higher infiltration capacities of water flow can prevent soil erosion (Kramer 2010). On the other hand, afforestation causes increasing POM in the soil (Mao, Zeng 2010), and POM affects ASI and soil fertility (Haynes 2005; Onweremadu et al. 2010). Little research has been done in this regard, and most research on the size and concentration of POM has been carried out for different management practices and land use (Meyer et al. 2012; Mujuru et al. 2013). Thus, the effect of different species on POM have been rarely studied (Yu et al. 2013), and little information is available. Nevertheless, the issue of which planted species can be more effective on ASI, POM-C, POM-N, and finally, soil protection remained unclear. This study aims to investigate the effect of the afforestation with different tree species on soil ASI, POM-C, and POM-N.

MATERIAL AND METHODS

Study area. The study was conducted on the southern coast of the Caspian Sea, Mazandaran Province, Northern Iran: Klodeh ($36^{\circ}35'N$, $52^{\circ}10'E$)




Figure 1. Location of the study area in the Mazandaran province, north of Iran

(Figure 1). The region has a temperate climate, with the mean annual temperature of 16.9 °C. The mean annual precipitation of the last 10-year period is 802–823.5 mm and the majority of precipitation falls between September and March. This region has a distinct dry season that stretches from April to August. The topography is characterized by flatlands to low hills at an elevation of 5 m a.s.l. According to WRB (World Reference Base), the soil type is Dystric Cambisol (WRB, IUSS 2014). The soil texture is silty loam with poor drainage (Eslamdoust 2015). The average pH ranges between 7.6 and 8.1.

The study plantations are composed of *Alnus subcordata* L. (AS), *Populus deltoides* L. (PD), and *Taxodium distichum* L. Rich. (TD). These treatments were planted in 2001. A description of the study sites is presented in Table 1 (Eslamdoust, Sohrabi 2018). An adjacent abandoned land (BL) was also selected as control.

Soil sampling and laboratory analysis. Due to the homogeneity of each afforestation in the study site, a network with dimensions of 200 m × 200 m with systematic random sampling was used. Soil samples (25 cm × 25 cm × 15 cm) were taken within 16 quadrats of each plantation and BL from two depths of 0–15 cm (D_1) and 15–30 cm (D_2) during the summer. The soil samples were stored at 4 °C for two weeks. Soils were air-dried and passed through a 2-mm sieve (aggregates were broken to pass through a 2-mm sieve) to laboratory analysis. Soil texture was measured using the Bouyoucos hydrometer method (Bouyoucos 1962). Bulk density was measured by the clod method. Soil organic C was determined using the Walkley-Black technique (Allison 1975). Total N was measured using a semi-micro Kjeldahl technique (Bremner, Mulvaney 1982). Available P

was determined with a spectrophotometer and by the Olsen method (Homer, Pratt 1961). POM-C and POM-N were determined by physical fractionation. 25 g of air-dried soil samples were dispersed with 100 mL of 5 g·L⁻¹ of sodium hexametaphosphate. The soil solution mixture was shaken for 1 h at high speed (= 500 rpm) on an end-to-end shaker and poured over a 0.053-mm sieve with several deionized water rinses. The soil remaining on the sieve was washed into a pre-weighed aluminium dish and dried at 60 °C for 24 h, then ground and analysed for C and N (Kooch et al. 2019).

Data analysis. The normality of variables was checked by the Kolmogorov-Smirnov test. The equality of the variances was checked using Levene's test. Two-way analysis of variance (ANOVA) was used to compare data on soil properties among plantations and depths. Duncan's test was employed to test for differences at the $P = 0.05$ level. All statistical analyses were conducted using the SPSS statistical software package (Version 19.0; 2010).

RESULTS

Clay and sand contents were not significantly different between afforested sites (Table 2). However, the highest value of clay and sand content was found under PD and AS, respectively. Soil silt contents decreased significantly in the order BL > TD > PD > AS. Soil bulk density was significantly higher in the AS treatment (1.58 gr·m⁻³) compared with PD, TD, and BL. Soil C was significantly higher under TD (2.10%) than under BL (2.02%) > PD (1.61%) > AS (1.30%). Soil N was found in ranked order of AS > TD > PD > BL with more than twofold value in the AS treatment (8.99%) compared to BL (3.68%). Higher values

Table 1. Site properties and plantation characteristics (Eslamdoust, Sohrabi 2018)

Plantation types		AS	PD	TD
DBH (cm)		31.32 ± 7.68 (15.92–50.96)*	27.43 ± 6.34 (12.42–43.94)	28.44 ± 6.83 (17.83–47.04)
Tree height (m)		21.98 ± 4.38 (10.1–31.8)	29.79 ± 4.77 (16.9–39.5)	17.74 ± 2.66 (9.8–24.1)
Crown height (m)		8.91 ± 2.69	11.45 ± 3.30	6.67 ± 1.94
Crown diameter (m)		4.02 ± 1.38	3.12 ± 1.20	4.25 ± 0.96
Stem density (trees·ha ⁻¹)		459	556	556
Main understory vegetation	shrubs	<i>Ruscus hyrcanus</i> ; <i>Smilax excels</i>	<i>Ruscus hyrcanus</i> ; <i>Smilax excels</i>	–
	herbaceous plants	<i>Carex remota</i> ; <i>Carex sylvatica</i> ; <i>Poa</i> sp.	<i>Carex remota</i> ; <i>Carex sylvatica</i> ; <i>Poa</i> sp.	

AS – *Alnus subcordata*; PD – *Populus deltoides*; TD – *Taxodium distichum*; * (min-max)

<https://doi.org/10.17221/20/2021-JFS>Table 2. Means (\pm SE, $n = 16$) of the soil variables analysed in the studied land covers and depths

Properties	Depth	AS	PD	TD	BL	Statistic description
Clay (%)	D ₁	33.5 (\pm 4.03)	35.7 (\pm 3.51)	26.5 (\pm 1.76)	33.3 (\pm 2.16)	Land cover: $F = 1.003$; $P = 0.398$ Depth: $F = 0.289$; $P = 0.593$ L \times D: $F = 0.672$; $P = 0.769$
	D ₂	32.3 (\pm 5.61)	33.0 (\pm 4.72)	3.7 (\pm 3.02)	27.5 (\pm 2.16)	PD > AS > BL > TD D ₁ > D ₂
Silt (%)	D ₁	34.3 (\pm 1.82)	38.3 (\pm 4.14)	41.0 (\pm 4.50)	42.8 (\pm 2.90)	Land cover: $F = 3.643$; $P = 0.018$ Depth: $F = 1.863$; $P = 0.178$ L \times D: $F = 0.337$; $P = 0.798$
	D ₂	35.0 (\pm 5.69)	41.3 (\pm 4.30)	43.8 (\pm 2.25)	50.8 (\pm 2.60)	BL ^a > TD ^{ab} > PD ^{ab} > AS ^b D ₂ > D ₁
Sand (%)	D ₁	32.3 (\pm 4.70)	26.0 (\pm 5.14)	32.5 (\pm 5.23)	24.0 (\pm 2.61)	Land cover: $F = 1.191$; $P = 0.321$ Depth: $F = 0.353$; $P = 0.555$ L \times D: $F = 0.198$; $P = 0.897$
	D ₂	32.8 (\pm 6.16)	25.8 (\pm 6.95)	25.5 (\pm 4.74)	21.8 (\pm 2.80)	TD > AS > PD > BL D ₁ > D ₂
Bulk density (gr·m ⁻³)	D ₁	1.58 (\pm 0.07)	1.41 (\pm 0.08)	1.42 (\pm 0.11)	1.35 (\pm 0.05)	Land cover: $F = 4.308$; $P = 0.008$ Depth: $F = 0.500$; $P = 0.482$ L \times D: $F = 0.648$; $P = 0.578$
	D ₂	1.61 (\pm 0.09)	1.48 (\pm 0.09)	1.28 (\pm 0.10)	1.23 (\pm 0.07)	AS ^a > TD ^{ab} > PD ^b > BL ^b D ₁ > D ₂
C (%)	D ₁	1.30 (\pm 0.03)	1.61 (\pm 0.14)	2.10 (\pm 0.20)	2.02 (\pm 0.21)	Land cover: $F = 6.65$; $P = 0.001$ Depth: $F = 14.427$; $P = 0.000$ L \times D: $F = 1.910$; $P = 0.138$
	D ₂	1.25 (\pm 0.07)	1.37 (\pm 0.03)	1.49 (\pm 0.13)	1.49 (\pm 0.07)	TD ^a > BL ^{ab} > PD ^{bc} > AS ^c D ₁ > D ₂
N (%)	D ₁	8.99 (\pm 1.33)	5.30 (\pm 1.42)	7.82 (\pm 2.07)	3.68 (\pm 0.64)	Land cover: $F = 3.588$; $P = 0.019$ Depth: $F = 1.74$; $P = 0.19$ L \times D: $F = 1.11$; $P = 0.35$
	D ₂	5.10 (\pm 1.92)	6.27 (\pm 0.69)	6.51 (\pm 1.11)	2.88 (\pm 0.73)	AS ^a > TD ^a > PD ^{ab} > BL ^b D ₁ > D ₂
C.N	D ₁	3.87 (\pm 0.49)	4.29 (\pm 0.37)	5.36 (\pm 0.72)	10.87 (\pm 1.41)	Land cover: $F = 17.37$; $P = 0.00$ Depth: $F = 0.002$; $P = 0.965$ L \times D: $F = 2.32$; $P = 0.085$
	D ₂	6.20 (\pm 0.79)	5.16 (\pm 0.47)	3.96 (\pm 0.34)	9.18 (\pm 1.55)	BL ^a > TD ^b > PD ^b > AS ^b D ₂ > D ₁

D1 – 0–15 cm soil depth; D2 – 15–30 cm soil depth; AS – *Alnus subcordata*; PD – *Populus deltoides*; TD – *Taxodium distichum*; BL – bare land; L \times D – Land cover \times Depth; F – F -statistic; different letters indicate significant differences ($P < 0.05$ by Duncan test) between land covers

of C/N were observed in the BL treatment (10.87) compared with TD (5.36), PD (4.29), and AS (3.87). Values of clay, silt, sand, bulk density, N and C/N were not found significantly different between the studied depths. However, soil C was significantly higher at D₁ (1.76%) than at D₂ (1.40%).

Our results showed that there were significant differences in ASI, POM-C, and POM-N between AS, PD, TD, and BL (Table 3). A significantly higher ASI was found in TD (57.49) in comparison with PD (53.10), BL (51.23), and AS (36.57) (Figure 2). ASI was insignificantly higher at D₁ (53.18) than at D₂ (46.01). POM-C was as follows: TD (0.209%) > PD (0.141%) > AS (0.139%) > BL (0.075%) (Figure 3). POM-C was significantly higher at D₁ (0.244%) than at D₂ (0.098%). The highest POM-N was found under TD (0.035), followed by AS (0.0284%), PD (0.0288%), and BL (0.007%) (Figure 4). POM-N was significantly higher at D₁ (0.039%) than at D₂ (0.021%).

ASI had a positive correlation with C and POM-N (Table 4). Soil clay and silt had a negative correlation with soil sand. Soil silt was positively correlated with C. C was positively correlated with POM-C.

DISCUSSION

Afforestation improves soil quality by supplies of nutrition resources and reduces soil erosion (Berthrong et al. 2009). Our findings showed that soils under coniferous TD have a significantly higher C than soils under broadleaved AS and PD. The changes in C in different land cover also reflect the differences in quantity and quality of litter inputs, litter decomposition rates, and root biomass C (Zheng et al. 2008). Contrary to our findings, previous reports (Salamon et al. 2004; Wardle et al. 2006) claimed that the type of aboveground vegetation has no remarkable effect on the soil features. According to Kooch et al. (2012), soil total N content is the main factor that affects the decomposition of litter, and the litter decomposition rate is a source of soil CO₂ emission. In the present study, N values were in ranked order




Figure 2. Mean (\pm SE; $n = 16$) of Aggregate Stability Index (ASI) under different land covers




Figure 3. Mean (\pm SE; $n = 16$) of particulate organic carbon (POM-C) under different land covers




Figure 4. Mean (\pm SE; $n = 16$) of particulate organic nitrogen (POM-N) under different land covers

AS – *Alnus subcordata*; PD: *Populus deltoides*; TD – *Taxodium distichum*; BL – bare land; different letters indicates significant differences and same letters indicates non-significant differences

Table 3. Statistical descriptions of Agregate Stability index (ASI), POM-C and POM-N in the studied land covers and depths

Properties	Land cover		Depth		Land cover \times Depth	
	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>	<i>F</i>	<i>P</i>
ASI	5.78	0.01	3.68	0.06	1.97	0.13
POM-C	1.59	0.02	7.91	0.01	0.55	0.07
POM-N	3.28	0.03	4.01	0.05	0.56	0.67

POM-C – particulate organic matter-carbon; POM-N – particulate organic matter-nitrogen; *F* – *F*-statistic

<https://doi.org/10.17221/20/2021-JFS>

Table 4. Pearson correlation coefficients (*r*) of soil properties across different land covers

	ASI	Clay	Silt	Sand	Bulk Density	C	N	POM-C	POM-N
ASI	1	0.10	0.22	−0.24	−0.08	0.25*	0.07	0.20	0.25*
Clay		1	0.02	−0.65**	0.03	0.007	0.15	0.007	−0.09
Silt			1	−0.73**	0.01	0.30*	0.05	0.02	−0.14
Sand				1	−0.03	−0.23	−0.06	−0.24	0.17
Bulk Density					1	−0.10	0.15	0.04	−0.02
C						1	−0.02	0.35**	−0.01
N							1	−0.09	0.11
POM-C								1	0.19
POM-N									1

*correlation significant at the 0.05 level (2-tailed); **correlation significant at the 0.01 level (2-tailed); ASI – Aggregate Stability Index; POM-C – particulate organic matter-carbon; POM-N – particulate organic matter-nitrogen

of TD > PD > AS, which may be related to soil silt. The C/N ratio did not significantly differ between the coniferous TD and deciduous AS and PD at 0–15 cm depth. Different results have been reported based on a research review on the variability of soil C and N under various land covers. Soil chemical features are the influential factors affecting surface soil C and N (Zhou et al. 2015). Afforested lands with different species produced litterfall with various litter mass and chemicals that can affect C and N concentrations (Van den Berg et al. 2012).

Higher ASI in PD and TD compared to BL was expected because of conditions of undisturbed lands. The results of Nael et al. (2004) are consistent with our findings. Afforested lands influence soil structure by adding SOM to the soil through litterfall (Tajik 2004; Khazaee et al. 2008; Carpenter, Chong 2010; Fattet et al. 2011). However, lower ASI under AS is probably more related to the soil texture. The factors affecting ASI have been reported in numerous studies. Soil texture and clay content (Kemper, Koch 1966), acidity, SOM content (Kandiah 1976), and management practices and land use (i.e. natural forest, afforested lands, or agricultural lands) (Emadodin et al. 2009; Oades 1993) are among the factors influencing the aggregate stability. Thus, according to the above-mentioned factors and the effect of soil texture on aggregate stability, reduced ASI in AS could be explained by higher sand content (light texture) in soil under AS (Pearson correlation results). Based on the Pearson correlation, ASI is more closely related to soil C and POM-N. Studies of Khazaee et al. (2008) and Onweremadu et al. (2010) indicate a positive effect of C on ASI. It is well-known that there is a direct linear rela-

tionship between ASI and soil C (Six et al. 2002; Carpenter, Chong 2010). Higher ASI at the depth of 0–15 cm compared to 15–30 cm can be pertinent to higher accumulation of SOM in the soil surface layers. SOM is considered a dominant indicator that is influencing aggregate stability (Cantón et al. 2009). Soil dispersion decreases significantly with increasing SOM and ASI (Emadi et al. 2009).

Land use changes individually the cultivation operations and land covers can differently affect SOM compounds in temperate regions. POM-C like SOM, are indices of soil sensitivity to land-use changes (Hajabbasi, Fallahzadeh 2010). In AS, PD, and TD, a higher value of POM (POM-C and POM-N) identifying aboveground litter was found through different land covers considering an influential POM source (Poeplau, Don 2013; Kooch et al. 2019). However, the differences were probably majorly due to the physical and chemical nature of the litterfall inputs by different tree species (Mendham et al. 2004). Increased POM-C in forested lands usually results from adding organic materials from trees of large size because entirely different from SOM of bare lands (and agricultural lands) in terms of diverse chemical composition and particle size (Mokhtari Karchegani et al. 2011). The measured POM values were reported to vary considerably in different soils (Christensen 2001). However, generally soils with perennial vegetation and a more prolonged period of litter return into the soil (e.g. forest soils) show the highest POM. Mokhtari Karchegani et al. (2011) reported a 60–70% reduction in SOM in bare lands compared to afforested lands. Thus, higher POM in afforested lands compared to bare lands is predictable due to the influence of tree cover and litter from the trees. In the present study, the highest

ASI was below TD, followed by PD. According to the Pearson correlation results, the higher ASI under TD and PD may be associated with higher POM. Also, the highest POM-C and POM-N were under TD, which in turn may impact on ASI. However, inconsistently with our results Soucémariadin et al. (2019) indicated that soils under deciduous trees were significantly higher in POM than under coniferous trees.

POM-C and POM-N were significantly higher at 0–15 cm soil depth than at 15–30 cm depth. It is not surprising given the higher SOM in the soil surface layers due to litter presence. Generally, it has been reported that the soil surface behaves more actively in terms of the fertility parameters while not so accurate for the deeper layers (Rumpel, Kogel-Knabner 2011). Soil C and N are the most labile C and N fractions in the biogeochemical cycling of topsoil (Zhou et al. 2015) while being affected by vegetation type and litter quality (Kooch et al. 2021). POM in topsoil is more likely to originate from the organic layers than directly from fresh leaves/needles (Soucémariadin et al. 2019). Thus, the topsoil functions as nutrient provision, storage, and finally, increased stand growth is clear (Wang et al. 2014).

CONCLUSION

POMs are intermediate matters between newly shed litter and decomposed SOM while a source of SOM determining the future status of nutrient and soil quality is considered. POM can play a significant role as a sensitive indicator reflecting negative impacts of forest degradation on soil quality. Based on our findings, ASI decreased in ranks order of TD > PD > BL > AS, which can be assigned to the lower values of soil silt and C. Our results revealed a positive effect of afforestation with AS, PD, and TD on POM-C and POM-N. Afforested lands have more remarkably enhanced POM-C and POM-N compared to BL. Especially in terms of POM-N, there were significant differences between afforested lands and BL. Our findings indicate that afforestation with this species improves the properties of soil fertility and stability. Thus afforestation is recommended to increase soil quality and prevent erosion caused by rainfall in these areas. It can be concluded that afforested lands increase soil ASI and POMs, finally ensuring future soil fertility.

Contribution of the authors: Masoomah Soleimany: field works and collecting the data, the labora-

tory analysis; Jamshid Eslamdoust: running the data analysis, writing the paper; Moslem Akbarinia: designing the experiment; Yahya Kooch: designing the experiment, writing the paper, supervising the work.

Acknowledgement: Here we cherish the memory of Dr. Akbarinia (third author of the paper) that we lost because of COVID-19. We will always remember the everlasting of his smile, kindness, and passion.

We also acknowledge the helpful comments and suggestions of two anonymous reviewers.

REFERENCES

- Allison L.E. (1975): Organic carbon. In: Black C.A. (ed.): Methods of Soil Analysis, Part 2. Madison, American Society of Agronomy: 1367–1378.
- Angers D.A., Carter M.R. (1995): Aggregation and organic matter storage in cool, humid agricultural soils. In: Carter M.R., Stewart B.A. (eds.): Structure and Organic Matter Storage in Agricultural Soils. Boca Raton, CRC Press: 193–211.
- Antunes S.C., Pereira R., Sousa J.P., Santos M.C., Goncalves F. (2008): Spatial and temporal distribution of litter arthropods in different vegetation covers of Porto Santo Island (Madeira Archipelago, Portugal). European Journal of Soil Biology, 44: 45–56.
- Ayres E., Steltzer H., Berg S., Wallenstein M.D., Simmons B.L., Wall D.H. (2009): Tree species traits influence soil physical, chemical, and biological properties in high elevation forests. PLoS ONE, 4: e5964.
- Barthès B.G., Kouakoua E., Larré-Larrouy M.C., Razafimbelo T.M., de Luca E.F., Azontonde A., Neves C.S.V.J., de Freitas P.L., Feller C.L. (2008): Texture and sesquioxide effects on water stable aggregates and organic matter in some tropical soils. Geoderma, 143: 14–25.
- Berthrong S.T., Finzi A.C. (2006): Amino acid cycling in three cold-temperate forests of the northeastern USA. Soil Biology and Biochemistry, 38: 861–869.
- Berthrong S.T., Jobbágy E., Jackson R.B. (2009): A global meta-analysis of soil exchangeable cations, pH, carbon, and nitrogen with afforestation. Ecological Applications, 19: 2228–2241.
- Boley J.D., Drew A.P., Andrus R.E. (2009): Effects of active pasture, teak (*Tectona grandis*) and mixed native plantations on soil chemistry in Costa Rica. Forest Ecology and Management, 257: 2254–2261.
- Bouyoucos G.J. (1962): Hydrometer method improved for making particle size analysis of soils. Agronomy Journal, 54: 464–465.
- Bremner J.M., Mulvaney C.S. (1982): Nitrogen - total. In: Page A.L., Miller R.H., Keeney R.R. (eds): Methods of Soil Analysis, Part 2. Madison, American Society of Agronomy: 595–624.

<https://doi.org/10.17221/20/2021-JFS>

- Bronick C.J., Lal R. (2005): Manuring and rotation effect on soil organic carbon concentration for different aggregate size fractions on two soils in northeastern Ohio, USA. *Soil Tillage Research*, 81: 239–252.
- Cantón Y., Solé-Benet A., Asensio C., Chamizo S., Puigdefábregas J. (2009): Aggregate stability in range sandy loam soils Relationships with runoff and erosion. *CATENA*, 77: 192–199.
- Cannell M.G.R. (2003): Carbon sequestration and biomass energy offset: theoretical, potential and achievable capacities globally, in Europe and the UK. *Biomass and Bioenergy*, 24: 97–116.
- Carpenter D.R., Chong G.W. (2010): Patterns in the aggregate stability of Mancos Shale derived soils. *CATENA*, 80: 65–73.
- Christensen B.T. (2001): Physical fractionation of soil and structural and functional complexity in organic matter turnover. *European Journal of Soil Science*, 52: 345–353.
- Emadi M., Baghernejad M., Memarian H.R. (2009): Effect of land-use change on soil fertility characteristics within water-stable aggregates of two cultivated soils in northern Iran. *Land Use Policy*, 26: 452–457.
- Emadodin I., Reiss S., Bork H.R. (2009): A study of the relationship between land management and soil aggregate stability (case study near Albersdorf, Northern-Germany). *ARP Journal of Agricultural and Biological Science*, 4: 48–53.
- Eslamdoust J. (2015): Allometric equations, biomass estimation and carbon sequestration of *Populus deltoids*, *Alnus subcordata* and *Taxodium distichum* trees in 20 years old-plantation, northern Iran. [M.Sc. Thesis.] Tehran, Tarbiat Modares University..
- Eslamdoust J., Sohrabi H. (2018): Carbon storage in biomass, litter, and soil of different native and introduced fast-growing tree plantations in the South Caspian Sea. *Journal of Forestry Research*, 29: 449–457.
- Evans J. (1992): *Plantation Forestry in the Tropics*. Oxford, Oxford University Press: 403.
- Fattet M., Fu Y., Ghestem M., Ma W., Foulonneau M., Nespoulous J., Le Bissonnais Y., Stokes A. (2011): Effects of vegetation type on soil resistance to erosion: Relationship between aggregate stability and shear strength. *CATENA*, 87: 60–69.
- Fisher R.F., Binkley D. (2000): *Ecology and Management of Forest Soils*. New York, John Wiley and Sons: 489.
- Hajabbasi M.A., Fallahzadeh J. (2010): Aggregation, carbohydrate, total and particulate organic carbon changes by cultivation of an arid soil in Central Iran. In: Gilkes R., Prakhongkep N. (eds): *Proceedings of the 19th World Congress of Soil Science, Soil Solutions for a Changing World*, Brisbane Aug 1–6, 2010: 3934–3937.
- Haynes R.J. (2005): Labile organic matter fractions as central components of quality of agricultural soil: An overview. *Advances in Agronomy*, 85: 221–268.
- Homer C.D., Pratt P.F. (1961): *Methods of Analysis for Soils, plants and waters*. Berkeley, University of California, Agricultural Sciences Press: 309.
- Kandiah A. (1976): Influence of organic matter on the erodibility of a saturated illitic soil. *Mededelingen van de Faculteit Landbouwwetenschappen Rijksuniversiteit Gent*, 41: 397–406.
- Kemper W.D., Koch E.J. (1966): *Aggregate Stability of Soils from Western United States and Canada*. Washington, Agricultural Research Service, U.S. Department of Agriculture: 52.
- Khazaei A., Mosaddeghi M., Mahboub A. (2008): Evaluation of aggregate stability in 21 series of Hamedan Province Soil by wet sieve method and its relationship with the inherent characteristics of the soil. *Journal of Agriculture: Water, Soil and Plants in Agriculture*, 8: 171–181. (in Persian)
- Kooch Y., Zoghi Z. (2014): Comparison of soil fertility of *Acer insigne*, *Quercus castaneifolia*, and *Pinus brutia* in the Hyrcanian forests of Iran. *Chinese Journal of Applied and Environmental Biology*, 20: 899–905.
- Kooch Y., Hosseini S.M., Zaccane C., Jalilvand H., Hojjati S.M. (2012): Soil organic carbon sequestration as affected by afforestation: the Darab Kola forest (north of Iran) case study. *Journal of Environmental Monitoring*, 14: 2438–2446.
- Kooch Y., Sanji R., Tabari M. (2019): The effect of vegetation change in C and N contents in litter and soil organic fractions of a Northern Iran temperate forest. *CATENA*, 178: 32–39.
- Kooch Y., Ghorbanzadeh N., Wirth S., Novara A., Piri A.S. (2021): Soil functional indicators in a mountain forest-rangeland mosaic of northern Iran. *Ecological Indicators*, 126: 107672.
- Kramer I. (2010): Rainfall partitioning and soil water dynamics along a tree species diversity gradient in a deciduous old-growth forest in Central Germany. [Ph.D. Thesis.] Göttingen, Mathematisch-Naturwissenschaftlichen Fakultäten der Georg-August-Universität Göttingen.
- Luan J., Xiang C., Liu S., Luo Z., Gong Y., Zhu X. (2010): Assessments of the impacts of Chinese fir plantation and natural regenerated forest on soil organic matter quality at Longmen mountain, Sichuan, China. *Geoderma*, 156: 228–236.
- Mao R., Zeng D.H. (2010): Changes in soil particulate organic matter, microbial biomass, and activity following afforestation of marginal agricultural lands in a semi-arid area of Northeast China. *Environmental Management*, 46: 110–116.
- Mendham D.S., Heagney E.C., Corbeels M., O'Connell A.M., Grove T.S., McMurtrie R.E. (2004). Soil particulate organic matter effects on nitrogen availability after afforestation with *Eucalyptus globulus*. *Soil Biology and Biochemistry*, 36: 1067–1074.
- Meyer S., Leifeld J., Bahn M., Fuhrer J. (2012): Land-use change in subalpine grassland soils: Effect on particulate organic carbon fractions and aggregation. *Journal of Plant Nutrient and Soil Science*, 175: 401–409.

- Mokhtari Karchegani P., Ayoubi S., Mosaddeghi M., Malekian M. (2011): Effects of land use and slope gradient on soil organic carbon pools in particle-size fractions and some soil physico-chemical properties in hilly regions, western Iran. *Electronic Journal of Soil Management and Sustainable Production*, 1: 23–41.
- Mueller K.L., Eissentat D.M., Hobbie S.E., Oleksyn J., Jagodzinski A.M., Reich P.B., Chadwick O.A., Chorover J. (2012): Tree species effect on coupled cycles of carbon, nitrogen, and acidity in mineral soils at a common garden experiment. *Biogeochemistry*, 111: 601–614.
- Mujuru L., Mureva A., Velthorst E.J., Hoosbeek M.R. (2013): Land use and management effects on soil organic matter fractions in Rhodic Ferralsols and Haplic Arenosols in Bindura and Shamva districts of Zimbabwe. *Geoderma*, 209: 262–272.
- Nael M., Khademi H., Hajabbasi M.A. (2004): Response of soil quality indicators and their spatial variability to land degradation in central Iran. *Apply Soil Ecology*, 27: 221–231.
- Oades J.M. (1993): The role of biology in the formation, stabilization and degradation of soil structure. In: Brussaard L., Kooistra M.J. (eds.): *Soil Structure/Soil Biota Interrelationships*, International Workshop on Methods of Research on Soil Structure/Soil Biota Interrelationships, Wageningen, Nov 24–28, 1991: 377–400.
- Obalum S.E., Chibuike G.U., Peth S., Ouyang Y. (2017): Soil organic matter as sole indicator of soil degradation. *Environmental Monitoring and Assessment*, 189: 176.
- Onweremadu E., Izuogu O., Akamigbo F. (2010): Aggregation and pedogenesis of seasonally inundated soils of a tropical watershed. *Chiang Mai Journal of Science*, 37: 74–84.
- Poepplau C., Don A. (2013): Sensitivity of soil organic carbon stocks and fractions to different land-use changes across Europe. *Geoderma*, 192: 189–201.
- Rumpel C., Kögel-Knabner I. (2011): Deep soil organic matter – a key but poorly understood component of terrestrial C cycle. *Plant and Soil*, 338: 143–158.
- Salamon J.A., Schaefer M., Alpehi J., Schmid B., Scheu S. (2004): Effects of plant diversity on Collembola in an experimental grassland ecosystem. *Oikos*, 106: 51–60.
- Six J., Callewaert P., Lenders S., De Gryze S., Morris S.J., Gregorich E.G., Paul E.A., Paustian K. (2002): Measuring and understanding carbon storage in afforested soils by physical fractionation. *Soil Science Society of America Journal*, 66: 1981–1987.
- Soucémariadin L., Cécillon L., Chenu C., Baudin F., Nicolas M., Girardin C., Delahaie A., Barré P. (2019): Heterogeneity of the chemical composition and thermal stability of particulate organic matter in French forest soils. *Geoderma*, 342: 65–74.
- Spaccini R.J., Mbagwu J.S.C., Igwe C.A., Conte P., Piccolo A. (2004): Carbohydrates and aggregation in lowland soils of Nigeria as influenced by organic input. *Soil and Tillage Research*, 75: 161–172.
- Tajik F. (2004): Evaluation of aggregate stability in some parts of Iran. *Water and Soil Science (Journal of Science and Technology of Agriculture and Natural Resources)*, 8: 123. (in Persian)
- Van den Berg L.J., Shotbolt L., Ashmore M.R. (2012): Dissolved organic carbon (DOC) concentrations in UK soils and the influence of soil, vegetation type and seasonality. *Science of the Total Environment*, 427: 269–276.
- Vesterdal L., Schmidt I.K., Callesen I., Nilsson L.O., Gundersen P. (2008): Carbon and nitrogen in forest floor and mineral soil under six common European tree species. *Forest Ecology and Management*, 255: 35–48.
- Wang H.M., Wang W.J., Chen H., Zhang Z., Mao Z., Zu Y.G. (2014): Temporal changes of soil physico-chemical properties at different soil depths during larch afforestation by multivariate analysis of covariance. *Ecology and Evolution*, 4: 1039–1048.
- Wardle D.A., Yeates G.W., Barker G.M., Bonner K.I. (2006): The influence of plant litter diversity on decomposer abundance and diversity. *Soil Biology and Biochemistry*, 38: 1052–1062.
- WRB, IUSS (2014): *World Reference Base for Soil Resources 2014: International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*. 3rd ed. Rome, Food and Agriculture Organization: 192.
- Xie H., Tang Y., Yu M., Wang G.G. (2021): The effects of afforestation tree species mixing on soil organic carbon stock, nutrients accumulation, and understory vegetation diversity on reclaimed coastal lands in Eastern China. *Global Ecology and Conservation*, 26: e01478.
- Yoosefi M., Shariatmadari H., Hajabbasi M.A. (2008): Determination of some of labile C pools as soil quality index. *Journal of Water and Soil Science*, 11: 429–439. (in Persian)
- Yu Y., Lu L., Du H., Peng W., Zeng F., Song T. (2013): Distribution of organic carbon and aggregation of calcareous soil in different forest types. *Acta Botanica Boreali-Occidentalia Sinica*, 33: 1011–1019.
- Zheng H., Ouyang Z., Xu W., Wang X., Miao H., Li X., Tian Y. (2008): Variation of carbon storage by different reforestation types in the hilly red soil region of southern China. *Forest Ecology and Management*, 255: 1113–1121.
- Zhou W.-J., Sha L.-Q., Schaefer D.A., Zhang Y.-P., Song Q.-H., Tan Z.-H., Guan H.-L. (2015): Direct effects of litter decomposition on soil dissolved organic carbon and nitrogen in a tropical rainforest. *Soil Biology and Biochemistry*, 81: 255–258.

Received: February 20, 2021

Accepted: May 25, 2021