

Biomass and carbon stocks in *Schima superba* dominated subtropical forests of eastern China

A. ALI^{1,2}, W.J. MA^{1,2}, X.D. YANG^{1,2}, B.W. SUN^{1,2}, Q.R. SHI^{1,2}, M.S. XU^{1,2}

¹Department of Environmental Sciences, East China Normal University, Shanghai, P. R. China

²Tiantong National Forest Ecosystem Observations and Research Station, Ningbo, Zhejiang, P. R. China

ABSTRACT: Quantitative relationships between stand indices and carbon dioxide (CO₂) stocking are missing in the evergreen broadleaved forests (EBLFs) in eastern China and this hinders to estimate carbon (C) budget in the subtropical region. We determined the vegetation-soil C pool and CO₂ stocking using stand indices [diameter at breast height (DBH), total height (H) and wood density] in *Schima superba* dominated EBLFs in the Tiantong National Forest Park in eastern China. Vegetation biomass was determined by a non-destructive method using the tree volume and wood density approach while soil C concentration was determined using the oil bath-K₂CrO₇ titration method. Finally, multiple regression and one-way ANOVA with LSD test were used for data analysis. Results showed that total C stocks in the vegetation and the 0–20 cm surface soil were 90.53 t·ha⁻¹ and 116.24 t·ha⁻¹, respectively. The study revealed that the total amount of CO₂ stocks in the studied forest is 331.87 t·ha⁻¹. One-way ANOVA with LSD test showed that CO₂ stocks varied significantly ($P < 0.05$) between the tree growth stages. There was a significant variation in CO₂ stocking capacity within sapling and pole growth stages but no significant variation within standard stage. The stepwise multiple regression analysis showed that DBH, BA and H were related to the C stocking while wood density had no significant effect. The significant amount of C stocking in EBLFs in the Tiantong National Forest Park of eastern China showed the potential and significant C stocks by trees. As the C pool structure changes due to a change in the forest type and location, therefore this study is important to estimate C stocks and predict CO₂ stocks from stand indices in EBLFs which serve as a scientific basis for sustainable forestry operations, rational utilization of forest resources and global warming reduction in EBLFs in subtropical regions of China.

Keywords: carbon stocks; evergreen broadleaved forest; forest inventory-based approaches; tree growth; wood density

The increase of carbon dioxide (CO₂) in the atmosphere is becoming a global issue. Carbon (C) is sequestered by the plant photosynthesis and stored as biomass in different components of the tree. During photosynthesis, trees act as a sink for CO₂ by fixing C and sequester excess C as biomass in different tree organs. The net long-term CO₂ sequestration of forest changes through the life span as trees grow up, die and decay (NOWAK, CRANE 2002). Trees absorb CO₂ from the atmosphere by getting an increase in their biomass through growth and sequester it in the plant tissues (MATHEW et al. 2000) resulting in development of different tree components. As the tree biomass performs growth, the C held by the tree also increases the CO₂ stock

(TAGUPA et al. 2010). The CO₂ sequestration rate depends on the growth (diameter) characteristics of the tree species (HUY, ANH 2008). The plant having a higher quantity of biomass reflects the higher CO₂ sequestration in the whole tree as well as in tree components (overbark stem, branches and leaves) (JANA et al. 2011). Forest trees are recognized as very important in the global C cycling, because the amount of C stored in plant biomass globally exceeds that of atmospheric CO₂, and nearly 90% of the plant biomass C is stored in tree biomass (MOONEY et al. 2001). The quantification of forest biomass has a long history and has received renewed attention in the last decades because forest biomass represents about 44% of the

Supported by the National Natural Science Foundation of China, Grants No. 31070383 and 31270475, and by the ECNU, Project for Ecology No. '211'.

globe forest C pool (PAN et al. 2011) and therefore plays a crucial role in climate change mitigation. Furthermore, sequestration of CO₂ among forests depends on forest type, dominant tree species and forest stand age (HUY, ANH 2008). This highlights the need to precisely determine the amount of C stored and CO₂ sequestered in each of specific forest ecosystems.

For estimating C stocks, forest inventory-based approaches have been used all over the world (KURZ, APPS 1999; LISKI et al. 2002; NIZAMI 2012). The output variables of traditional forest inventories are tree height and diameter which can later be converted into tree volume, biomass and C stocks. Stand-level biomass is an aggregation of single-tree biomass. In various process models the relative proportion of biomass components in trees changes after canopy closure (BERNINGER, NIKINMAA 1997), which is the time point-based variable as light conditions change considerably during forest development (OLIVER, LARSON 1996). For example, the relative proportion of foliage biomass decreases, while the relative biomass of stems continues to increase through the tree ontogenetic growth (SATO, MADGWICK 1982). Therefore, the stand-level biomass cannot be measured directly. Instead, an estimation for each tree can be done and then summed to give the total stand estimation (ZIANIS et al. 2005). Estimations of tree biomass can be measured directly by using harvesting methods, but estimation is preferred to avoid damaging the forests (MONTÈS et al. 2000). Using allometric equations (species-specific or generalized; CHAVE et al. 2005) or general volume equation (BROWN, LUGO 1992; NIZAMI et al. 2009; NIZAMI 2012) is a non-destructive approach to calculate total tree biomass. Then the estimation of C stocks can be derived from the biomass using an international standard conversion factor (0.5) and assuming that 50% of the tree biomass has elemental C (DIXON et al. 1994; NIZAMI et al. 2009). After that the ratio of CO₂ to C, i.e. 3.6663, can be used to calculate CO₂ stocks from total C stocks (ANH 2007; TAGUPA et al. 2010).

The rate of sequestered and stored CO₂ of a given tree is related to the tree size such as stem basal area (BA) and total tree height (H). At a sapling stage, the biomass and CO₂ stocking varies among tree species but as trees grew bigger and reached the pole and standard stages, no more significant variation existed in terms of their C stocking capacity (TAGUPA et al. 2010). During the tree growth, BA and H should allometrically relate to biomass accumulation and CO₂ stocks of a tree. However,

the stem wood density might have no significant effect on CO₂ stocks because of a small variation in wood density with the tree growth for a given species (ZHANG 1995). If this is true, we thus hypothesized that CO₂ stocks should vary with the tree growth from sapling to standard stage. Rather than wood density, tree CO₂ stocks should be associated with BA, diameter at breast height (DBH) and H, because of the negligible variation in wood density among the growth stages of a tree.

In absolute terms, soil C stocks are much larger than C sequestered in tree biomass (LAL 2004). A recent calculation of C storage values, including soil C stock estimates, revealed significantly higher estimates in nearly all biomes, including an approximately threefold increase in soil organic C stocks estimates for tropical forests (EGLIN et al. 2011). The moist tropical or sub-tropical forests contained more than 50% of C stocks in soils and almost less than 50% in the vegetation because the dead biomass rapidly decomposes in the warm, humid conditions and the minerals rapidly leach out of tropical forest soils (ROSS 2009). Therefore, the C concentration in soils is highly significant; it is assumed that soils have approximately three times more C than the vegetation and twice more as to that present in the atmosphere (BATJES, SOMBROEK 1997). However, little is currently understood how C stocks partitioned between vegetation and soil in subtropical forests in eastern China.

Evergreen broadleaved forests (EBLFs) are a zonal vegetation type located in subtropical China (SONG, WANG 1995; FENG et al. 1999). In this study, we determined the vegetation-soil C pool and estimated CO₂ stocks in one of the typical EBLFs using different stand indices through forest inventory based approaches. Specifically, we were interested in (1) how the C stock partitions between vegetation and soil pools; (2) whether forest CO₂ stocks relate to the tree growth stage, and (3) whether site indices can be used for predicting CO₂ stocks of trees in EBLFs.

MATERIAL AND METHODS

Study site and plot. The study was conducted in Tiantong National Forest Park (29°48'N, 121°47'E, 200 m a.s.l.), Zhejiang province, China, covering an area of 349 ha. The region has warm and humid subtropical climate with an average annual temperature of 16.2°C, average annual precipitation of 1,374.7 mm (mostly concentrated in the summer), annual average relative humidity is 82 % (shows

low intra-annual variability) and mean annual evaporation is 1,320.1 mm (less than annual precipitation). The soil is mostly mountain yellow-red soil, with the parent material mostly including Mesozoic sedimentary rocks, some acidic igneous rocks, and granite residual weathered material (SONG, WANG 1995).

The zonal vegetation in this region is subtropical EBLFs. In Tiantong National Forest Park, the majority of the EBLFs are *Schima superba* Gardn. et Champ dominated forests, which are considered as sub-climax monsoon EBLFs and have been severely disturbed in the history with only small tracks (approximate 10 ha) of semi-intact forests left around a Buddhist temple (YAN et al. 2009). Since forest age, community structure and plant species composition are similar in this area (YAN et al. 2013), six square plots, each of 10 × 10 m in size, were established to represent the ranges of both community and environmental properties in this study. The tree stratum is 15 to 18 m high and cover percentage is 80~90%, occupied by evergreen broadleaf species. The shrub stratum is

< 4 m in height and coverage is 45–50%. The herb stratum is < 0.5 m in height with coverage being 10 to 30 % and the dominant species are normally ferns. Totally, eighteen woody plant species were found in the studied plots. The species list and their DBH (cm), BA (m²·ha⁻¹), H (m), stem volume (m³·ha⁻¹), wood density (g·cm⁻³) and stem density (trees·ha⁻¹) for each species are shown in detail in Table 1.

Measurements of forest structure and tree biomass. The vegetation C stocks were estimated based on the general volume equation by getting the relationships between tree H, DBH and wood density (NIZAMI et al. 2009; TAGUPA et al. 2010; NIZAMI 2012). For all woody plants in the studied plot, base diameter and DBH were measured for tall trees, while base diameter and 45 cm diameter (D₄₅) were measured for small trees (of less than 1.50 m in height). The H to the top of trees was measured with a telescopic pole for heights up to 15 m, and with a clinometer for heights > 15 m.

The tree individuals of *S. superba* and *Lithocarpus glaber* were classified into three growth stages on the

Table 1. Characteristics of the studied plots in *Schima superba* dominated evergreen broadleaved forest in the Tiantong National Forest Park in eastern China

Species name	DBH (cm)	Height (m)	Basal area (m ² ·ha ⁻¹)	Volume (m ³ ·ha ⁻¹)	Wood density (g·cm ⁻³)	Stem density (tree·ha ⁻¹)	Biomass (t·ha ⁻¹)
<i>Schima superba</i>	14.70 ± 0.56	14.65 ± 0.50	26.95	176.29	0.6	1,063	105.77
<i>Pinus massoniana</i>	33.30 ± 3.20	18 ± 1.32	15.72	121.82	0.5	125	60.91
<i>Lithocarpus glaber</i>	6.81 ± 1.52	7.93 ± 1.73	2.37	15.55	0.5	250	7.77
<i>Rhododendron ovatum</i>	1.39 ± 0.08	2.21 ± 0.06	0.32	0.37	0.6	1,225	0.22
<i>Camellia fraterna</i>	1.03 ± 0.08	1.84 ± 0.10	0.05	0.05	0.6	387	0.027
<i>Eurya muricata</i>	1.30 ± 0.18	2.07 ± 0.19	0.03	0.03	0.6	137	0.019
<i>Eurya rubiginosa</i> var. <i>attenuata</i>	1.19 ± 0.07	1.90 ± 0.06	0.14	0.13	0.6	763	0.08
<i>Symplocos stellaris</i>	1.58 ± 0.22	2.32 ± 0.20	0.10	0.14	0.5	275	0.08
<i>Symplocos sumuntia</i>	1.17 ± 0.06	1.73 ± 0.05	0.12	0.10	0.5	725	0.05
<i>Symplocos heishanensis</i>	1.08 ± 0.06	1.80 ± 0.08	0.07	0.06	0.4	513	0.03
<i>Castanopsis fargesii</i>	1.70 ± 0.60	2.09 ± 0.47	0.50	2.57	0.5	362	1.29
<i>Castanopsis sclerophylla</i>	10.55 ± 1.76	9.77 ± 1.75	1.47	6.74	0.5	100	3.37
<i>Loropetalum chinensis</i>	0.84 ± 0.12	1.93 ± 0.13	0.02	0.01	0.7	162	0.009
<i>Vaccinium mandarinorum</i>	1.06 ± 0.16	1.77 ± 0.07	0.02	0.02	0.8	138	0.01
<i>Helicia cochinchinensis</i>	1.32 ± 0.34	1.92 ± 0.18	0.03	0.03	0.7	113	0.02
<i>Myrica rubra</i>	2.75 ± 1.05	2.96 ± 0.70	0.45	1.74	0.8	187	1.39
<i>Syzygium buxifolium</i>	1.20 ± 0.20	2.31 ± 0.18	0.02	0.02	0.7	88	0.01
<i>Castanopsis carlesii</i>	1.003 ± 0.08	1.78 ± 0.07	0.05	0.04	0.5	437	0.02
Grand total				188.34		7,050	181.075

values of diameter at breast height (DBH) and height for each species are mean ± SE while basal area, volume, and biomass for each species are summarized (total) by all individuals for each species in the plot; basal area and volume were estimated by the ratio of the sample plot area to the area of 1 ha (10,000 m²)

basis of DBH regardless of the age, i.e. sapling stage (< 10 cm DBH), pole stage (10–20 cm DBH) and standard stage (> 20 cm DBH) by following the method standard of TAGUPA et al. (2010) while the other species of the study plot could not be classified into growth stages according to the standard criteria.

Aboveground biomass for each individual tree was calculated by a non-destructive method by multiplying the tree volume by the species-specific dry wood density (BROWN, LUGO 1992; NIZAMI et al. 2009; NIZAMI 2012). The biomass obtained by the volume-density equation was highly consistent ($R^2 = 0.87$; $P < 0.0001$) with biomass obtained by the site and species-specific allometric equation for *Schima superba* (DBH ranges from 3.5 to 25.5 cm; the number of trees cut = 6; YANG et al. 2010) (Fig. 1). Tree volume per hectare for each species was calculated using general Eq. (1) (LU et al. 2003):

$$V = 0.42 \times BA \times H \quad (1)$$

where:

- V – volume of a tree ($\text{m}^3 \cdot \text{ha}^{-1}$),
- 0.42 – fixed general value used for form factor because it can be used in the absence of local equations to estimate the cubic volume of a standing tree (MANGNUSSEN et al. 2004),
- BA – basal area ($\text{m}^2 \cdot \text{ha}^{-1}$) using the formula $BA = \pi (d/2)^2 \times 10^{-4}$ (SMALL et al. 2004),
- H – total tree height (m),
- d – DBH for big trees or D_{45} for small trees (cm).

The per-hectare volume was estimated by the ratio of the sample plot area to the area of 1 ha.

For measuring wood density, wood samples were taken from the trunk of 7–10 trees of the same spe-




Fig. 1. Comparison of the individual tree biomass of *Schima superba* estimated from the volume equation, i.e. tree biomass = tree volume ($\text{m}^3 \cdot \text{ha}^{-1}$) \times species-specific wood density ($\text{g} \cdot \text{cm}^{-3}$) (BROWN, LUGO 1992; BROWN, IVERSON 1992; NIZAMI et al. 2009) and species-specified allometric equation for *Schima superba* (YANG et al. 2010)

cies having different DBH using a 5 mm-diameter increment corer. In the laboratory, the length of the tree core was measured with an electronic vernier calliper, and then the volume of the tree core was calculated. Next, core samples were dried at 75°C in an oven for 72 h to determine dry mass used to calculate the wood density (CHAVE et al. 2006).

It should be noted that the studied plots did not include so much deadwood and debris because of clearing by the forest manager every year for a landscape view. Therefore, we decided to take the soil and vegetation as major C pools by ignoring dead woody debris.

Calculation of forest C and CO_2 stocks. Total C stock in each species was determined by multiplying the total tree biomass by a conversion factor (0.5), which is representative of the average C content in tree biomass. This conversion factor shows that 50% of the total tree biomass is equal to elemental C (BROWN, LUGO 1982; DIXON et al. 1994; NIZAMI et al. 2009).

The CO_2 stock in tree was computed using Eq. 2 (TAGUPA et al. 2010).

$$\text{CO}_2 \text{ stock} = C_{\text{veg}} \times 3.6663 \quad (2)$$

where:

- C_{veg} – C stocks in vegetation ($\text{t} \cdot \text{ha}^{-1}$),
- 3.6663 – universal conversion factor of C content values to CO_2 values.

The ratio of CO_2 to C is determined by atomic mass (ANH 2007; CFS 2009; SUPERAK 2010). Furthermore, ANH (2007) also suggested on the basis of an experimental approach that the CO_2 amount which vegetation absorbs and O_2 are harmonized in the atmosphere and can be calculated by means of the chemistry equation, i.e. $\text{CO}_2 = \text{C} + \text{O}_2$ and thus $\text{CO}_2 = 3.6663\text{C}$.

CO_2 stocks in different tree components for each individual were estimated using tree components specific allometric Eqs. 3–6, developed by HUYNH and ANH (2008) for *Schima superba* dominated EBLFs in Vietnam:

$$\text{CO}_2 \text{ stem} = 6.15398 + 1.02468 \times \text{Ln}(V) \quad (3)$$

$$\text{CO}_2 \text{ bark} = 4.11447 + 1.06381 \times \text{Ln}(V) \quad (4)$$

$$\text{CO}_2 \text{ branch} = -4.11248 + 2.70337 \times \text{Ln}(d) \quad (5)$$

$$\text{CO}_2 \text{ leaf} = -2.941 + 1.72414 \times \text{Ln}(d) \quad (6)$$

where:

- CO_2 – stocks in each tree component, converted to $\text{t} \cdot \text{ha}^{-1}$;
- V – volume of a tree stem (m^3),
- d – DBH for big trees or D_{45} for small trees (cm).

However, to avoid the overestimation of the total CO₂ content, the amount of CO₂ in bark and stem was considered as an overall percentage, i.e. CO₂ in overbark stem (TAGUPA et al. 2010).

Measurement of soil carbon stock. Ten soil samples were taken with a metal corer from randomly chosen spots in each of the studied plot (0–20 cm soil layer). The litter layer was removed before the soil was taken. The soil bulk density was determined using a steel corer of the known volume. Before weighing, approximately 5 soil cores per plot were placed in a 105°C oven for > 48 hours. Bulk density (g·cm⁻³) was calculated by dividing the oven dry weight (g) of the soil sample by the volume of the soil sample in the steel corer (cm³). Then the remaining soil samples were air-dried for 30 days, and passed through a 2-mm sieve to determine the total C concentration using the oil bath-K₂CrO₇ titration method (NELSON, SOMMERS 1975). Finally, C stock in the top 0–20 cm soils was estimated using BATJES (1996) formula.

Data analysis. The data analysis included three steps. The first step was to determine relationships between H, DBH, BA, tree volume and CO₂ stocks of all plant species using linear regression methods. In the second step, one-way ANOVA with the Least Significant Difference (LSD) test was used to find out the significant variance of CO₂ stocks between tree

growth stages while regression ANOVA was used to explore the significant variation of CO₂ stocks with the response of DBH in each growth stage of *Schima superba* and *Lithocarpus glaber*. In the third step, the stepwise multiple regression analysis was used to find out whether CO₂ stocks were affected by different tree variables (DBH, H and wood density) of all studied species in the sampling plots. All statistical tests were considered significant at the $P < 0.05$ level.

RESULTS

Forest biomass

H in the studied forest increased with increasing DBH. Among all species, on average basis *Pinus massoniana* had greater H (18 m) and DBH (33.3 cm) than *Schima superba* with 14.65 m H and 14.70 cm DBH (Table 1). For most species, H and DBH showed a positive linear relationship. Stem volume increased linearly with increasing BA ($r = 0.98$) for all species.

Tree biomass was related to BA significantly and linearly for each species (Table 2). Total tree biomass was greater in *Schima superba* (105.77 t·ha⁻¹) than in *Pinus massoniana* (60.911 t·ha⁻¹). In the studied forest, grand total tree biomass being cal-

Table 2. Linear regression analysis for tree biomass (TB) against basal area (BA)

Species name	Regression equation	R ²	P
<i>Schima superba</i>	TB = -0.151 + 73.356 (BA)	0.875	0.000
<i>Pinus massoniana</i>	TB = -0.578 + 70.697 (BA)	0.904	0.000
<i>Lithocarpus glaber</i>	TB = -0.001 + 54.568 (BA)	0.934	0.000
<i>Rhododendron ovatum</i>	TB = -0.001 + 14.822 (BA)	0.935	0.000
<i>Camellia fraterna</i>	TB = 0.000 + 12.162 (BA)	0.914	0.000
<i>Eurya muricata</i>	TB = 0.000 + 12.535 (BA)	0.939	0.000
<i>Eurya rubiginosa</i> var. <i>attenuata</i>	TB = 0.000 + 11.662 (BA)	0.964	0.000
<i>Symplocos stellaris</i>	TB = -0.001 + 15.891 (BA)	0.926	0.000
<i>Symplocos sumuntia</i>	TB = 0.000 + 8.248 (BA)	0.942	0.000
<i>Symplocos heishanensis</i>	TB = -0.00009 + 6.388 (BA)	0.886	0.000
<i>Castanopsis fargesii</i>	TB = -0.009 + 51.722 (BA)	0.979	0.000
<i>Castanopsis sclerophylla</i>	TB = -0.039 + 41.846 (BA)	0.906	0.000
<i>Loropetalum chinensis</i>	TB = 0.000 + 13.022 (BA)	0.935	0.000
<i>Vaccinium mandarinorum</i>	TB = 0.000 + 11.677 (BA)	0.978	0.000
<i>Helicia cochinchinensis</i>	TB = 0.000 + 15.006 (BA)	0.996	0.000
<i>Myrica rubra</i>	TB = -0.006 + 54.293 (BA)	0.997	0.000
<i>Syzygium buxifolium</i>	TB = 0.00004 + 11.230 (BA)	0.947	0.000
<i>Castanopsis carlesii</i>	TB = 0.000 + 8.703 (BA)	0.976	0.000

significant at $P < 0.05$

culated by summing each individual tree of all species was 181.07 t·ha⁻¹.

Forest C and CO₂ stocks

Assuming that 50% of the vegetation biomass is C, the greater amount of total C stocks was found in the species *Schima superba* (52.88 t·ha⁻¹), followed by *Pinus massoniana* (30.45 t·ha⁻¹). Total vegetation C stocks in the studied forest were 90.53 t·ha⁻¹ (Table 3). Total soil C stocks at the 0–20 cm surface soil were 116.24 t·ha⁻¹ (Fig. 2) having 1.3 ± 0.01 (g·cm⁻³) bulk density. Totally, the overall C stocks on the ecosystem (vegetation + soil) level were 206.77 t·ha⁻¹, with 43.79% in vegetation, while with 56.21% in the surface soil (Fig. 2).

The total amount of CO₂ stocks in the studied forest was 331.87 t·ha⁻¹. The amount of CO₂ sequestered among tree components was largest in stems, intermediate in branches, and lowest in leaves (Table 3). Considering particular species, the greater amount of total CO₂ stocks was found in *Schima superba* (193.90 t·ha⁻¹) and *Pinus massoniana* (111.66 t·ha⁻¹).

Relationship between CO₂ stock capacity and tree growth

To test whether the CO₂ stocking capacity depends on tree ontogeny, the two most abundant species *Schima superba* and *Lithocarpus glaber* were classified into three developmental stages on the basis of DBH; sapling stage (< 10 cm DBH), pole stage (10–20 cm DBH), standard stage (> 20 cm DBH). One-way ANOVA with the LSD test showed that the CO₂ stock var-




Fig. 2. Carbon stocks in each type of the aboveground vegetation and the 0–20 cm surface soil in *Schima superba* dominated evergreen broadleaved forest in the Tiantong National Forest Park in eastern China

ies significantly ($P < 0.05$) between the growth stages (Fig. 3). Furthermore, according to the regression ANOVA result, significant variation in CO₂ stocking capacity with DBH variation exists within sapling stage ($P = 0.037$) and pole stage ($P = 0.00$) while no significant ($P = 0.13$) variation exist within standard stage (Table 4).

Relationships between CO₂ stocks and stand indices

The regression equation of each species showed that CO₂ stocks increase with the increasing diameter of the tree. Wood density, however, did not affect the C stocks. Also, P -values for each species showed that there was a significant relationship between CO₂ stocks and DBH of trees ($P < 0.05$, Table 5).

In each unit area or each forest stand, it is necessary to calculate CO₂ stocks to find out which stand indices influence the effectiveness of predicting CO₂ stocks. The stepwise regression analysis showed that DBH and H influenced the effectiveness of predicting CO₂ stocks and wood density was excluded from the regression model [i.e. Eq. 7, where $R^2 = 0.80$, $P < 0.001$, DBH – diameter breast high (cm), H – total tree height (m)].

$$\text{CO}_{2 \text{ stock}} = -0.27 + 0.35 \text{ DBH} - 0.14 \text{ H} \quad (7)$$

DISCUSSION

Effectiveness of stand indices in predicting forest CO₂ stocks

According to HUY and ANH (2008) findings the absorbed CO₂ amount depends on the species, DBH and H but they did not assess the role of wood density and BA in terms of their effect on C stocks for the specific forest type. In this study, wood density and BA were taken as indicators for predicting the CO₂ stocking capacity in EBLFs. The result showed that BA strongly affects the CO₂ stocks because of a change in DBH and stem area occupied by the tree during growth. This result is consistent with the findings of HUY and ANH (2008) that CO₂ has a firm relationship with DBH which has the advantage to identify the CO₂ amount sequestered in each forest stand. Also, this result was quite close to the finding of TAGUPA et al. (2010) that the variation in the amount of CO₂ sequestered and stored in the trees within a forest stand is affected greatly by the stand density of trees. The estimation of per-hectare tree volume using BA as a function is more significant as compared

Table 3. Estimates of C and CO₂ stocks based on tree components in *Schima superba* dominated evergreen broad-leaved forest in the Tiantong National Forest Park in eastern China

Species name	Carbon stock (t·ha ⁻¹)	CO ₂ stock capacity (t·ha ⁻¹)			
		stems	branches	leaves	total
<i>Schima superba</i>	52.89	139.61	50.41	3.883.87	193.90
<i>Pinus massoniana</i>	30.45	80.4	29.03	2.23	111.66
<i>Lithocarpus glaber</i>	3.89	10.27	3.7	0.29	14.26
<i>Rhododendron ovatum</i>	0.11	0.3	0.1	0.008	0.408
<i>Camellia fraterna</i>	0.01	0.03	0.01	0.001	0.041
<i>Eurya muricata</i>	0.01	0.02	0.008	0.0006	0.0286
<i>Eurya rubiginosa</i> var. <i>attenuata</i>	0.04	0.1	0.03	0.002	0.132
<i>Symplocos stellaris</i>	0.04	0.1	0.03	0.002	0.132
<i>Symplocos sumuntia</i>	0.02	0.07	0.02	0.001	0.091
<i>Symplocos heishanensis</i>	0.01	0.03	0.01	0.0008	0.0408
<i>Castanopsis fargesii</i>	0.64	1.7	0.61	0.04	2.35
<i>Castanopsis sclerophylla</i>	1.69	4.44	1.6	0.12	6.16
<i>Loropetalum chinensis</i>	0.004	0.01	0.004	0.0003	0.0143
<i>Vaccinium mandarinorum</i>	0.005	0.01	0.006	0.0004	0.0164
<i>Helicia cochinchinensis</i>	0.01	0.03	0.01	0.0008	0.0408
<i>Myrica rubra</i>	0.70	1.83	0.67	0.05	2.55
<i>Syzygium buxifolium</i>	0.005	0.01	0.005	0.0003	0.0153
<i>Castanopsis carlesii</i>	0.01	0.02	0.01	0.0007	0.0307
Grand total	90.534	238.98	86.263	6.6279	331.8709

C and CO₂ stocks were obtained from the summation of all individuals for each species in the plot

with using DBH only because BA is the area of a given section of land that is occupied by the cross-sections of tree trunks and stems at their base, which can be further applied to more precise estimation of tree biomass and then CO₂ stocks per hectare. In this study, wood density had no significant effect on CO₂ stocks in a mixed forest stand because wood density have a negative correlation with DBH ($r = -0.125$, $P < 0.05$) and H ($r = -0.191$, $P < 0.05$) growth (ROCHON et al. 2007), but on the other hand, wood density is a significant parameter for predicting tree biomass and C stocks (NOGUERIA et al. 2005). Therefore, it is important to mention here that the role of wood density cannot be ignored in tree biomass estimation but it has no significant effect on the CO₂ stocks with tree growth in a mixed forest stand.

Change in CO₂ stocking capacity with tree growth

C sequestration varies from forest type to forest type and for the growth of forest which potentially relies on the tree diameter size class, i.e. growth stages according to DBH (TERAKUNPISUT et al. 2007).

The CO₂ stocks were dependent on the amount of biomass of trees, specifically, on the variables trunk diameter and H. This confirmed the result of TERAKUNPISUT et al. (2007) and TAGUPA et al. (2010), who mentioned that the C stocking potential in different forest types tends to be correlated with DBH and H. The oneway ANOVA result of *Schima superba* and *Lithocarpus glaber* species showed that the biomass and CO₂ stock varied between growth stages. Furthermore, within sapling and pole stages, CO₂ stocks varied significantly with DBH growth but as trees grow bigger and reach standard sizes,

Table 4. Results of regression ANOVA for the variability of CO₂ stocks induced by individual trees differing in DBH within each growth stage (sapling, pole and standard stages)

Growth stage	CO ₂ stock (t·ha ⁻¹)	F	P
Sapling	6.17	3.328	0.037
Pole	132.17	5.469	0.00
Standard	69.85	76.838	0.13

CO₂ stock data – summation of *Schima superba* and *Lithocarpus glaber*, significant at $P < 0.05$

Table 5. Results of linear regression analysis for CO₂ stocking capacity (Y) against stem diameter at breast height (X) for large trees or diameter at 45 cm height (X) for small trees in *Schima superba* dominated evergreen broadleaved forest in the Tiantong National Forest Park in eastern China

Species name	Regression equation	R ²	P
<i>Schima superba</i>	$Y = -2.550 + 0.329 (X)$	0.819	0.000
<i>Pinus massoniana</i>	$Y = -10.886 + 0.662 (X)$	0.855	0.000
<i>Lithocarpus glaber</i>	$Y = -0.193 + 0.133 (X)$	0.890	0.000
<i>Rhododendron ovatum</i>	$Y = -0.006 + 0.007 (X)$	0.794	0.000
<i>Camellia fraternal</i>	$Y = -0.003 + 0.004 (X)$	0.798	0.000
<i>Eurya muricata</i>	$Y = -0.004 + 0.005 (X)$	0.877	0.000
<i>Eurya rubiginosa</i> var <i>attenuata</i>	$Y = -0.004 + 0.005 (X)$	0.877	0.000
<i>Symplocos stellaris</i>	$Y = -0.008 + 0.009 (X)$	0.739	0.000
<i>Symplocos sumuntia</i>	$Y = -0.002 + 0.003 (X)$	0.849	0.000
<i>Symplocos heishanensis</i>	$Y = -0.001 + 0.002 (X)$	0.836	0.000
<i>Castanopsis fargesii</i>	$Y = -0.264 + 0.165(X)$	0.964	0.000
<i>Castanopsis sclerophylla</i>	$Y = -0.417 + 0.109(X)$	0.759	0.005
<i>Loropetalum chinensis</i>	$Y = -0.002 + 0.003(X)$	0.829	0.000
<i>Vaccinium mandarinorum</i>	$Y = -0.002 + 0.004(X)$	0.898	0.000
<i>Helicia cochinchinensis</i>	$Y = -0.007 + 0.009(X)$	0.913	0.000
<i>Myrica rubra</i>	$Y = -0.131 + 0.109(X)$	0.985	0.000
<i>Syzygium buxifolium</i>	$Y = -0.002 + 0.004(X)$	0.970	0.000
<i>Castanopsis carlesii</i>	$Y = -0.002 + 0.003(X)$	0.889	0.000

significant at $P < 0.05$

no significant variation exists any more in terms of their CO₂ stocking because the growth rate (change in DBH) will gradually slow down for mature trees (TERAKUNPISUT et al. 2007). The factors such as wood density, H, BA and DBH may have interplayed, giving complementary effects. The lower wood density values may be compensated by greater H and DBH, and vice versa.

Carbon stock pattern in typical subtropical forest

Estimating C stocks by species is an important trend and should be studied to a greater extent especially when the tree species composition might change in a mixed forest stand (KELLOMÄKI, KOLSTRÖM 1992). Therefore, the study of C stocks between vegetation and soils is more important to clear the complex relationship for each forest type and location. A linear regression equation was developed for the tree biomass of each species as a function of BA, showing that the tree biomass increases with the increasing stem diameter (NIZAMI et al. 2009). In this study, the result is quite close to the findings of YANG et al. (2010), who reported that the total biomass in Tiantong *Schi-*

ma superba community was $225.3 \pm 30.1 \text{ t}\cdot\text{ha}^{-1}$. The overall C stock (vegetation plus soil) in subtropical EBLFs was $206.77 \text{ t}\cdot\text{ha}^{-1}$ which consists of 43.79% in vegetation biomass and 56.21% in the topsoil. Additionally, our results are quite close to the findings of Ross (2009), who reported that the subtropical forest contained more than 50% (i.e. up to 50.5%) C in soil and less than 50% (i.e. up to 49.5%) C in vegetation. The overall C stocks are relatively close to the estimate by DIXON et al. (1994), who reported that the Asian tropical forest holds a C density of $132\text{--}174 \text{ t}\cdot\text{ha}^{-1}$. BROWN and LUGO (1984) estimated that the tropical forests of Bangladesh hold approximately $55\text{--}90 \text{ t}\cdot\text{ha}^{-1}$ of C in forest ecosystems. Overall, soil C stocks are the interactive result of climate, soil properties, litter production by vegetation, and litter quality; the more productive and dense vegetation in the site, the more litter is fed into soil (LISKI 1995). In subtropical EBLFs, the C stocks in soils are a little higher than those in vegetation because of the rapid decomposition of dead biomass (YAN et al. 2007), higher litter production due to the dense mixed forest stand (approx; plant to plant and row to row distance was $3 \times 3 \text{ m}$, respectively, in the study plot), and moist and humid climate (YAN et al. 2008).

CONCLUSIONS

The significant amount of C stocking in the typical EBLF in the Tiantong National Forest Park of eastern China showed the potential and significant C stocks by trees. Despite of the tree age, the bigger trees, particularly at their standard sizes, have the greatest C stocks, sequestered the greatest amount of CO₂, but have a low ability to sequester more CO₂ from the atmosphere for future because of the slow growth in trunk diameter as compared to the trees of pole and sapling sizes. Assuming that the trees are allowed to grow and are not cut for any purpose at all, they continue to provide the safety C sinks for the adverse effects of climate change. As the C pool structure changes due to a change in the forest type, class and location, therefore this study is important to estimate C stocks and predict CO₂ stocks from stand indices in EBLFs which serve as a scientific basis for sustainable forestry operations, rational utilization of forest resources and global warming reduction in EBLFs in subtropical regions of China.

Acknowledgements

We are grateful to Prof. EN-RONG YAN for insightful comments during the development of this work.

References

- ANH T.P. (2007): Forecasting CO₂ absorptivity capacity on natural broad-leaved evergreen forests in tuy duc district, Daknong province. [Summary of Master's Thesis.] Đak Läk, Ministry of Agriculture and Rural Development (MARD), Forestry University-Tay Nguyen University in Vietnam: 24.
- BATJES N.H. (1996): Total carbon and nitrogen in the soils of the world. *European Journal of Soil Science*, **47**: 151–163.
- BATJES N.H., SOMBROEK W.G. (1997): Possibilities for carbon sequestration in tropical and subtropical soils. *Global Change Biology*, **3**: 161–173.
- BERNINGER F., NIKINMAA E. (1997): Implications of varying pipe model relationships on Scots Pine growth in different climates. *Functional Ecology*, **11**: 146–156.
- BROWN S., LUGO A.E. (1982): The storage and production of organic matter in tropical forests and their role in the global carbon cycle. *Biotropica*, **14**: 161–187.
- BROWN S., LUGO A.E. (1984): Biomass of tropical forests: A new estimate based on forest volumes. *Journal of Forest Science*, **22**: 1290–1293.
- BROWN S., LUGO A.E. (1992): Above ground biomass estimates for tropical moist forests of the Brazilian Amazon. *Interciencia*, **17**: 8–18.
- CHAVE J., ANDALO C., BROWN S., CAIRNS M.A., CHAMBERS J.Q., EAMUS D., FÖLSTER H., FROMARD F., HIGUCHI N., KIRA T., LESURE J.P., NELSON B.W., OGAWA H., PUIG H., RIÉRA B., YAMAKURA T. (2005): Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, **145**: 87–99.
- CFS (2009): CarbonFix Standard. Available at <http://www.carbonfix.info/chameleon/outbox/public/216/Example-Calculations.pdf?PHPSESSID=cyqkigazki>
- CHAVE J., MULLER-LANDAU H., BAKER T., EASDALE T., TEER-STEERGE H.T., WEBB C.O. (2006): Regional and phylogenetic variation of wood density across 2,456 neotropical tree species. *Ecological Applications*, **16**: 2356–2367.
- DIXON R.K., BROWN S., SOLOMON R.A., TREXLER M.C., WISNIEWSKI J. (1994): Carbon pools and flux of global forest ecosystems. *Science*, **263**: 185–190.
- EGLIN T., CIAIS P., PIAO L.S., BARRE P., BELASSEN V., CADULE P., CHEN C., GASSER T., REICHSTEIN M., SMITH P., SAUER J.T., NORMAN M.J., SIVAKUMAR K.V.M. (2011): Overview on Response of Global Soil Carbon Pools to Climate and Land-Use Changes. In: SAUER T.J. et al. (eds): *Sustaining Soil Productivity in Response to Global Climate Change*: 183–199.
- FENG Z.W., WANG X.K., WU G. (1999): *The Biomass and Productivity of Forest Ecosystem in China*. Beijing, China Science Press: 241. (in Chinese)
- HUY B., ANH T.P. (2008): Estimating CO₂ sequestration in natural broad-leaved evergreen forest in Vietnam. *Asia-Pacific Agroforestry Newsletter*, **32**: 7–10.
- JANA B.K., BISWAS S., MAJUMDER M., ROY P.K., MAZUMDAR A. (2011): Carbon sequestration rate and aboveground biomass carbon potential of three young species in lower Gangetic plain. *Journal of Environmental Science and Engineering*, **53**: 299–308.
- KELLOMÄKI S., KOLSTRÖM M. (1992): Simulation of tree species composition and organic matter accumulation in Finnish boreal forests under changing climatic conditions. *Vegetatio*, **102**: 47–68.
- KURZ W.A., APPS M.J. (1999): Developing Canada's national forest carbon monitoring, accounting and reporting system to meet the reporting requirements of the Kyoto protocol. *Mitigation and Adaptation Strategies for Global Change*, **11**: 33–43.
- LAL R. (2004): Soil carbon sequestration impacts on global climate change and food security. *Science*, **304**: 1623–1627.
- LISKI J. (1995): Variation in soil organic carbon and thickness of soil horizons within a boreal forest stand – effect of trees and implications for sampling. *Silva Fennica*, **29**: 255–266.
- LISKI J., PERRUCHOD D., KARJALAINEN T. (2002): Increasing carbon stocks in the forest soils of Western Europe. *Forest Ecology and Management*, **169**: 159–175.
- LU D., MAUSEL P., BRONDI'ZIO E., MORAN E. (2003): Classification of successional forest stages in the Brazilian Amazon basin. *Forest Ecology and Management*, **181**: 301–312.

- MANGNUSSEN S., REED D. (2004): Modeling for estimation and monitoring. Available at <http://www.fao.org/forestry/17109/en/>
- MATTHEWS E., PAYNE R., ROHWEDER M., MURRAY S. (2000): Forest ecosystem: carbon storage sequestration. *Global Climate Change Digest*, **12**: 157–180.
- MONTÈS N., GAUQUELIN T., BADRI W., BERTAUDIÈRE V., ZAOUÏ E.H. (2000): A nondestructive method for estimating above-ground forest biomass in threatened woodlands. *Forest Ecology and Management*, **130**: 37–46.
- MOONEY H., ROY J., SAUGIER B. (2001): *Terrestrial Global Productivity*. San Diego, Academic Press: 573.
- NELSON D.W., SOMMERS L.F. (1975): A rapid and accurate method for estimating organic carbon in soil. *Proceedings of the Indian Academy of Sciences*, **84**: 456–462.
- NIZAMI M.S., MIRZA N.S., LIVESLEY S., ARNDT S., FOX C.J., KHAN A.I., MAHMOOD T. (2009): Estimating carbon stocks in sub-tropical pine (*Pinus roxburghii*) forests of Pakistan. *Pakistan Journal of Agricultural Sciences*, **4**: 266–270.
- NIZAMI M.S. (2012): Assessment of the carbon stocks in sub tropical forests of Pakistan for reporting under Kyoto protocol. *Journal of Forest Research*, **23**: 377–384.
- NOGUEIRA E.M., NELSON B.W., FEARNISID P.M. (2005): Wood density in dense forest in central Amazonia, Brazil. *Forest Ecology and Management*, **208**: 261–286.
- NOWAK D.J., CRANE D.E. (2002): Carbon storage and sequestration by urban trees in the United States. *Environmental Pollution*, **116**: 381–389.
- OLIVER C.D., LARSON B.C. (1996): *Forest Stand Dynamics*. New York, John Wiley & Sons: 518.
- PAN Y., BIRDSEY R.A., FANG J., HOUGHTON R., KAUPPI P.E., KURZ W.A., PHILLIPS O.L., SHVIDENKO A., LEWIS S.L., CANADELL J.G., CIAIS P., JACKSON R.B., PACALA S.W., MCGUIRE A.D., PIAO S., RAUTIAINEN A., SITCH S., HAYES D. (2011): A large and persistent carbon sink in the world's forests. *Science*, **333**: 988–993.
- ROCHON C., MARGOLIS H.A., WEBER J.C. (2007): Genetic variation in growth of *Guazuma crinita* (Mart.) trees at an early age in the Peruvian Amazon. *Forest Ecology and Management*, **243**: 291–298.
- ROSS W.G. (2009): Carbon sequestration in forests. Available at <http://www.fas.org/sgp/crs/misc/RL31432.pdf>
- SATOO T., MADGWICK H.A.I. (1982): *Forest Biomass*. Hague, Martinus Nijhoff/Dr W. Junk Publishers: 152.
- SMALL A., MARTIN T.G., KITCHING R.L., WONG K.M. (2004): Contribution of tree species to the biodiversity of a 1 ha old World rainforest in Brunei, Borneo. *Biodiversity and Conservation*, **13**: 2067–2088.
- SONG Y.C., WANG X.R. (1995): *Vegetation and flora of Tiantong national forest park Zhejiang province*. Shanghai, Shanghai Scientific & Technological Literature Publishing House: 16. (in Chinese with English abstract)
- SUPERAK C. (2010): Rural tree decline in Tasmania's Midlands: stand structure, substrate geology, and carbon content analysis. Available at http://digitalcollections.sit.edu/isp_collection/866
- TAGUPA C., LOPEZ A., CAPERIDA F., PAMUNAG G., LUZADA. A. (2010): Carbon dioxide (CO₂) sequestration capacity of Tampilisan Forest. *E-International Scientific Research Journal*, **2**: 182–191.
- TERAKUNPISUT J., GAAJASENI N., RUANKAWE N. (2007): Carbon sequestration potential in aboveground biomass of Thong Pha Phum National Forest, Thailand. *Applied Ecology and Environmental Research*, **5**: 93–102.
- YAN E.R., YANG X.D., CHANG S.X., WANG X.H. (2013): Plant trait-species abundance relationships vary with environmental properties in subtropical forests in eastern China. *PLoS ONE*, **8**, e61113. doi:10.1371/journal.pone.0061113
- YAN E.R., WANG X.H., GUO M. et al. (2009): Temporal patterns of net soil N mineralization and nitrification through secondary succession in the subtropical forests of eastern China. *Plant and Soil*, **320**: 181–194.
- YAN E.R., WANG X.H., ZHOU W. (2008): Characteristics of litter fall in relation to soil nutrients in mature and degraded evergreen broad-leaved forests of Tiantong, Eastern China. *Chinese Journal of Plant Ecology*, **32**: 1–12.
- YAN E.R., WANG X.H., HUANG J.J., ZENG F.R., GONG L. (2007): Long-lasting legacy of forest succession and forest management: Characteristics of coarse woody debris in an evergreen broad-leaved forest of Eastern China. *Forest Ecology and Management*, **252**: 98–107.
- YANG T.H., SONG K., DA L.J. LI X.P., WU J.P. (2010): The biomass and aboveground net primary productivity of *Schima superba*–*Castanopsis carlesii* forests in east China. *Science China Life Sciences*, **53**: 811–821.
- ZHANG S.Y. (1995): Effect of growth rate on wood specific gravity and selected mechanical properties in individual species from distinct wood categories. *Wood Science and Technology*, **29**: 451–465.
- ZIANIS D., MUUKKONEN P., MÄKIPÄÄ R., MENCUCCINI M. (2005): *Biomass and Stem Volume Equations for tree Species in Europe*. *Silva Fennica Monographs Vol. 4* Tampere, The Finnish Society of Forest Science: 63.

Received for publication February 15, 2014

Accepted after corrections May 12, 2014

Corresponding author:

ARSHAD ALI, Department of Environmental Sciences, East China Normal University, Shanghai 200241, P. R. China; e-mail: arshadforester@gmail.com
