

# Influence of salt stress on ecophysiological parameters of *Periploca sepium* Bunge

J.K. Sun<sup>1,2</sup>, T. Li<sup>2</sup>, J.B. Xia<sup>2</sup>, J.Y. Tian<sup>2</sup>, Z.H. Lu<sup>1,2</sup>, R.T. Wang<sup>1</sup>

<sup>1</sup>*Institute of Restoration Ecology, China University of Mining and Technology, Beijing, P.R. China*

<sup>2</sup>*Shandong Provincial Key Laboratory of Eco-Environmental Science for Yellow River Delta, Binzhou University, Binzhou, P.R. China*

## ABSTRACT

An experiment was carried out to investigate the effect of salt stress on *Periploca sepium* Bunge seedlings using three levels of salinity, 50 mmol/L, 100 mmol/L, and 200 mmol/L sodium chloride (NaCl) solution. The results showed that growth parameters and net photosynthetic rate ( $P_n$ ), stomatal conductance ( $G_s$ ) of *Periploca sepium* Bunge were enhanced under low salinity levels (50 mmol/L NaCl), which reduced strongly with increasing salinity levels. Under 100 mmol/L NaCl and 200 mmol/L NaCl stress, the decline of  $P_n$  was mainly caused by non-stomatal factors. The water use efficiency (WUE), apparent light use efficiency (LUE), carboxylation efficiency (CUE) were enhanced under low salinity levels (50 mmol/L NaCl), the maximum value of WUE was observed at 100 mmol/L NaCl, the minimum value of WUE was observed at 200 mmol/L NaCl, the LUE, CUE were reduced by 52% and 47%, at 200 mmol/L NaCl, respectively, compared to control. Activities of the antioxidative enzymes superoxide dismutase (SOD), peroxidase (POD), and catalase (CAT) were enhanced by low salinity treatment (50 mmol/L NaCl), but CAT activity decreased at 200 mmol/L NaCl stress. Malondialdehyde (MDA) was non-significant compared to the control under low salinity levels (50 mmol/L NaCl), the maximum value was observed at 200 mmol/L NaCl. These results suggest a possibility to improve saline soil utilization of *Periploca sepium* Bunge in Yellow River Delta region.

**Keywords:** salinity; growth; gas exchange; protective enzyme; lipid peroxidation

High salinity is the most widespread abiotic stress and constitutes the most stringent factor in limiting plant distribution and productivity (Yildirim et al. 2009, Qin et al. 2010). Salt stress has various effects on plant ecophysiological processes such as changes in plant growth (Maeda and Nakazawa 2008), protective enzyme activities (Wu et al. 2010), properties of photosynthesis (Ashraf and Shahbaz 2003, Kao et al. 2003), mineral distribution (Yang et al. 2008a), membrane permeability (Dogan et al. 2010), and so on.

The Yellow River Delta differs in many respects from other major deltaic systems of the world.

The tremendous annual sediment load combined with many dramatic changes in the course of the river resulted in a complex and episodic deltaic history (Bornhold et al. 1986). It is the only large delta in China to undergo extensive development which is characterized by extensive coverage of saline soils (Fang et al. 2005). Salt stress is a major abiotic stress problem in the region. Utilization of salt-tolerant plants is expected to be an effective method to improve saline soil. Hence, better understanding of the mechanisms that enable plants to adapt to salt stress is necessary for exploiting saline soil (Patel and Pandey 2008).

---

Supported by the Eleventh Five-year Plan for Supporting Science and Technology of China, Project No. 2006BAC01A13; by the National Natural Science Foundation of China, Project No. 30770412; by the Natural Science Foundation of Shandong Province, Project No. Y2006D01; by the Science and technology projects of Shandong Province, Projects No. 2008GG3NS07005 and 2008GG10006024, and by the Environmental Protection Key Projects of Shandong Province, Project No. 2006 006-2.

*Periploca sepium* Bunge is a woody climbing vine belonging to the family Asclepiadaceae. It is widely distributed in northeastern area, northern area, northwestern area and southwestern area of China. The root bark of *Periploca sepium* Bunge has been used as a traditional Chinese herbal medicine for more than two thousand years in China (Ma et al. 2008). To our knowledge, no information is available on salt tolerance for *Periploca sepium* Bunge, thus it is necessary to study its mechanisms. In this research, the effect of sodium chloride on growth, photosynthetic gas exchange, and protective enzyme activities of *Periploca sepium* Bunge were studied so as to provide some fundamental bases for vegetation restoration in yellow river delta region.

## MATERIALS AND METHODS

**Plant materials.** The seeds of *Periploca sepium* Bunge were originally collected from the National Nature Reserve of Shell Islands and Wetlands in Binzhou, P.R. China. The experiment was conducted in a greenhouse supplied with natural sunlight in the Shandong Provincial Key Laboratory of Eco-environmental Science for Yellow River Delta, Binzhou University, Binzhou, P.R. China from March to June 2008. The greenhouse temperatures were from 28°C to 12°C, relative humidity ranged from 35% to 65%. The experiment was set as four treatments including control, each treatment with three replications. The pots of 15 cm diameter lined with polythene sheets were filled with 2.0 kg of well washed pure shell sand. Thirty seeds were sown in each pot. At 15 days after germination, plants were thinned to 6 plants per pot. Full Hoagland's nutrient solution was applied to each pot every week before salt treatment.

**Stress treatment.** The salt stress was applied on the 30<sup>th</sup> day after germination. *Periploca sepium* Bunge seedlings uniform in size, were subjected to salt stress by adding NaCl solutions of four concentrations, 0 (control), 50, 100, and 200 mmol/L NaCl. The final salt concentration was progressively adjusted with increasing NaCl solutions (50 mmol/L per day). At the same time, the plants were irrigated with fresh water every three days to maintain sand water.

**Indices measurement.** Leaf gas exchange measurements were conducted on three to five medial leaves for each treatment on 61<sup>st</sup> day after salt treatment. Net photosynthetic rate ( $P_n$ ,  $\mu\text{mol}/\text{m}^2/\text{s}$ ) transpiration rate ( $T_r$ ,  $\text{mmol}/\text{m}^2/\text{s}$ ), stomatal conductance ( $G_s$ ,  $\text{mmol}/\text{m}^2/\text{s}$ ), intercellular  $\text{CO}_2$

concentration ( $C_i$ ,  $\mu\text{mol}/\text{mol}$ ) and environmental factors such as photosynthetically active radiation (PAR,  $\mu\text{mol}/\text{m}^2/\text{s}$ ), atmospheric  $\text{CO}_2$  concentrations ( $C_a$ ,  $\mu\text{mol}/\text{mol}$ ), temperature ( $T_a$ , °C) and atmospheric humidity (RH, %) et al. were measured simultaneously with an LI-6400 Portable Photosynthesis System in the morning between 9:00 a.m. and 10:00 a.m. In the native environment, with the following specifications and adjustments: the photosynthetic active radiation  $1000.8 \pm 9.2 \text{ mmol}/\text{m}^2/\text{s}$ , ambient  $\text{CO}_2$  concentration  $390.5 \pm 5.2 \mu\text{mol}/\text{mol}$ , air temperature  $24.7 \pm 1.6^\circ\text{C}$ , and relative humidity  $32.4 \pm 3.6\%$ . The following indicators were calculated: stomatal conductance limitation ( $L_s$ ) =  $1 - C_i/C_a$  (Berry and Downton 1982), water use efficiency (WUE,  $\mu\text{mol}/\text{mmol} = P_n/T_r$  (Nijs et al. 1997), apparent light use efficiency (LUE) =  $P_n/\text{PAR}$  (Long et al. 1993), carboxylation efficiency (CUE,  $\text{mol}/\text{m}^2/\text{s}$ ) =  $P_n/C_i$  (He and Xu 2000).

On the 62<sup>nd</sup> day after salt treatment, extraction for determination of SOD, POD, CAT activities were prepared from 0.3 g of tissues homogenized under ice-cold conditions in 3 ml of extraction buffer, containing 50 mmol/L phosphate buffer (pH 7.4) at 4°C. The homogenates were centrifuged at 12 000 g for 20 min and the supernatant fraction was used for the assays. The activities of SOD, POD, CAT were measured by the standard analytical methods of Groppa et al. (2003), and Trevor et al. (1994). Lipid peroxidation in the leaf tissue was measured as formation of malondialdehyde (MDA, a product of lipid peroxidation) determined by thiobarbituric acid reaction using the method of Selvakumar (2008).

Height growth and biomass of the seedlings were measured on the 63<sup>rd</sup> day after salt treatment. All plants at each treatment were taken for biomass measurements and were divided into roots, leaves and stems. Then dry weight (dried at 105°C for 20 min and then dried at 70°C for 24 h) were measured on each root, leaf and stem sample.

**Statistical data analysis.** Statistical analysis was performed using the software SPSS 13.0. The results were subjected to a one-way ANOVA using the Tukey test to check for significant differences between means ( $P < 0.05$ ). Error bars in figures represent standard deviation.

## RESULTS AND DISCUSSION

**Effect of salt stress on seedling growth.** Seedling growth is normally limited by increasing concentration of NaCl (Sreenivasulu et al. 2000). In our

study, with increasing salinity levels, the plant height, root length, root fresh mass, root dry mass, stem fresh mass, stem dry mass, leaf dry mass of *Periploca sepium* Bunge were enhanced under low salinity levels (50 mmol/L NaCl), which reduced strongly with increasing salinity levels (Figure 1). Maximum reduction of all growth parameters were observed at 200 mmol/L NaCl. The plant height, root length, root fresh mass, root dry mass, stem fresh mass, stem dry mass, leaf fresh mass, dry mass were reduced by 40%, 47%, 65%, 58%, 60%, 57%, 54% and 38%, respectively, compared to control. The present results were in line with those of Parveen and Farrukh (2009), who reported decline in fresh and dry weights of shoots under high salinity stress. The stem dry mass, leaf fresh mass, dry mass were non-significant at 100 mmol/L NaCl compared to control, whereas they were significant at 50 mmol/L NaCl and at 200 mmol/L NaCl.

**Effect of salt stress on seedlings photosynthetic gas exchange.** Salinity reduces  $P_n$  and  $G_s$  in most plant species (Ashraf and Shahbaz 2003, Koyro

2006). In the present study, the  $P_n$  and  $G_s$  were enhanced under low salinity levels (50 mmol/L NaCl), and were reduced by 15%, 27%, at 150 mmol/L NaCl, and 50%, 47%, at 200 mmol/L NaCl, respectively, compared to control with increasing salinity levels (Figure 2). The changes are similar to the results of Yang et al. (2008b). Effect of salinity was non-significant on  $T_r$  at 50 mmol/L NaCl, but the  $T_r$  declined by 41%, 52%, at 150 mmol/L NaCl and 200 mmol/L NaCl, respectively, compared to control. With increasing salinity levels, the  $C_i$  reduced by 9%, 22% and 5% compared to control, but the  $L_s$  was enhanced by 10%, 28% and 4%. According to Farquhar and Sharkey's views (1984), at 100 mmol/L NaCl and 200 mmol/L NaCl stress, the decline of net photosynthetic rate of *Periploca sepium* Bunge was mainly caused by non-stomatal factors. The WUE, LUE, CUE were enhanced under low salinity levels (50 mmol/L NaCl) (Figure 2). With increasing salinity levels, the maximum value of WUE was observed at 100 mmol/L NaCl, the minimum value of WUE was observed at




Figure 1. Response of growth of *Periploca sepium* Bunge to salt stress. The different small letters were significantly different at 0.05 according to the Duncan test, the same as followed




Figure 2. Effect of salt stress on photosynthetic characteristics of *Periploca sepium* Bunge seedlings




Figure 3. Response of systems of anti-oxidation of *Periploca sepium* Bunge seedlings to salt stress

200 mmol/L NaCl; the LUE reduced by 17% at 150 mmol/L NaCl, 52% at 200 mmol/L NaCl compared to control; the CUE was enhanced by 8% at 150 mmol/L NaCl, while it was reduced by 47% at 200 mmol/L NaCl.

**Effect of salt stress on seedlings protective enzyme activities.** The level of the antioxidant enzymes SOD, POD, and CAT may determine the sensitivity of plants to lipid peroxidation (Kanazawa et al. 2000). In our study, with increasing salinity levels, activities of the antioxidative enzymes of SOD, POD, and CAT were enhanced by salts treatment of *Periploca sepium* Bunge seedlings, but CAT activity decreased at 200 mmol/L NaCl salt stress (Figure 3), indicating that the ability of these antioxidant enzymes to eliminate ROS is limited. The activity of enzymes of POD and CAT is higher than that of SOD at 200 mmol/L NaCl, which suggests that POD and CAT provide a better defense mechanism against salt stress-induced oxidative damage in *Periploca sepium* Bunge seedlings. The results are similar to Yeonghoo et al. (2004).

**Effect of salt stress on seedlings malondialdehyde.** The increased accumulation of lipid peroxides is indicative of enhanced production of toxic oxygen (Markovska et al. 2009). In this study, MDA concentration was non-significant compared to the control under low salinity levels (50 mmol/L NaCl), which was enhanced with increasing salinity levels, the maximum value was observed at 200 mmol/L NaCl (Figure 3). The results indicated that salt stress (> 100 mmol/L NaCl) produced superoxide radicals, resulting in increased lipid peroxidative products and oxidative stress in *Periploca sepium* Bunge seedlings.

In conclusion, *Periploca sepium* Bunge possesses moderate salt tolerance capacity, which can help to adapt to 100 mmol/L NaCl level. These results suggest a possibility to improve saline soil by utilizing *Periploca sepium* Bunge in Yellow River Delta region, where salinity is often the common major abiotic stress for plants. This study was done in the laboratory; although it may provide some information, it may not infer the actual field condition (Hameed et al. 2006). Therefore, further study is required to obtain some meaningful conclusions.

## REFERENCES

- Ashraf M., Shahbaz M. (2003): Assessment of genotypic variation in salt tolerance of early CIMMYT hexaploid wheat germplasm using photosynthetic capacity and water relations as selection criteria. *Photosynthetica*, 41: 273–280.
- Berry J.A., Downton W.J.S. (1982): Environmental regulation of photosynthesis. In: Covindjee (ed.): *Photosynthesis*. Vol. II. Academic Press, New York, 263–343.
- Bornhold B.D., Yang Z.S., Keller G.H., Prior D.B., Wiseman Jr W.J., Wang Q., Wright L.D., Xu W.D., Zhuang Z.Y. (1986): Sedimentary framework of the modern Huanghe (Yellow River) delta. *Geo-Marine Letters*, 6: 77–83.
- Dogan M., Tipirdamaz R., Demir Y. (2010): Salt resistance of tomato species grown in sand culture. *Plant, Soil and Environment*, 56: 499–507.
- Fang H.L., Liu G.H., Kearney M.S. (2005): Georelational analysis of soil type, soil salt content, landform, and land use in the Yellow River Delta, China. *Environmental Management*, 35: 72–83.
- Farquhar G.D., Sharkey T.D. (1982): Stomatal conductance and photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, 33: 317–345.

- Groppa M.D., Benavides M.P., Tomaro M.L. (2003): Polyamine metabolism in sunflower and wheat leaf discs under cadmium or copper stress. *Plant Sciences*, *164*: 293–299.
- Hameed A., Ahmed M.Z.M., Khan A. (2006): Comparative effects of NaCl and sea salt on seed germination of coastal halophytes. *Pakistan Journal of Botany*, *38*: 1605–1612.
- He W.M., Xu F.Y. (2000): Effects of water gradient on fluorescence characteristics and gas exchange in *Sabina vulgaris* seedlings. *Acta Phytoecologica Sinica*, *24*: 630–634.
- Kanazawa S., Sano S., Koshiha T. (2000): Changes in antioxidative in cucumber cotyledons during natural senescence: comparison with those during dark-induced senescence. *Physiologia Plantarum*, *109*: 211–216.
- Kao W.Y., Tsai T.T., Shih C.N. (2003): Photosynthetic gas exchange and chlorophyll a fluorescence of three wild soybean species in response to NaCl treatments. *Photosynthetica*, *41*: 415–419.
- Koyro H.W. (2006): Effect of salinity on growth, photosynthesis, water relations and solute composition of the potential cash crop halophyte *Plantago coronopus* (L.). *Environmental and Experimental Botany*, *56*: 136–146.
- Long S.P., Baker N.R., Rains C.A. (1993): Analyzing the responses of photosynthetic CO<sub>2</sub> assimilation to long-term elevation of atmospheric CO<sub>2</sub> concentration. *Vegetatio*, *104*: 33–45.
- Ma Y.M., Shi Q.H., Kong Y. (2008): Chemical constituents of twig bark from *Periploca sepium*. *Natural Product Research and Development*, *20*: 280–282.
- Maeda Y., Nakazawa R. (2008): Effects of the timing of calcium application on the alleviation of salt stress in the maize, tall fescue, and reed canarygrass seedlings. *Biologia Plantarum*, *52*: 153–156.
- Markovska Y.K., Gorinova N.I., Nedkovska M.P. (2009): Cadmium-induced oxidative damage and antioxidant responses in *Brassica juncea* plants. *Biologia Plantarum*, *53*: 151–154.
- Nijs I., Ferris R., Blum H. (1997): Stomatal regulation in a changing climate: A field study using free air temperature increase (FATL) and free air CO<sub>2</sub> enrichment (FACE). *Plant, Cell and Environment*, *20*: 1041–1050.
- Parveen A., Farrukh H. (2009): Growth performance of *Vicia satival* under saline conditions. *Pakistan Journal of Botany*, *41*: 3075–3080.
- Patel A.D., Pandey A.N. (2008): Growth, water status and nutrient accumulation of seedlings of *Holoptelea integrifolia* (Roxb.) Planch in response to soil salinity. *Plant, Soil and Environment*, *54*: 367–373.
- Qin J., Dong W.Y., He K.N., Yu Y., Tan G.D., Han L., Dong M., Zhang Y.Y., Zhang D., Li Z.A., Wang Z.L. (2010): NaCl salinity-induced changes in water status, ion contents and photosynthetic properties of *Shepherdia argentea* (Pursh) Nutt. seedlings. *Plant, Soil and Environment*, *56*: 325–332.
- Selvakumar V. (2008): Ultraviolet-B radiation (280–315 nm) invoked antioxidant defence systems in *Vigna unguiculata* (L.) Walp. and *Crotalaria juncea* L. *Photosynthetica*, *46*: 98–106.
- Sreenivasulu N., Grimm B., Wobus U. (2000): Differential response of antioxidant compounds to salinity stress in salt-tolerant and salt-sensitive seedlings of foxtail millet (*Setaria italica*). *Physiologia Plantarum*, *109*: 435–443.
- Trevor E., Kraus R., Austin F. (1994): Paclobutrazol protects wheat seedlings from heat and paraquat injury is detoxification of active oxygen involved. *Plant, Cell and Physiology*, *35*: 45–52.
- Wu Q.S., Zou Y.N., Liu W., Ye X.F., Zai H.F., Zhao L.J. (2010): Alleviation of salt stress in citrus seedlings inoculated with mycorrhiza: changes in leaf antioxidant defense systems. *Plant, Soil and Environment*, *56*: 470–475.
- Yang C.W., Shi D.C., Wang D.L. (2008a): Comparative effects of salt and alkali stresses on growth, osmotic adjustment and ionic balance of analkali-resistant halophyte *Suaeda glauca* (Bge.). *Plant Growth Regulation*, *56*: 179–190.
- Yang C.W., Wang P., Li C.Y., Shi D.C., Wang D.L. (2008b): Comparison of effects of salt and alkali stresses on the growth and photosynthesis of wheat. *Photosynthetica*, *46*: 107–114.
- Yeonghoo Kim, Joji Arihara, Takuji Nakayama, Norikazu Nakayama, Shinji Shimadaand, Kenji Usui (2004): Antioxidative responses and their relation to salt tolerance in *Echinochloa oryzicola* Vasing and *Setaria viridis* (L.) Beauv. *Plant Growth Regulation*, *44*: 87–92.
- Yildirim E., Karlidag H., Turan M. (2009): Mitigation of salt stress in strawberry by foliar K, Ca and Mg nutrient supply. *Plant, Soil and Environment*, *55*: 213–221.

Received on August 17, 2010

---

*Corresponding author:*

Prof. Zhaohua Lu, Binzhou University, Shandong Provincial Key Laboratory of Eco-Environmental Science for Yellow River Delta, Binzhou 256603, P.R. China  
 phone: + 865 433 195 580, fax: + 865 433 195 580, e-mail: lu-zhh@263.net

---