

## Evaluation of Fusarioses in Selected Grass Species

B. VOŽENÍLKOVÁ\* and F. KLIMEŠ

Faculty of Agriculture, University of South Bohemia, 370 05 České Budějovice,  
Czech Republic

\*E-mail: vozenil@zf.jcu.cz

### Abstract

In the years 1999–2001 we observed the spread and harmful effects of pathogenic fungi causing the dying of some grass species in permanent grass growths. The most common and frequent cause of the dying (winter freezing injury) is the fungus *Microdochium nivale* known also as *Fusarium nivale*, or *Gerlachia nivalis* (imperfect stage) – snow mould. The evaluation was carried out in dominant species (*Deschampsia cespitosa*, *Festuca rubra*, *Holcus mollis*) in permanent grass growths in the central part of the Bohemian Forest (locality Zhůří (1050–1080 m a.s.l.). The experiment involved three treatments: (1) Mulched growth (mulched only once), (2) Unharvested growth (fallow land) and (3) Mown growth (mown once in a year). The phytopathological evaluation took place during the vegetation periods in of 1999–2001 at three particular times (May, June, July). Our investigation has shown the most considerable development of the fungi of the genus *Fusarium* in the unharvested growth with serious decline of *Festuca rubra*, which deteriorates the conditions for subsequent fodder crops. On the other hand, the mown growth showed the least extent of fungi-attacked dominant grass species. The least occurrence of harmful *Fusarium* sp. div. disease (fusariosis) was found in *Deschampsia cespitosa* whereas the most serious damage occurred in *Festuca rubra*. From the viewpoint of temporal dynamics of fungal disease, the increase took place at the beginning of the second (1999) and first (2000, 2001) decade of May. In the tested dominant grasses we determined the genus *Fusarium* as the cause of fungal disease. Apart from further undetermined species, *Microdochium nivale* and *Fusarium solani* were the most frequent species.

**Keywords:** *Deschampsia cespitosa*; *Festuca rubra*; *Holcus mollis*; *Fusarium* sp. div.

### INTRODUCTION

The last years have witnessed considerable decline of interest in mountain meadows management. From the often inappropriate intensive management of some areas at the time of the former agrarian boom we have moved in the opposite direction – to abandonment of many originally cultivated areas (KLIMEŠ & KVĚT 1997b).

The remaining anthropogenic non-forested areas in the Bohemian Forest depend on human management. On selected managed sites this means preservation of favorable conditions for sustainable reproduction of the genetic pool of mountain grassland biota (KLIMEŠ & KVĚT 1997a).

We concentrated on the spread and harmfulness of pathogenic fungi causing damage to and death of some grass species (*Deschampsia cespitosa* (L.) P. Beauv., *Festuca rubra* L., *Holcus mollis* L.). The most common harmful agent is the fungus *Microdochium nivale* (Fr.: Fr.) Samuels & I. C. Hallet, of the order *Hypocreales*, class *Ascomycetes*, known better as *Fusarium nivale* (Fr.) Sorauer or *Gerlachia nivalis* (Ces. Ex Berl. & Voglino) W. Gams et E. Müller (imperfect stage) – snow mould (CAGAŠ 1998). According to SCHUMANN and BACKHAUS (1988), this species is dominant in the development of fusarioses occurring on the shoots of *Lolium multiflorum* Lam. var. *multiflorum* from the start of the growing season to the second grass cut. SNIJDERS and WINKELHORST (1996)

---

Supported by the Grant Agency of the Czech Republic (Grant No. 206/99/1410), by the Ministry of Education, Youth and Sports of the Czech Republic (Grants No. MSM J06/98:122200002, J06/98:122200003) and by the Ministry of Agriculture of the Czech Republic (Grant No. QL 1074).

investigated a grass turf in western Europe and proved that it was not the snow mould (*Microdochium nivale*) but other species of *Fusarium* [*F. cerealis* (Cooke) Sacc., *F. graminearum* Schwabe, *F. culmorum* (Wm. G. Sm.) Sacc. and *F. acuminatum* (Ellis & Everh.)] that were causing serious damage to grassland in the areas where *Lolium perenne* L. and *Festuca rubra* L. were dominant.

## MATERIAL AND METHODS

The spread and harmfulness of pathogenic fungi suppressing dominant species of grasses (*Deschampsia cespitosa*, *Festuca rubra*, *Holcus mollis*) was observed mainly from the phytopathological viewpoint. The experimental investigation itself was carried out in 1999–2001. The experimental site (Zhůří in the Šumava National Park) is situated at 1150–1180 m a. s. l. In our experiments, we tested three treatments (K – mowing, M – mulching, L – unharvested stand). Mowing was accomplished on July 10, 1999, July 15, 2000, and on July 15, 2001. Mulching was on July 11, 1999, July 14, 2000, and on July 16, 2001. Observations of each treatment were carried out in three replicate plots (3 × 30 m<sup>2</sup>).

Phytopathological analysis of plants with symptoms indicating an attack by *Fusarium* fungi was carried out in all treatments and replicates in May, June and July i.e. before mowing or mulching. In this experiment, we applied the method of symptoms evaluation in plants according to DIXON and DOODSON (1971).

We also evaluated the impact of the attack of *Fusarium* fungi on dominant grass species considering their phytocenological dynamics.

The isolation, cultivation and determination of the fungi of the genus *Fusarium* was performed regularly at each sampling time, on agar nutrient media (2% potato-glucose agar and SNA-Synthetischer Nährstoffärmer Agar), according to GERLACH and NIRENBERG (1982) and FASSATIOVÁ (1979).

The basic meteorological data during the observation period were summarized by KVĚT (1999, 2000, 2001).

## RESULTS AND DISCUSSION

On the experimental site at Zhůří we observed grasses dominant in permanent grass growths attacked by fungi of the genus *Fusarium*. Of the dominant grasses, we concentrated mainly on *Deschampsia cespitosa*, *Festuca rubra* and *Holcus mollis*.

Quality assessment was accomplished in each treatment and replicate at 10 points where we always evaluated ten plants according to the following scale (DIXON & DOODSON 1971):

- I – healthy plant, no symptoms of attack (0%)
- II – moderate symptoms of attack (1–20%)
- III – medium symptoms of attack (20–50%)
- IV – strong symptoms of attack (50–75%)
- V – withering or dead plant (75–100%).

During the entire vegetation period, we observed the development of fusarioses in the selected grasses.




Figure 1. The effect of different treatment on reduction of the *Fusarium* fungi




Figure 2. The development of the *Fusarium* fungi on selected grass species depending on the evaluation date




Figure 3. The extent of attack of selected grass species by the *Fusarium* fungi

The strongest impact of fungus infection was recorded at the beginning of the second decade of May in 1999 and of the first May decade in 2000, 2001. Statistical evaluation was made by means of non-parametric Kruskal-Wallis test, and the influence of both sampling date and treatment was highly significant in *Festuca rubra*, *Deschampsia cespitosa* and *Holcus mollis*. For the assessment of differences between particular categories we used Tukey HSD test. As to the date of evaluation, a statistically highly significant difference was found between the first (May), second (June) and third (July) assessments of quality in all observed grasses (Figure 2). In the experimental area, where

we observed possible influences of various types of management on the occurrence of fusarioses in grass growths, statistically highly significant differences were recorded between the following treatments: mulching and mowing, mulching and fallow land, and mowing and fallow land. Total differences between individual treatments applied to the grass growth were highly statistically significant (Figure 1).

During the observation of the attack of *Fusarium* fungi on selected grass species, considerable statistical differences existed between the extent of attack observed in *Deschampsia cespitosa* and *Festuca rubra*, and also in *Festuca rubra* and *Holcus mollis*. The least

extent of fusarioses was recorded in *Deschampsia cespitosa* whereas the most serious damage was found in *Festuca rubra*, especially in unharvested growths (Figure 3).

In these growths *Festuca rubra* also exhibited the most serious decline of occurrence: from the original cover degree of 30–40% to only 3–6%. In mulched growths of *Festuca rubra*, its moderate retreat was also recorded, approximately to 20–30%, owing to the attack of infection. The increased attack by fusarioses found in mulched and mainly unharvested *Festuca rubra* stands creates worse conditions for subsequent fodder production. The retreat of *Festuca rubra* in the observed stands was serious mainly with respect to the fact that this species represents the only cultural grass in the cenoses of interest.

Also SNIJDERS and WINKELHORST (1996) found serious damage caused by *Fusarium* fungi to grass stands dominated by *Festuca rubra*.

The entire evaluation of differences between particular years (1999 and 2001) was statistically non-significant.

In the dominants from the mulched, unharvested and mown growths tested *in vitro*, we determined the agents causing fungal diseases – fungi of the genus *Fusarium*. Apart from other unidentified species, *Microdochium nivale* and *Fusarium solani* (Mart.) Sacc., identified by J. HÝSEK and M. ZEMÁNKOVÁ, at the Research Institute of Crop Production, Division of Plant Medicine in Prague-Ruzyně belonged to the most numerous species.

## References

- CAGÁŠ B. (1998): Choroby a škůdci píceňích a travníkových trav. OSEVA PRO, s.r.o., Praha: 59.
- DIXON G.R., DOODSON J.K. (1971): Assessment keys for some diseases of vegetable, fodder and herbage crops. J. Nat. Inst. Agric. Bot., **12**: 299–307.
- FASSATIOVÁ O. (1979): Plísňě a vláknité houby v technické mikrobiologii. SNTL, Praha: 185–196.
- GERLACH W., NIRENBERG H. (1982): The Genus *Fusarium* – a Pictorial Atlas. Mitteilungen aus der biologischen Bundesanstalt für Land- und Forstwirtschaft, Berlin-Dahlem, 209, Verl. Paul, Berlin.
- KLIMEŠ F. (1997): Lukařství a pastvinářství. Ekologie travních porostů. Jihočeská Univ., České Budějovice.
- KLIMEŠ F., KVĚT J. (1997a): Travní porosty Šumavy na přelomu tisíciletí. Léto. Šumava, Vimperk, 2: 27–28.
- KLIMEŠ F., KVĚT J. (1997b): Travní porosty Šumavy na přelomu tisíciletí. Podzim. Šumava, Vimperk, 3: 24–25.
- KVĚT J. (1999): Funkce horských luk při různých způsobech jejich obhospodařování. [Zpráva.] GA ČR, České Budějovice.
- KVĚT J. (2000): Funkce horských luk při různých způsobech obhospodařování. [Zpráva.] GA ČR, České Budějovice.
- KVĚT J. (2001): Funkce horských luk při různých způsobech obhospodařování. [Zpráva.] GA ČR, České Budějovice.
- SCHUMANN K., BACKHAUS R. (1988): Untersuchungen zur Variabilität des Schneeschimmelerregers bei Futtergräsern. Nachr.-Blatt Pfl.-Schutz DDR, **42** (4): 75–78.
- SNIJDERS C.H.A., WINKELHORST G.D. (1996): An artificial inoculation method to screen for resistance to *Fusarium* rot in grasses. In: 2<sup>nd</sup> Int. Conf. Harmful and Beneficial Microorganisms in Grassland, Pastures and Turf., **19** (7): 265–271.