

Effect of long-term differential application of inorganic fertilizers and manure on soil CO₂ emissions

H.S. Dhadli, B.S. Brar

Department of Soil Science, Punjab Agricultural University, Ludhiana, India

ABSTRACT

Carbon dioxide (CO₂) fluxes from agricultural soils have been considered as one of the important environmental impact issue, due to their role in global warming and also its mitigation by carbon (C) sequestration in soils. Substantial scope of C sequestration with the application of inorganic fertilizers and manures has been reported, but the long-term effects of continuous application need to be critically examined. To study the effect of continuous differential application of NPK fertilizers and farmyard manure (FYM) in maize-wheat cropping system, CO₂ fluxes were measured via closed chambers and gas chromatography in a long-term experiment in progress for the past 42 years. The average daily CO₂ fluxes differed significantly amongst various treatments and were 55, 26 and 92% higher in NPK, N and NPK + FYM treatments over the control in the maize crop season and 43, 8 and 83% in the wheat crop season. Highly significant correlation of CO₂ emissions was found with soil organic carbon and total nitrogen in the maize and the wheat crop seasons. Although, CO₂ emissions were higher from long-term inorganic fertilizers and FYM treatments, still they are environmentally sustainable management practices, as they increased soil fertility and crop yields which consequently resulted in higher atmospheric CO₂ capture by plants and carbon sequestration in soils.

Keywords: fertilization; soil C-forms; soil N-forms; *Zea mays*; *Triticum* spp.

Terrestrial ecosystems are important sink of carbon (C) and could partially offset the increase in atmospheric CO₂ (Smith 1999). Conflicting results were reported about the changes in soil organic carbon (OC) due to use of inorganic fertilizers under intensive agriculture. Application of fertilizers, along with tillage and crop residue management practices, influence emissions of greenhouse gases (GHG) from agricultural soils (Adviento-Borbe et al. 2010, Mbonimpa et al. 2015, Dhadli et al. 2016). Further, cropping systems can influence CO₂ emission by affecting the quality and quantity of C returned to the soil as root biomass or as straw incorporation or mulching (Mapanda et al. 2011, Gong et al. 2012, Ding et al. 2014). Additionally, due to variations in management practices in different cropping systems, soil temperature and water content which directly control CO₂ emission are also influenced (Peng et al. 2011, Allaire et al. 2012, Zhang et al. 2013, Dhadli et al. 2015).

Although CO₂, CH₄ and N₂O emissions from agricultural fields under rice-wheat system in India have been reported from some field studies conducted over short term, information about CO₂ emissions from soils under long-term maize-wheat cropping system is lacking. The objectives of the present study were to study the impact of long-term application of inorganic fertilizers and farmyard manure (FYM) in a maize-wheat cropping system on CO₂ emissions and relationship of CO₂ emissions with various soil C and N (nitrogen) forms.

MATERIAL AND METHODS

An experiment under maize-wheat cropping system in progress since 1971 at the Research Farm, Punjab Agricultural University, Ludhiana, India, was selected to study the effect of long-term differential application of inorganic fertilizers and

doi: 10.17221/266/2015-PSE

FYM on CO₂ emissions in the maize and the wheat crop seasons. The experimental site is located at 75°47'09"E, 30°54'19"N at an altitude of about 247 m a.s.l. and the climate is subtropical, semiarid, hot and monsoon type with cold winters and hot summers. The soils of the area are developed above the flood plains sediments and are classified as Fluvisol (FL-dr) according to World Reference Base for Soil Resources (WRB) and have loamy sand texture. At the beginning of the experiment in 1971, the initial status of soil organic carbon (SOC), available P and K was 2.2, 6.1 and 58.7 g/kg, respectively. In the maize and the wheat crop seasons, CO₂ fluxes from five treatments: 100% NPK (T₁); 150% NPK (T₂); 100% N (T₃); 100% NPK + FYM (T₄), and control (T₅) were measured to study the effect of long-term application of inorganic fertilizers and FYM on CO₂ emissions. In different treatments, 100% NPK refers to the application of inorganic fertilizers viz., nitrogen (N), phosphorus (P) and potassium (K) at recommended level of 120 kg N + 26.2 kg P + 25.0 kg K/ha to each of maize and wheat crops. In T₃ treatment 100% N refers to the application of fertilizer N only at 120 kg/ha and in the control treatment (T₅) no input has been added in the past 42 years. Urea, single super phosphate (SSP) and muriate of potash (MOP) were used to supply N, P and K to maize and wheat crops. In T₄, FYM was applied at 10 t/ha (dry weight basis) before sowing of maize, which supplied 3700 kg C, 90 N, 140 kg P and 150 K/ha annually. Fertilizer N was applied in 3 equal doses in maize and 2 equal doses in wheat, while the whole of P and K were applied at sowing. Maize was sown at the end of May 2013 and at sowing 1/3 of N was applied, while remaining N was applied in two equal parts at 21 and 56 days after sowing (DAS). Wheat crop was sown in November 2013 and in all treatments except T₅, half of fertilizer N was applied at sowing and remaining after 30 days. Soil temperature in maize and wheat seasons ranged between 31–39°C and 11–24°C, respectively. Total rainfall received in maize crop season was 662 mm, while total rainfall in wheat crop season was 143 mm. Grain and straw yields were recorded at harvest and straw of both the crops was removed from field.

CO₂ fluxes were measured via closed chambers (10 cm internal diameter and 15 cm height) and gas chromatography. Air samples were collected in triplicate (within a minute) using a 10-mL sy-

ringe fitted with a hypodermic needle at 0, 15 and 30 min after chamber closure. The samples were immediately stored in 5-mL pre-evacuated glass vials equipped with a butyl rubber septum. The collected samples were taken to laboratory and CO₂ concentration in the samples was determined on the same day via a gas chromatograph (model 2100 Shimadzu Inc., Kyoto, Japan) equipped with a packed column, a flame ionization detector (FID) and a methanizer having Ni catalyst, 320°C. Daily CO₂ fluxes (CO₂-C kg/ha/day) were calculated from the linear temporal increase in mixing ratio of CO₂ in chamber headspace as elaborated by Parkin et al. (2012). Seasonal total CO₂ emissions (CO₂-C kg/ha) for each of maize and wheat crop seasons were calculated by interpolating flux between two successive measurements.

Surface soil (0–15 cm) samples were collected from different treatments before sowing of maize and wheat crops for the determination of various soil C and N forms. Soil organic carbon was determined by oxidation with potassium dichromate as described in Walkley and Black (1934) method and mineral nitrogen (N_{min}) by rapid titration method of Subbiah and Asija (1965). Total carbon (C_{tot}) and nitrogen (N_{tot}) were determined on CHN analyzer (Elementar Analysensysteme GmbH, Hanau, Germany) equipped with TC detector.

Statistical analysis of the data was performed via SAS 9.3 software (SAS Institute Inc., Cary, USA). A generalized linear mixed model (Glimmix) was used to perform ANOVA to study the significance of treatments as factorial structure, sample collection dates as repeated measures. In addition, an ANOVA was also run out to determine the effects of long-term treatments on various soil C and N forms, using the generalized linear model (GLM) of SAS 9.3 software. Long-term application of fertilizers and manures affected various soil C and N forms, therefore to study the effect of long-term treatments, correlations analysis between various soil C and N forms and CO₂ emissions from different treatments via Proc CORR was conducted.

RESULTS AND DISCUSSION

Crop yields and CO₂ emissions. Long-term application of fertilizers and manures in different treatments has a significant effect on grain and straw yields of both the maize and wheat crops.

Maize grain yield was highest in T₄ but statistically at par with T₂, and was followed by T₁. Grain yield in T₃ was lower than in T₁ but higher than in T₅. Wheat grain yields were significantly higher in NPK (T₁ and T₂) and FYM (T₄) treatments over N alone (T₃) and were lowest in control (T₅) (Table 1). Maize straw yield was significantly high in T₂, followed by T₄ and minimum in T₅, while the highest wheat biomass yield was observed in T₄ followed by T₂, T₁, T₃ and T₅. This implies that higher application of NPK fertilizer in maize contributed only to biomass production without any gains in grain yields. Continuous application of inorganic fertilizers and manure increased soil fertility, which helped in achieving higher crop yields. Higher dry matter yields indicated more capture of atmospheric CO₂ by plants.

Average of daily CO₂ fluxes during the maize crop season was the highest for T₄, followed by T₁ and T₂ and minimum for T₅ (Table 1). Seasonal total flux for the maize crop season was significantly higher by 26% to 55% in three fertilizer treatments and by 92% in FYM treatment over the control. In the wheat crop season, average of the daily fluxes was the highest for T₄ and lowest for T₅. Seasonal total CO₂ emissions were significantly higher by 43% in T₁ and T₂, 8% in T₃ and 83% in T₄, over T₅. Higher application of fertilizers (150% NPK) in T₂ had no significant effect on CO₂ emissions as compared to recommended level in T₁ (100% NPK), but CO₂ emissions and crop yields were significantly lower in T₃ receiving N fertilizer only. Continued applications of FYM along with NPK fertilizers, resulted in significantly high CO₂ emissions, in both the maize and the wheat crop

seasons. An increase in CO₂ emissions with the input of inorganic fertilizers and organic inputs has been reported from many studies (Song and Zhang 2009, Mapanda et al. 2011, Zhang et al. 2013).

Temporal variations in CO₂ fluxes. The daily CO₂ fluxes in the maize crop season ranged from 4.7 to 14.1 kg CO₂-C/ha/day in various treatments (Figure 1). Different long-term treatments, gas sampling dates and their interactions had a significant effect on daily CO₂ fluxes. CO₂ fluxes were comparatively higher at the beginning of maize crop season and thereafter declined sharply till 14 DAS (Figure 1). Significant surge in CO₂ fluxes after the application of N fertilizer on 21 and 56 DAS was registered in all the treatments, except T₅, and depression in fluxes was observed after rainfall associated with drop in soil temperature.

Amongst different treatments, CO₂ fluxes from T₄ treatment remained at significantly higher level throughout the maize crop season, except on 21 DAS, and similarly fluxes in T₁ and T₂ remained significantly higher than T₃ and T₅ till 77 DAS with one exception on 21st day (Figure 1). CO₂ fluxes in T₅ increased slightly between 21 to 49 DAS and declined gradually thereafter. They were at par with T₃ on 42 DAS, but were still significantly lower than T₁, T₂, T₃ and T₄. At the end of season, CO₂ fluxes were still significantly high in T₄, and T₁ was at par with T₂ at lower level, while T₃ was at par with T₅ at the lowest level. It became evident that CO₂ emissions were significantly higher in FYM treatment than in all the other treatments. The effect of fertilizer application on CO₂ fluxes at higher rates (T₂) as compared to recommended

Table 1. Effect of long-term application of inorganic fertilizers and farmyard manure (FYM) on CO₂ emissions and crop yields (t/ha)

	Average of daily flux (CO ₂ -C kg/ha/day)		Seasonal total emissions (CO ₂ -C kg/ha)		Grain yield		Straw yield	
	maize	wheat	maize	wheat	maize	wheat	maize	wheat
T ₁	9.29 ^b	5.7 ^b	898 ^b	813 ^b	4.59 ^{ab}	5.08 ^a	5.07 ^c	8.50 ^a
T ₂	9.34 ^b	5.7 ^b	903 ^b	819 ^b	5.04 ^a	5.43 ^a	8.67 ^a	8.75 ^a
T ₃	7.58 ^c	4.3 ^c	735 ^c	614 ^c	3.33 ^{bc}	4.23 ^b	3.70 ^d	5.77 ^b
T ₄	11.6 ^a	7.3 ^a	1112 ^a	1040 ^a	5.91 ^a	5.75 ^a	7.17 ^b	9.79 ^a
T ₅	6.01 ^d	4.0 ^d	578 ^d	563 ^c	2.25 ^c	1.72 ^c	2.99 ^d	2.87 ^c

Values followed by the same letters in a column are not significantly different ($P < 0.05$). T₁ – 100% NPK; T₂ – 150% NPK; T₃ – 100% N; T₄ – 100% NPK + FYM; T₅ – control

doi: 10.17221/266/2015-PSE




Figure 1. Temporal variations in weather variables and CO₂ fluxes in different treatments in the maize crop seasons (solid arrows indicate N-fertilization and dotted arrows irrigation). T₁ – 100% NPK; T₂ – 150% NPK; T₃ – 100% N; T₄ – 100% NPK + FYM; T₅ – control

level (T₁) was not significant. Application of N alone had some effect on CO₂ fluxes just after the fertilizer application, but generally they declined gradually in season. Another interesting observation was that in control treatment, CO₂ fluxes increased during the middle of season, which coincided with vegetative and flowering stages (Sey et al. 2010, Dhadli et al. 2015). Soil CO₂ fluxes were found associated with soil temperature and soil water content (Adviento-Borbem et al. 2010) and N-fertilization (Zhang et al. 2013).

In the wheat crop season, CO₂ fluxes were also significantly affected by long-term treatments, gas sample collection dates and their interactions. Before the sowing of wheat crop, CO₂ fluxes in various treatments ranged from 4.2 to 7.1 kg CO₂-C/ha/day and were significantly higher in T₄ than other treatments (Figure 2). Just after the sowing and application of basal dose of fertilizer-N, CO₂

emissions increased in all the treatments. Generally throughout the wheat crop season, CO₂ fluxes were significantly high in T₄, followed by T₁ and T₂ at par with each other and then T₃, while in T₅ fluxes remained at constant level. CO₂ fluxes decreased in all the treatments 11 DAS and remained stable up to 27 DAS. After the application of 2nd dose of urea on 37 DAS, fluxes increased significantly in all treatments except T₅. Between 49 DAS and 59 DAS, all season low CO₂ flux was recorded. Thereafter CO₂ fluxes gradually increased in all treatments till 102 DAS, but amazingly in control (T₅) after registering an increase, fluxes attained a plateau between 83 DAS and 93 DAS, and declined thereafter. On 116 day depression in CO₂ fluxes was observed in all treatments except T₄ and thereafter CO₂ emissions also started rising in all the other treatments, but the increase was significant in T₁, T₂ and T₄ only. The depressions




Figure 2. Temporal variations in weather variables and CO₂ fluxes in different treatments in the wheat crop seasons (solid arrows indicate N-fertilization and dotted arrows irrigation). T₁ – 100% NPK; T₂ – 150% NPK; T₃ – 100% N; T₄ – 100% NPK + FYM; T₅ – control

in CO₂ emissions in middle of wheat season coincided with drop in temperature, while towards the end of season soil moisture and temperature together controlled quantum of CO₂ emissions. Rise of flux in T₅ in the middle of season and then drop afterwards could have been the effect of higher root activity resulting in higher root exudation during vegetative and flowering growth stages (Dhadli et al. 2015). It was reported that specific root-derived respiration is higher during early vegetative growth and decline thereafter during reproductive and senescence stages (Fu et al. 2002, Sey et al. 2010, Jans et al. 2010). Further, soil CO₂ emissions have found to be regulated by soil temperature and moisture (Peng et al. 2011, Allaire et al. 2012, Dhadli et al. 2015). Soil moisture coupled with soil temperature and other soil-atmospheric interactions had been found to greatly affect soil CO₂ emissions, and an increase for shorter period

after N application was also recorded (Dhadli et al. 2015, Mbonimpa et al. 2015).

Relationship of CO₂ emission with soil C and N status. In the soil samples collected before the sowing of both the maize and wheat crops, SOC and C_{tot} were significantly high in T₄ followed by T₁ and T₂ at par with each other, while T₃ and T₅ were at par at lower level (Table 2). Effect of various long-term treatments was significant on N_{tot} and N_{min} and they were found significantly higher in T₄ and at lowest levels in T₅. Increase in SOC due to continuous cropping with the input of inorganic fertilizers and manures has been reported by Ding et al. (2014). During maize, average and total CO₂ flux have higher significant ($P < 0.01$) correlation with SOC and N_{tot} as compared to C_{tot} and N_{min} (Table 3). In wheat season, all C and N forms have highly significant correlation with average and total CO₂ emissions, but it was

doi: 10.17221/266/2015-PSE

Table 2. Effect of long-term application of inorganic fertilizers and farmyard manure (FYM) on soil carbon (C) and nitrogen (N) forms before sowing of maize and wheat crops

	SOC	C _{tot}	N _{min}	N _{tot}
	(g/kg)			
Maize				
T ₁	4.2 ^b	4.6 ^b	0.08 ^{bc}	0.48 ^{bc}
T ₂	4.2 ^b	4.5 ^b	0.09 ^{ab}	0.52 ^b
T ₃	3.3 ^c	3.9 ^c	0.07 ^c	0.44 ^c
T ₄	5.3 ^a	6.1 ^a	0.10 ^a	0.60 ^a
T ₅	3.0 ^c	3.6 ^c	0.06 ^c	0.39 ^d
Wheat				
T ₁	4.0 ^b	4.7 ^b	0.08 ^{ab}	0.51 ^{bc}
T ₂	4.0 ^b	4.8 ^b	0.09 ^{ab}	0.52 ^b
T ₃	3.3 ^c	3.7 ^c	0.07 ^{bc}	0.46 ^c
T ₄	5.2 ^a	5.9 ^a	0.10 ^a	0.68 ^a
T ₅	3.0 ^c	3.6 ^c	0.06 ^c	0.39 ^c

Values followed by the same letters in a column are not significantly different ($P < 0.05$). SOC – soil organic carbon; C_{tot} – total carbon; N_{min} – mineral nitrogen; N_{tot} – total nitrogen; T₁ – 100% NPK; T₂ – 150% NPK; T₃ – 100% N; T₄ – 100% NPK + FYM; T₅ – control

higher with both the soil C forms. Wheat grain and straw yield was found significantly correlated with N_{min}. The highest increase in C_{tot} upon long-term application of FYM followed by balanced NPK fertilizers in a long-term experiment under maize-wheat was recorded in China (Gong et al. 2009). CO₂ emissions were found related to SOC in maize and wheat seasons in a long-term experiment in China (Ding et al. 2007).

As a result of long-term differential application of inorganic fertilizers and FYM in the maize-wheat cropping system, the status of various soil C and N forms in various long-term treatments differed significantly after 42 years, which consequently resulted in significant differences in soil CO₂ emissions and crop yields amongst various long-term treatments. Although CO₂ emissions were higher from NPK fertilizer and FYM treatments, these treatments present status of soil C was significantly higher than the soil C status at the beginning of the experiment and the current status of control treatment. It can be concluded that although long-term application of balanced

Table 3. Correlation of soil carbon (C) and nitrogen (N) forms with CO₂ emissions and crop yields

	SOC	C _{tot}	N _{min}	N _{tot}
Maize				
Average flux	0.99**	0.96*	0.96*	0.98**
Cumulative flux	0.98**	0.96*	0.96*	0.98**
Grain yield	0.96*	0.91*	0.98**	0.97*
Straw yield	0.77	0.66	0.91*	0.82
Wheat				
Average flux	0.99**	1.00**	0.98**	0.98**
Cumulative flux	0.99**	1.00**	0.98**	0.98**
Grain yield	0.84	0.81	0.90*	0.86
Straw yield	0.92*	0.91*	0.96**	0.92*

* $P \leq 0.05$; ** $P \leq 0.01$; SOC – soil organic carbon; C_{tot} – total carbon; N_{min} – mineral nitrogen; N_{tot} – total nitrogen

inorganic fertilizers and FYM resulted in higher CO₂ emissions, still they are environmentally sustainable management practices, as they helped in achieving higher yields along with carbon sequestration in plants and soils.

REFERENCES

- Adviento-Borbe M.A.A., Kaye J.P., Bruns M.A., McDaniel M.D., McCoy M., Harkcom S. (2010): Soil greenhouse gas and ammonia emissions in long-term maize-based cropping systems. *Soil Science Society of America Journal*, 74: 1623–1634.
- Allaire S.E., Lange S.F., Lafond J.A., Pelletier B., Cambouris A.N., Dutilleul P. (2012): Multiscale spatial variability of CO₂ emissions and correlations with physico-chemical soil properties. *Geoderma*, 170: 251–260.
- Dhadli H.S., Brar B.S., Black T.A. (2015): Influence of crop growth and weather variables on soil CO₂ emissions in a maize-wheat cropping system. *Agricultural Research Journal*, 52: 28–34.
- Dhadli H.S., Brar B.S., Kingra P.K. (2016): Temporal variations in N₂O emissions in maize and wheat crop seasons: Impact of N-fertilization, crop growth, and weather variables. *Journal of Crop Improvement*, 30: 17–31.
- Ding W.X., Meng L., Yin Y.F., Cai Z.C., Zheng X.H. (2007): CO₂ emission in an intensively cultivated loam as affected by long-term application of organic manure and nitrogen fertilizer. *Soil Biology and Biochemistry*, 39: 669–679.
- Ding X.L., Yuan Y., Liang Y., Li L.J., Han X.Z. (2014): Impact of long-term application of manure, crop residue, and mineral

- fertilizer on organic carbon pools and crop yields in a Mollisol. *Journal of Soils and Sediments*, 14: 854–859.
- Fu S.L., Cheng W.X., Susfalk R. (2002): Rhizosphere respiration varies with plant species and phenology: A greenhouse pot experiment. *Plant and Soil*, 239: 133–140.
- Gong W., Yan X., Wang J., Hu T., Gong Y. (2009): Long-term manuring and fertilization effects on soil organic carbon forms under a wheat-maize cropping system in North China Plain. *Plant and Soil*, 314: 67–76.
- Gong W., Yan X.Y., Wang J.Y. (2012): The effect of chemical fertilizer on soil organic carbon renewal and CO₂ emission – A pot experiment with maize. *Plant and Soil*, 353: 85–94.
- Jans W.W.P., Jacobs C.M.J., Kruijt B., Elbers J.A., Barendse S., Moors E.J. (2010): Carbon exchange of a maize (*Zea mays* L.) crop: Influence of phenology. *Agriculture, Ecosystems and Environment*, 139: 316–324.
- Mapanda F., Wuta M., Nyamangara J., Rees R.M. (2011): Effects of organic and mineral fertilizer nitrogen on greenhouse gas emissions and plant-captured carbon under maize cropping in Zimbabwe. *Plant and Soil*, 343: 67–81.
- Mbonimpa E.G., Hong C.O., Owens V.N., Lehman M.R., Osborne S.L., Schumacher T.E., Clay D.E., Kumar S. (2015): Nitrogen fertilizer and landscape position impacts on CO₂ and CH₄ fluxes from a landscape seeded to switchgrass. *Global Change Biology – Bioenergy*, 7: 836–849.
- Parkin T.B., Venterea R.T., Hargreaves S.K. (2012): Calculating the detection limits of chamber-based soil greenhouse gas flux measurements. *Journal of Environment Quality*, 41: 705–715.
- Peng Q., Dong Y.S., Qi Y.C., Xiao S.S., He Y.T., Ma T. (2011): Effects of nitrogen fertilization on soil respiration in temperate grassland in Inner Mongolia, China. *Environmental Earth Sciences*, 62: 1163–1171.
- Sey B.K., Manceur A.M., Whalen J.K., Gregorich E.G., Rochette P. (2010): Root-derived respiration and nitrous oxide production as affected by crop phenology and nitrogen fertilization. *Plant and Soil*, 326: 369–379.
- Smith K.A. (1999): After the Kyoto Protocol: Can soil scientists make a useful contribution? *Soil Use and Management*, 15: 71–75.
- Song C.C., Zhang J.B. (2009): Effects of soil moisture, temperature, and nitrogen fertilization on soil respiration and nitrous oxide emission during maize growth period in northeast China. *Acta Agriculturae Scandinavica, Section B – Soil and Plant Science*, 59: 97–106.
- Subbiah B.V., Asija (1965): A rapid procedure for the estimation of available nitrogen in soils. *Current Science India*, 25: 259–260.
- Walkley A., Black I.A. (1934): An examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37: 29–39.
- Zhang J.J., Li Y.F., Jiang P.K., Zhou G.M., Shen Z.M., Liu J., Wang Z.L. (2013): Effects of fertilization on soil CO₂ flux in *Castanea mollissima* stand. *Journal of Applied Ecology*, 24: 2431–2439.

Received on April 24, 2015

Accepted on April 1, 2016

Corresponding author:

Dr. Harmanjit Singh Dhadli, Punjab Agricultural University, Department of Soil Science, Ludhiana, Punjab, India
e-mail: harman.dhadli@pau.edu
