

Long-term effect of low potassium fertilization on its soil fractions

M. Madaras¹, M. Koubová², M. Smatanová³

¹Crop Research Institute, Prague, Czech Republic

²Czech Geological Survey, Prague, Czech Republic

³Central Institute for Supervising and Testing in Agriculture, Brno, Czech Republic

ABSTRACT

In the Czech Republic, negative potassium (K) budget in agricultural soils is caused by non-fertilization by K and by a decline of manure application. We investigated soil available, fixed (acid-extractable, K_{fix}) and structural K pools in the field trial with graduated K application rate, established in 1972 at 8 sites of different climate and soils. The content of K-bearing minerals was evaluated on semi-quantitative scale by XRD diffraction. K-feldspars were a dominant source of structural K. Total soil K consisted of 1.7–7.1% of fixed K, which was in a positive relation to mixed-layer phyllosilicates. Differences in available K in treatments with K budget lower than -30 kg K/ha/year were small compared to those of fixed K. In control treatments, calculated average depletion of available K was -18 kg K/ha/year and the average depletion of fixed K was -12 kg K/ha/year; however at sites of higher altitude fixed K depletion prevailed. Fixed K accounted for 6–31% of the K budget. In negative K budget, monitoring of K_{fix} is advisable to avoid fertility loss of soil with low K supplying capacity.

Keywords: plant nutrition; soil mineralogy; X-ray diffraction; non-exchangeable potassium; long-term experiments

Potassium (K) is required by crop plants in large quantities as its concentration in plant biomass is at the second place just after the nitrogen (Syers 1998). The actual average annual K application rate in the Czech Republic is as low as 5.4 kg/ha of K in mineral fertilizers and 18 kg/ha in manures (Anonymous 2013), whereas overall outputs are 71 kg/ha on average (Klír et al. 2008). Since 1989, a highly negative K budget became standard for current agricultural practice in the Czech Republic, especially on farms without animals. The lack of K fertilization leads to lower K concentrations in plant tissues (Hejcman et al. 2013), which may result in higher plant susceptibility to low temperature, drought and salinity stress (Cakmak 2005). In many world areas, K deficiency leads to progressive decline in soil fertility and K status is recognized as an important factor restricting crop yields (Römheld and Kirkby 2010).

Mineral soils contain from 0.4 to 30 g/kg K, which is distributed between K available to plants (K_{avail} ; i.e. solution K and exchangeable K); fixed K (K_{fix}) and structural K (K_{struct}) in K-bearing feldspars and layer silicates (Huang 2005). Analysis of K_{avail} is standard for assessment of nutritional status of soils; however, a release of K_{fix} to the available forms takes place when K_{avail} is decreased by plant removal (Moritsuka et al. 2004), microbial activity and leaching (Martin and Sparks 1985). The measurement of K_{fix} is therefore often suggested to improve prediction of real availability of K to plants (Rees et al. 2013, Edmeades et al. 2014).

Knowledge about effects of non-fertilization on soil K fractions or soil K bearing minerals was usually gathered from one-site long-term experiments (Blake et al. 1999, Scherer et al. 2003, Barré et al. 2008, Lukin 2012, Hejcman et al. 2013). On the other hand, many studies relating K reserves to

Supported by the Ministry of Agriculture of the Czech Republic, Projects No. QI91C118 and RO0414.

soils of different mineralogy (Andrist-Rangel et al. 2010, Sarkar et al. 2013) do not include information about long-term K budget. In our research, we use the data from the experiment realized at eight sites covering a variety of soils and climatic conditions of arable soils of the Czech Republic, which allowed us to study soil and environmental interferences together with the influences of agricultural management. The aim of our research was to investigate the extent of soil K changes in conditions of negative K budget and how it is influenced by soil mineralogy and site characteristics.

MATERIAL AND METHODS

Experimental design. The long-term experiment (LTE) was established within the years 1972 and 1983 by the Central Institute for Supervising and Testing in Agriculture (CISTA). For this study, we selected two sites in warm regions (sugar beet growing region – SBGR) and six in moderately-warm region (potato growing region – PGR; Table 1). Until 1990, 9-year crop rotation was applied; afterwards crops were grown in 8-year crop rotation: 1. oat – alfalfa (SBGR)/oat – clover (PGR); 2. alfalfa/clover; 3. winter wheat; 4. potato; 5. winter wheat; 6. spring barley; 7. sugar beet (SBGR)/potato (PGR); 8. spring barley.

At each site, treatments were replicated 6-times. The trial included four treatments $N_2P_2K_{0,1,2,3}$ of graduated K mineral application rates: 0, 51, 81 and 119 kg K/ha/year (as KCl) in SBGR and 0, 57,

90 and 131 kg K/ha/year in PGR (in text referred to as K0–K3). Farmyard manure (FYM) was applied every 4 years at a rate of 40 t/ha (as fresh matter). Farmyard manure increased mean annual K input by 35 kg K/ha. Average N application rate was 112 kg N/ha/year and those of P 42 kg P/ha/year. Crop yields and K contents in main and side products were recorded every year by CISTA. Information about simple nutrient budget calculated as input of fertilizers minus above-ground crop uptake was taken from the report of Klement et al. (2012).

Soil sampling and analyses. Soil samples of plough horizon (depth of 0–25 cm) were taken after harvest 2010, dried at room temperature and sieved through 2 mm mesh. Mixed samples consisting of 6 subsamples (plot replications) were used for further analyses. K_{avail} was extracted by 1 mol/L ammonium acetate (1:10 w/v). K_{fix} was calculated as K extracted by 1 mol/L HCl (1:10 w/v, incubation 20 h at 50°C) subtracted by K_{avail} . *Aqua regia* extractable pool ($K_{aq.r.}$) was calculated as K extracted by *aqua regia* subtracted by K_{avail} and K_{fix} . Total K was analyzed after HF-HClO₄ digestion. Structural K was calculated as a difference between total K and K in *aqua regia* extract. Each soil sample was extracted twice and each extract was analyzed twice by ICP EOS (Thermo Jarrel Ash, Trace Scan, Franklin, USA). Organic carbon (C_{ox}) was analyzed after wet combustion. For soil texture specification, sedimentation method was used (ISO 11277). Cation exchange capacity (CEC) was analyzed by BaCl₂ method (ISO 13536). The soil mineralogical composition was determined from soils of K0 treatment on the basis of

Table 1. Characteristics of experimental sites and soils

Location	Establishment of the trial	Altitude (m)	Region	Mean annual precipitation (mm)	Mean annual air temperature (°C)	Soil type	C_{ox} (%)	Texture (%)			CEC	Exchangeable Ca+Mg (mmol ₊ /100 g)
								< 0.001	< 0.01	< 0.05		
								(mm)			(mmol ₊ /100 g)	
Uherský Ostroh	1972	196	SB	551	9.2	Luvisol	1.2	19.4	28.7	50.8	13.7	13.7
Žatec	1972	247	SB	451	8.3	Chernozem	2.1	26.8	45.4	75.6	26.8	25.6
Chrastava	1977	345	P	798	7.1	Luvisol	1.4	11.3	29.4	90.5	13.8	12.2
Staňkov	1981	370	P	511	7.8	Luvisol	1.3	12.5	27.9	67.1	10.9	10.9
Jaroměřice	1975	425	P	535	7.5	Luvisol	1.3	19.1	39.4	79.3	15.3	12.2
Svitavy	1981	460	P	624	6.5	Luvisol	1.1	9.8	24.4	47.6	11.0	11.0
Lípa	1974	505	P	632	7.7	Cambisol	1.4	7.5	31.6	67.7	13.4	8.9
Vysoká	1983	595	P	655	7.4	Stagnosol	1.6	9.8	32.8	59.1	14.8	12.9

Region – growing region of sugar beet (SB) or potato (P); CEC – cation exchange capacity

Table 2. Contents of K fractions (mg/kg) in soils at the beginning of the trial and in 2010 (treatments K0 and K3)

Location	K _{avail} at the beginning of the trial	Soil K pools				Total K
		K _{avail}	K _{fix}	K _{aq.r.}	K _{struct}	
Uherský Ostroh	271	232–338	885–950	3700–4500	9600	15 000
Žatec	238	149–378	657–754	4500–4900	9900	15 600
Chrastava	191	88–186	316–347	2300–2700	15 400	18 400
Staňkov	126	88–274	733–770	3200–2900	14 100	18 000
Jaroměřice	195	173–304	1313–1435	5100–5400	13 800	20 700
Svitavy	230	95–176	339–400	2100–2300	12 400	15 400
Lípa	147	78–172	783–939	3700–3900	16 300	21 000
Vysoká	200	98–198	452–536	2700–2500	9300	13 400

K0 – 0 kg K/ha/year; K3 – 131 kg K/ha/year; K_{avail} – K available to plants; K_{fix} – fixed K; K_{aq.r.} – *Aqua regia* extractable pool; K_{struct} – structural K

XRD patterns, which were obtained using CuK radiation with the powder diffractometer X'Pert System Philips with a graphite monochromator (Almelo, the Netherlands). The relative presence of main soil minerals was evaluated on semi-quantitative 5-point scale in a range from the limit of detectability (+) to dominant mineral (+++++). Statistical evaluation of the results was performed using the Statistica 10 software (Statsoft, Tulsa, USA).

RESULTS AND DISCUSSION

Soil K pools and mineralogy. Total K content in investigated soils was in the range from 13 000

to 21 000 mg/kg (Table 2). Total soil K comprises of 1.1% K_{avail} (0.5–2.5%), 4.3% K_{fix} (1.7–7.1%), 23% K_{aq.r.} (12.8–34.2%) and 72% K_{struct} (58.5–85%). K pools and their proportions to total K were significantly related both to environmental conditions of sites and to soil properties, however these two groups of parameters were interrelated. There was a positive relationship between mean site temperature and the share of K_{avail} ($P = 0.033$), K_{fix} ($P = 0.056$) and K_{aq.r.} ($P = 0.049$) on the total K. Soils of higher clay content had higher average K_{avail} content ($P = 0.042$). The content of K_{fix} was in a positive relation to mixed-layer phyllosilicates (MLP) ($P = 0.015$), which is due to preferential dissolution of MLP by HCl (Andrist-

Table 3. Soil mineralogical composition (relative presence: +++++ – dominant mineral; + – present at the limit of detectability)

Location	Quartz	Feldspars		Mica group	K + C	K:C	Mixed-layer phyllosilicates	A + P	
		plagioclase	K-feldspar						
Uherský Ostroh	+++++	++/+++	++	+	+	K	++/+++	I/S	
Žatec	+++++	++	++	+ / ++	++	K	++	I/S	
Chrastava	+++++	++/+++	++/+++	++	++	C ≥ K	+ / ++	I/S	
Staňkov	+++++	+++	+++ / +++++	+ / ++	+ / ++	K > C	+ / ++	I/S + C/S	++
Jaroměřice	+++++	+++	+++	++ / +++++	++ / +++++	C ≥ K	++ / +++++	I/S	
Svitavy	+++++	+ / ++	++	> +	+	K ≥ C	+	I/S	
Lípa	+++++	+++	+++	++ / +++++	++ / +++++	C ≥ K	++ / +++++	C/S + I/S	
Vysoká	+++++	+++	+ / ++	++	++	K ≥ C	+ / ++	C/S	

C – chlorite; K – kaolinite; S – smectite; I – illite; A – amphibole; P – pyroxene

Table 4. Relation between soil K fractions and soil and site parameters for treatments with minimum (K0) and maximum (K3) K fertilization

		K_{fix}	$K_{\text{aq.r.}}$	K_{struct}	Clay content	K-feldspar	Mica-group minerals	Mixed-layer silicates	Altitude
K0	K_{avail}	ns	0.75*	ns	0.80*	ns	ns	ns	ns
	K_{fix}	–	0.73*	ns	ns	ns	ns	0.82*	ns
	K_{struct}	–	–	–	ns	0.79*	ns	ns	ns
K3	K_{avail}	ns	0.74*	ns	0.95***	ns	ns	ns	–0.75*
	K_{fix}	–	0.85**	ns	ns	ns	ns	0.83**	ns
	K_{struct}	–	–	–	ns	0.79*	ns	ns	ns

K_{avail} – K available to plants; K_{fix} – fixed K; $K_{\text{aq.r.}}$ – *Aqua regia* extractable pool; K_{struct} – structural K; ns – statistically not significant; * $P \leq 0.05$; ** $P \leq 0.01$; *** $P \leq 0.001$

Rangel et al. 2013). The share of $K_{\text{aq.r.}}$ was negatively correlated with precipitation ($P = 0.029$) and positively with the content of fine particles ($P = 0.007$), which confirmed the dependence of K fractions on soil texture (Fotyma 2007). As a result of these complex influences, soils of SBGR had higher K_{avail} and also higher proportion of K_{avail} to total K than soils of PGR.

XRD analyses showed that the dominant soil mineral is quartz. All soils contained detectable amounts of main K-bearing minerals (K-feldspar, mica and mixed-layer phyllosilicates), but in different proportions (Table 3). This variability is typical for soils derived from different parent material (Andrist-Rangel et al. 2010). Dominant source of K_{struct} in the studied soils is K-feldspar, as we found a significant positive relationship between K_{struct} and its content ($P = 0.02$), whereas there was no significant relation to mica-mineral content (Table 4). Light-texture soils tend to contain less minerals of the mica group, especially at PGR ($P = 0.042$) and they also contained less MLP ($P = 0.063$; exception of this trend was the soil of Lípa).

Soil K changes affected by unbalanced fertilization. Since the beginning of the trial, K_{avail} changed markedly depending on differentiated K input; an average decrease by 38% with respect to initial value was observed at K0 while K_{avail} increased on average by 31% at K3 (Table 2). At five sites, the highest K application rates resulted in an increase of K_{avail} . However for the other three sites, the highest K application rate was sufficient only for keeping the initial K_{avail} level (Chrastava and Vysoká) or K_{avail} decreased even at the highest rate (Svitavy). It is possible that K application rates before the LTE establishment

were even higher than K3, as was speculated in a similar LTE (Madaras et al. 2010). Eventually, the land use might have been different in past; unfortunately details of the management before LTE establishment are unknown. These three sites also had K_{fix} content lower than 600 mg K_{fix} /kg. Soils of K_{fix} under this threshold tend to have decreasing K_{avail} even at positive K budget (Vopěnka and Macháček 1985). We found that the content of K_{fix} is connected with MLP content, which is known not only to release K to plants but also to fix K from fertilizer (Barré et al. 2008). Lower K_{fix} therefore indicates lower capability of soil to retain K. This can contribute to fertilizer K leaching together with highly water-permeable soil at Svitavy and increased precipitation at Chrastava.

Unbalanced fertilization induced a significant differentiation of both K_{avail} and K_{fix} with respect to treatments (Figure 1). At near-zero K budget, contents of K_{fix} in neighbour treatments K2 and K3 were approximately equal, whereas K_{avail} changed in the course of K budget surplus or shortage. At this situation, it appears that dynamics of K exchange among fertilizer, soil and crops is realized mainly through K_{avail} . However, for budgets lower than -30 kg K/ha/year, there were only minor differences of K_{avail} between neighbouring treatments as it reached the constant level, which is often observed in control plots of LTEs (Blake et al. 1999, Madaras and Lipavský 2009). At Žatec, Uherský Ostroh and Jaroměřice, K_{avail} at K0 stabilized at higher level compared to the other sites, most probably due to their higher clay content. In the situation of a negative K budget we observed a depletion of K_{fix} at all sites regardless their K_{fix} content. The lowest K_{fix} content was detected for




Figure 1. Relation between K budget and contents of K available to plants (K_{avail}) and fixed K (K_{fix}) at 3 sites with K application from 0 to 166 kg K/ha/year (including unfertilized control treatments)

treatments of the lowest K input, which is in accordance with the results of other LTEs (Blake et al. 1999, Scherer et al. 2003). Obviously the demand of crops for K in conditions of long-term low K input was fulfilled mainly from non-exchangeable K pool either directly (Moritsuka et al. 2004) or indirectly by a transfer through exchangeable sites.

Based on regressions between K budgets and K contents, we calculated the expected extent of K pools for the situation of zero K budget and the approximate annual depletion of K fractions for treatment K0 with respect to this zero budget (Table 5). The average depletion of K_{avail} was -18 kg

K/ha/year and the average depletion of K_{fix} was -12 kg K/ha/year. At soils of higher altitude K_{fix} decrease exceeded the decrease of K_{avail} , which can be explained by a combined effect of lighter soils and a more intensive profile percolation. An efficiency of K fertilization by highly soluble K salts would be lower at these sites. However the correlation of the altitude vs. K_{avail} was not statistically significant ($P = 0.099$). K_{fix} decrease in a plough layer accounted for 6% to 31% of the K budget. A decrease of both forms accounted from 17% to 78%, giving evidence that substantial K uptake is realized from other soil sources, e.g. other forms and/or deeper layers (Andrist-Rangel et al. 2007). Lukin (2012) reported that crops at sandy-loamy soil in conditions of K deficit use from 31 to 41 kg/ha of non-exchangeable K.

Average change of K_{fix} per the unit of K budget (i.e. 1 kg K/ha/year) was 1.3 mg K_{fix} /kg soil. The variability of this parameter among sites appears to have relation especially to the soil texture and K_{fix} content; the multilinear regression including these two parameters explains 89% of data variation ($P < 0.001$). According to it, K_{fix} depletion per the unit of K budget is enhanced in heavier soils and in soils of low K_{fix} .

Finally we concluded that intensive crop production with unbalanced K fertilization leads to significant depletion of K_{fix} . The rate of this depletion can be larger than that of K_{avail} and it is dependent on K budget, soil and site characteristics. Even though the annual decrease of K_{fix} is relatively small compared to the extent of this K pool, it is advisable to monitor K_{fix} to avoid fertility loss in soil of low K supplying capacity.

Table 5. Calculated annual K depletion of available (ΔK_{avail}) and fixed (ΔK_{fix}) K pools for treatment K0 (assuming 25 cm plough layer and specific weight 1.4 g/cm³)

Location	K budget for treatment K0 (kg K/ha/year)	K calculated for zero budget		Difference between K0 and contents at zero K budget		
		K_{avail}	K_{fix}	ΔK_{avail}	ΔK_{fix}	sum
		(mg/kg)		(kg/ha/year)		
Uherský Ostroh	-117	340	944	-13	-7	-20
Žatec	-53	342	758	-24	-13	-37
Chrastava	-68	353	403	-40	-13	-53
Staňkov	-80	193	761	-19	-5	-24
Jaroměřice	-68	243	1391	-10	-11	-21
Svitavy	-90	154	407	-11	-12	-23
Lípa	-68	179	940	-14	-21	-35
Vysoká	-70	156	533	-12	-17	-29

REFERENCES

- Andrist-Rangel Y., Edwards A.C., Hillier S., Öborn I. (2007): Long-term K dynamics in organic and conventional mixed cropping systems as related to management and soil properties. *Agriculture, Ecosystems and Environment*, 122: 413–426.
- Andrist-Rangel Y., Hillier S., Öborn I., Lilly A., Towers W., Edwards A.C., Paterson E. (2010): Assessing potassium reserves in northern temperate grassland soils: A perspective based on quantitative mineralogical analysis and *aqua-regia* extractable potassium. *Geoderma*, 158: 303–314.
- Andrist-Rangel Y., Simonsson M., Öborn I., Hillier S. (2013): Acid-extractable potassium in agricultural soils: Source minerals assessed by differential and quantitative X-ray diffraction. *Journal of Plant Nutrition and Soil Science*, 176: 407–419.
- Anonymous (2013): Agriculture 2012. Ministry of Agriculture of the Czech Republic, Prague. (In Czech)
- Barré P., Montagnier C., Chenu C., Abbadie L., Velde B. (2008): Clay minerals as a soil potassium reservoir: Observation and quantification through X-ray diffraction. *Plant and Soil*, 302: 213–220.
- Blake L., Mercik S., Koerschens M., Goulding K.W.T., Stempen S., Weigel A., Poulton P.R., Powlson D.S. (1999): Potassium content in soil, uptake in plants and the potassium balance in three European long-term field experiments. *Plant and Soil*, 216: 1–14.
- Cakmak I. (2005): The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *Journal of Plant Nutrition and Soil Science*, 168: 521–530.
- Edmeades D.C., Morton J.D., Waller J.E., Metherell A.K., Roberts A.H.C., Carey P. (2014): The diagnosis and correction of potassium deficiency in New Zealand pastoral soils: A review. *New Zealand Journal of Agricultural Research*, 53: 151–173.
- Fotyra M. (2007): Content of potassium in different forms in the soils of southeast Poland. *Polish Journal of Soil Science*, 40: 19–32.
- Hejzman M., Berková M., Kunzová E. (2013): Effect of long-term fertilizer application on yield and concentrations of elements (N, P, K, Ca, Mg, As, Cd, Cu, Cr, Fe, Mn, Ni, Pb, Zn) in grain of spring barley. *Plant, Soil and Environment*, 59: 329–334.
- Huang P.M. (2005): Chemistry of potassium in soils. In: Tabatabai M.A., Sparks D.L. (eds): *Chemical Processes in Soils*. Soil Science Society of America, Madison.
- Klement V., Smatanová M., Trávník K. (2012): Forty Years of Long-Term Field Experiments in ÚKZÚZ. Central Institute for Supervising and Testing in Agriculture, Brno. (In Czech)
- Klír J., Kunzová E., Čermák P. (2008): *Frame Methodics of Plant Nutrition and Fertilization*. Crop Research Institute, Prague. (In Czech)
- Lukin S.M. (2012): Potassium status and budget in sandy loamy soddy-podzolic soil under long-term fertilization. *Agrokhimiya*, 12: 5–14. (In Russian)
- Madaras M., Koubová M., Lipavský J. (2010): Stabilization of available potassium across soil and climatic conditions of the Czech Republic. *Archives of Agronomy and Soil Science*, 56: 433–449.
- Madaras M., Lipavský J. (2009): Interannual dynamics of available potassium in a long-term fertilization experiment. *Plant, Soil and Environment*, 55: 334–343.
- Martin H.W., Sparks D.L. (1985): On the behavior of nonexchangeable potassium in soils. *Communications in Soil Science and Plant Analysis*, 16: 133–162.
- Moritsuka N., Yanai J., Kosaki T. (2004): Possible processes releasing nonexchangeable potassium from the rhizosphere of maize. *Plant and Soil*, 258: 261–268.
- Rees G.L., Pettygrove G.S., Southard R.J. (2013): Estimating plant-available potassium in potassium-fixing soils. *Communications in Soil Science and Plant Analysis*, 44: 741–748.
- Römheld V., Kirkby E.A. (2010): Research on potassium in agriculture: Needs and prospects. *Plant and Soil*, 335: 155–180.
- Sarkar G.K., Chattopadhyay A.P., Sanyal S.K. (2013): Release pattern of non-exchangeable potassium reserves in Alfisols, Inceptisols and Entisols of West Bengal, India. *Geoderma*, 207–208: 8–14.
- Scherer H.W., Goldbach H.E., Clemens J. (2003): Potassium dynamics in the soil and yield formation in a long-term field experiment. *Plant, Soil and Environment*, 49: 531–535.
- Syers J.K. (1998): *Soil and Plant Potassium in Agriculture*. International Fertilizer Society, Newcastle upon Tyne.
- Vopěnka L., Macháček V. (1985): Study of the balance of potassium reserve in the soil. *Rostlinná Výroba*, 31: 905–912. (In Czech)

Received on April 8, 2014

Accepted on June 12, 2014

Corresponding author:

RNDr. Mikuláš Madaras, Ph.D., Výzkumný ústav rostlinné výroby, v.v.i., Drnovská 507,
161 06 Praha-Ruzyně, Česká republika
e-mail: madaras@vurv.cz