

<https://doi.org/10.17221/296/2019-PSE>

Phosphorus affects enzymatic activity and chemical properties of cotton soil

YANG GAO¹, HUIYI HUANG¹, HONGYI ZHAO¹, HOUQIANG XIA¹, MIAO SUN², ZONGYUN LI¹, PENGCHENG LI², CANGSONG ZHENG², HELIN DONG^{2*}, JINGRAN LIU^{1*}

¹Jiangsu Key Laboratory of Phylogenomics and Comparative Genomics, School of Life Science, Jiangsu Normal University, Xuzhou, P.R. China

²State Key Laboratory of Cotton Biology, Institute of Cotton Research of Chinese Academy of Agricultural Sciences, Anyang, Henan, P.R. China

*Corresponding authors: liujingran_66@163.com; donghl668@sina.com

Citation: Gao Y., Huang H.Y., Zhao H.Y., Xia H.Q., Sun M., Li Z.Y., Li P.C., Zheng C.S., Dong H.L., Liu J.R. (2019): Phosphorus affects enzymatic activity and chemical properties in cotton soil. *Plant Soil Environ*, 65: 361–368.

Abstract: Pot experiments were conducted in 2017 with two cotton cultivars (CCRI 79 and LMY 28) and three phosphorus (P) levels: 3, 8 and 12 mg P₂O₅/kg as P₀, P₁ and P₂, respectively. In this study, the soil water-soluble organic carbon content increased as the soil available P (AP) increased, while there were no significant variations for soil total organic matter content among the three AP levels. The activities of invertase, cellulase and urease in cotton soil decreased significantly in the P₀. There were positive correlations between invertase and cellulase activities with soil organic carbon and inorganic-nitrogen (N); these correlated negatively with soil C/N ratio and AP level. In addition, high soil AP can raise soil AP and enhance alkaline phosphatase activity, which had a significant negative relationship with the soil C/P ratio. Urease activity had a significant positive relationship with soil NH₄⁺-N, C/P and N/P, as well as a negative correlation with soil C/N. Moreover, soil NH₄⁺-N and NO₃⁻-N in the P₁ and P₂ were lower than in the P₀, which might be an effect of high AP on soil N availability.

Keywords: soil available phosphorus; soil enzymes; C, N and P availability; plant nutrition; fertilization; soil microorganisms

Phosphorus (P) is an important nutrient in the soil; it limits plant growth and metabolism due to its low mobility and bioavailability. Low P availability is frequently a primary limiting factor for crop production in many soils (Shen et al. 2011, Mühlbachová et al. 2018). Indeed, it has been estimated that 5.7 billion hectares worldwide contain insufficient available P (AP) for optimal crop production (Cordell and White 2013, Niu et al. 2013), and about 30% of cultivated land in China has only 3–5 mg AP/kg soil (Wu 2013).

The management of soil quality is crucial to improve agricultural productivity and environmental quality. In the absence of fertilizer inputs, soil organic carbon (OC) is a primary source of plant nutrients. P application can increase soil C, N and P (Griffiths

et al. 2012), especially soil AP for plants (Djodjic and Mattsson 2013), and subsequently reduce the C/P ratio in the rhizosphere. In addition, increasing soil AP can affect the C/P ratio by decreasing the carbohydrate contents released per unit of plant root (Schilling et al. 1998, DeForest et al. 2012). However, little is known about soil C, N and P availability in low AP soil for cotton. Soil total organic carbon (TOC) can promote the accumulation of soil nutrients such as alkali-hydrolyzed nitrogen, AP and available potassium by soil enzymes (Bowles et al. 2014). Thus, the soil enzymes are generally used to evaluate the factors affecting fertilizer application, soil quality and plant litter decomposition (Bowles et al. 2014, Mi et al. 2018). These soil enzymes associated with

Yang Gao and Huiyi Huang contributed equally to this paper. Supported by the National Natural Science Foundations of China, Grant No. 31501257; by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), and by the China Agriculture Research System, Project No. CARS-18-17.

the C, N, and P cycles are invertase, β -glucosidase, and cellulase, urease, and phosphatase (Adetunji et al. 2017).

Invertase acts by the organic compounds degrading to cut down their molecular size and make smaller organic structures, thus promoting microbial enzymatic activities (Li et al. 2014). β -glucosidase can hydrolyze the cellobiose residue during the final stage of the cellulose degradation process (Adetunji et al. 2017). Cellulase catalyzes the hydrolysis of cellulose to glucose. Urease is a crucial enzyme regulating nitrogen (N) conversion in soil and is mainly derived from microorganisms and plants (Li et al. 2014). Alkaline phosphatases (ALP) act by hydrolyzing simple phosphomonoesters and acquiring orthophosphate (Fraser et al. 2017), and are only produced by soil microorganisms (Spohn and Kuzyakov 2013). It is of great significance to study the response characteristics of these soil enzymatic activities to different soil AP levels and their relationships with soil chemical properties in order to reveal the rules for transformation and utilization of soil C, N and P, and in addition to improve the utilization efficiency of P fertilizer in cotton soil.

At present, few studies have been published on the effects of soil AP levels on soil enzymatic activities and active organic matter in cotton soil (Chen et al. 2014). The present study used soil with three different soil AP levels to evaluate the effect of soil AP levels on soil enzymatic activities and chemical properties, to provide data support and a scientific basis for evaluation of cotton soil quality, establishment of a reasonable fertilization system and improvement of land productivity.

MATERIAL AND METHODS

Experimental design. A pot experiment was conducted in 2017 at the Baibi station, Anyang, Henan, China (36°06'N, 114°21'E). In a depth of 0–20 cm before sowing cotton, the TOC and total N, available N and available K of the soil were 16.2 g/kg, 1.0 g/kg, 71.2 mg/kg and 106.7 mg/kg, respectively. Cotton seeds of cvs. CCRI 79 and LMY 28 were planted into plastic pots on April 19th, 2017. The quantitative irrigation was applied as needed to minimize the moisture stress during each season.

The treatments for the pot experiments consisted of three AP (P) levels: (i) 3 ± 0.5 mg/kg (P_0 , P deficiency), (ii) 8 ± 0.5 mg/kg (P_1), and (iii) 12 ± 0.5 mg/kg using triple superphosphate (P_2 , moderate P under the AP in this experiment) (Zheng et al. 2015), as a control.

In each plot at the flower and boll stage and the boll-opening stage, three soil samples from 0–20 cm depth were chosen from three replicate plots randomly. The samples were then used for the measurement of soil enzymes and soil chemical properties.

Determination of soil chemical properties and soil enzymes. Soil TOC was measured using the dichromate oxidation method (Hai et al. 2010). Soil water-soluble organic carbon (WSOC) was measured using the dichromate oxidation method. Soil water-soluble total nitrogen (WSTN) was measured using the Kjeldahl acid-digestion method after extraction with sodium hydroxide-potassium persulfate. Soil ammonium nitrogen (NH_4^+ -N) and nitrate nitrogen (NO_3^- -N) were analysed with the automated discrete analyser (SmartChem 200, AMS Alliance, Rome, Italy) after soil samples were extracted with potassium chloride, and then shaken at 200 rpm for 1 h. Soil AP was analysed using the Olsen-P method (Yang and Jacobsen 1990). Kits for invertase, β -glucosidase, cellulase, urease and ALP were acquired from the Suzhou Coming Biotechnology, Suzhou, China.

Statistical analysis. Data were handled by SPSS V24.0 (IBM corp., New York, USA), and the differences between mean values with $P < 0.05$ were determined as significant. The ratios of WSOC to WSTN (C/N), SOC to AP (C/P), WSTN to AP (N/P) were also calculated.

RESULTS

Effect of phosphorus levels on soil enzymatic activity

Soil enzymes related to C cycle. During cotton growth, the activity of invertase in the soil increased (Figure 1a,d). With increasing soil AP level, soil invertase activity decreased significantly ($P < 0.05$). Compared to the P_0 , the soil invertase activity in the P_1 and P_2 for cv. CCRI 79 decreased by (–1.8)–15.0% and 25.3–37.6%, respectively, and by 20.5–25.4% and 35.1–69.0% for cv. LMY 28, respectively. The changing trend of soil cellulase activity was similar to that of invertase (Figure 1c,f).

The activity of β -glucosidase exhibited an increasing trend with cotton growth (Figure 1b,e). There were no significant differences in soil β -glucosidase activity by soil AP level. However, the soil β -glucosidase activity initially increased and then decreased as the soil AP level increased. Compared to the P_0 , soil β -glucosidase activity for cv. LMY 28 in the P_1 and P_2 increased by 8.1–27.1% and 1.0–13.6%, respectively.

<https://doi.org/10.17221/296/2019-PSE>




Figure 1. Effect of phosphorus levels on soil invertase activity, β -glucosidase activity and cellulase activity. Values followed by a different letter within the same figure mean significant difference at 0.05 probability level. Lowercase letters and uppercase letters indicate cvs. CCRI 79 and LMY 28, respectively. P₀ – 3, P₁ – 8, P₃ – 12 mg P₂O₅/kg; FBS – the flower and boll stage; BOS – the boll-opening stage

Soil enzyme related to N cycle-urease. Soil urease activity increased with cotton growth and decreased significantly as soil AP increased ($P < 0.05$, Figure 2). Analysis of variance showed that there was no significant difference in soil urease activity between the P₀ and P₁. During the flowering and boll stage, soil urease activity in the P₂ treatment decreased by

14.2% for cv. CCRI 79 and by 21.7% for cv. LMY 28, compared with the P₀. However, the soil urease activity of cv. CCRI 79 was not significantly affected by soil AP level during the boll-opening stage, and only the soil urease activity of cv. LMY 28 decreased by 27%.

Soil enzyme related to P cycle-ALP. The soil ALP activity increased significantly as the soil




Figure 2. Effect of phosphorus levels on soil urease activity. Values followed by a different letter within the same figure mean significant differences at 0.05 probability level. Lowercase letters and uppercase letters indicate cvs. CCRI 79 and LMY 28, respectively. P₀ – 3, P₁ – 8, P₃ – 12 mg P₂O₅/kg; FBS – the flower and boll stage; BOS – the boll-opening stage




Figure 3. Effect of phosphorus levels on soil alkaline phosphatase activity. Values followed by a different letter within the same figure mean significant differences at 0.05 probability level. Lowercase letters and uppercase letters indicate cvs. CCRI 79 and LMY 28, respectively. P₀ – 3, P₁ – 8, P₂ – 12 mg P₂O₅/kg; FBS – the flower and boll stage; BOS – the boll-opening stage

AP increased, and the range by which it changed during flowering and boll period was higher than that during boll-opening period ($P < 0.05$, Figure 3). With the increase of soil AP, the soil ALP activity for cv. CCRI 79 and cv. LMY 28 increased by 21.2–69.9% and 44.0–127.7%, respectively. In addition, the soil ALP activity for cv. CCRI 79 was significantly higher than that for cv. LMY 28.

Effect of P levels on soil chemical properties

Soil AP content. Soil AP level increased significantly with increasing soil AP level (Figure 4), and its content was basically within the range of soil AP level set by the current research.

Soil TOC and WSOC content. There were no significant differences among the three soil AP levels for soil TOC content ($P > 0.05$, Figure 5a). Nevertheless, the soil WSOC content showed an increasing trend with increasing soil AP levels (Figure 5b). Compared to the P₀, the soil WSOC content in the P₁ and P₂ increased by 5.9–10.0% and 15.0–17.0%, respectively. The C/P ratio in soil was > 500 and increasing AP levels gradually decreased the soil C/P ratio.

Soil NH₄⁺-N, NO₃⁻-N and WSTN content. The NH₄⁺-N content decreased with the increase of soil AP (Figure 6a). Analysis of variance showed that the P₀ and P₁ had no significant difference for NH₄⁺-N content in soil, and both reached significant levels compared with the P₂ ($P < 0.05$). Compared with the P₀, the NH₄⁺-N content for cv. CCRI 79 and cv. LMY 28 in the P₂ reduced by 22.8% and 15.1%, respectively. The NO₃⁻-N content showed a significant

downward trend with increasing soil AP (Figure 6b). Compared with the P₀, soil NO₃⁻-N content in the P₁ and P₂ decreased by 25.5–44.5% and 36.2–59.1%, respectively.

The WSTN content significantly decreased with increasing soil AP (Figure 6c). However, the soil WSTN for cv. CCRI 79 and cv. LMY 28 only showed a significant variation between the P₀ and P₂ ($P < 0.05$). Compared with the P₀, soil WSTN content in the P₂ was reduced by 21.2% for cv. CCRI 79, and by 13.7% for cv. LMY 28. However, soil C/N ratio showed the opposite trend as the soil AP increased (Table 1). In addition, the C/N ratio had a significant variation in




Figure 4. Effect of phosphorus levels on soil available phosphorus content. Values followed by a different letter within the same figure mean significant differences at 0.05 probability level. Lowercase letters and uppercase letters indicate cvs. CCRI 79 and LMY 28, respectively. P₀ – 3, P₁ – 8, P₂ – 12 mg P₂O₅/kg

<https://doi.org/10.17221/296/2019-PSE>




Figure 5. Effect of phosphorus levels on soil organic carbon (TOC) and water-soluble organic carbon (WSOC) content. Values followed by a different letter within the same figure mean significant differences at 0.05 probability level. Lowercase letters and uppercase letters indicate cvs. CCRI 79 and LMY 28, respectively. P₀ – 3, P₁ – 8, P₃ – 12 mg P₂O₅/kg

cotton soil only between the P₀ and P₂. Moreover, the soil C/P ratio was gradually reduced with increasing AP levels.

Correlation analysis between soil enzymatic activities and soil chemical properties. Soil invertase, cellulase and urease had significantly positive correlations with soil TOC and NH₄⁺-N, and nega-

tive correlations with soil WSOC and AP (Table 2). In addition, soil ALP had significantly positive correlations with soil WSOC and AP and negative correlations with NH₄⁺-N. Soil cellulase was significantly correlated with NO₃⁻-N. Moreover, these four enzymes also significantly correlated with each other, because they belong to the same group of hydrolases. The




Figure 6. Effect of phosphorus levels on soil NH₄⁺-N, NO₃⁻-N and water-soluble total nitrogen (WSTN) content. Values followed by a different letter within the same figure mean significant differences at 0.05 probability level. Lowercase letters and uppercase letters indicate cvs. CCRI 79 and LMY 28, respectively. P₀ – 3, P₁ – 8, P₃ – 12 mg P₂O₅/kg

Table 1. Effect of phosphorus levels on the ratio of C/P, C/N and N/P in cotton soil

Phosphorus level	C/P ratio		C/N ratio		N/P ratio	
	CCRI 79	LMY 28	CCRI 79	LMY 28	CCRI 79	LMY 28
P ₀	1767 ^a	1902 ^a	0.51 ^b	0.56 ^b	10.89 ^a	10.40 ^a
P ₁	926 ^{bc}	1072 ^b	0.58 ^b	0.69 ^a	5.79 ^b	5.83 ^b
P ₂	656 ^c	649 ^c	0.74 ^a	0.76 ^a	3.41 ^b	3.41 ^b

Values followed by the different letters within the same index are significantly different at $P < 0.05$ probability level. P₀ – 3, P₁ – 8, P₂ – 12 mg P₂O₅/kg

correlation between β -glucosidase and soil chemical properties did not reach significant levels (data not shown).

DISCUSSION

Soil C, N and P availability. WSOC is the intermediate product of soil OC transformation and microbial metabolism. These two competing processes contribute to soil management and/or plant nutrient availability, to microbial activity and a series of enzymes that determine plant productivity and/or soil fertility (Powlson et al. 2011). In this study, soil WSOC increased with increasing soil AP (Figure 5b), while there were no significant differences for soil TOM among the three soil AP levels (Figure 5a). Generally, the application of fertilizers can directly increase the soil OC and release more WSOC when decomposition by microorganisms occurs (Pospíšilová

et al. 2011, Burns et al. 2013). In addition, the C/N ratio in this study increased significantly with the increase of soil AP level (Figure 6c), indicating that the soil C/N ratio can respond sensitively to soil nutrient changes, and it also verified the importance of WSOC as an index for evaluating soil fertility. Moreover, because of higher soil C/N ratio and N fixation (Gu et al. 2018), soil N loss in the P₂ was less than that in the P₀ or P₁. For the above reasons, significantly higher soil N in the P₂ was observed than in the P₀ or P₁.

The mineralization of P by soil microorganisms is greater than the immobilization due to a low C/P ratio. However, more P is fixed than mineralized, leading to P competition between soil microorganisms and plants with higher C/P ratio (Zhang et al. 2014). High AP reduced the C/P ratio in the P₁ and P₂ soil with the direct P supply, indicating that high P may indirectly improve organic P mineralization mediated

Table 2. Correlation analysis between soil chemical properties and soil enzymatic activities

	OC	NH ₄ ⁺ -N	NO ₃ ⁻ -N	AP	WSOC	WSTN	C/P	C/N	N/P	Invertase	Cellulase	Urease
NH ₄ ⁺ -N	ns											
NO ₃ ⁻ -N	ns	ns										
AP	-0.507*	-0.636**	-0.704**									
WSOC	-0.556*	-0.517*	-0.633**	0.950**								
WSTN	0.476*	ns	0.748**	-0.512*	ns							
C/P	0.488*	0.471*	0.708**	-0.941**	-0.912**	ns						
C/N	-0.582*	-0.517*	-0.806**	0.782**	0.730**	-0.916**	-0.734**					
N/P	0.601**	0.483*	0.836**	-0.777**	-0.717**	0.927**	0.727**	-0.992**				
Invertase	0.693**	0.704**	ns	-0.754**	-0.739**	0.481*	0.653**	-0.680**	0.668**			
Cellulase	0.564*	0.511*	0.818**	-0.798**	-0.763**	0.747**	0.826**	-0.892**	0.880**	0.660**		
Urease	0.625**	0.505*	ns	-0.769**	-0.777**	ns	0.802**	-0.604**	0.590**	0.781**	0.666**	
ALP	ns	-0.623**	ns	0.893**	0.864**	ns	-0.817**	0.591**	-0.562*	-0.684**	-0.626**	-0.725**

ns – no significance at the $P < 0.05$; * $P < 0.05$; ** $P < 0.01$; $n = 18$, $R_{0.05} = 0.4683$; $R_{0.01} = 0.5897$; OC – organic carbon; AP – available phosphorus; WSOC – water-soluble organic carbon; WSTN – water-soluble total nitrogen; ALP – alkaline phosphatases

<https://doi.org/10.17221/296/2019-PSE>

by soil microorganism. This resulted in a significant negative relationship between the soil AP and C/P ratio (Table 2). In addition, the osmotic metabolites, including sugars, amino acids and carboxylic acids, increase under AP deficient conditions in plants (Spohn and Kuzyakov 2013), possibly to significantly improve soil microbial-mediated mineralization of organic P to meet the demand of microorganism.

The soil N/P ratio can indicate the relative restriction of soil nutrients to plant growth. N is relatively sufficient for a higher N/P ratio and P is relatively sufficient for a lower N/P ratio (Liu et al. 2010). The P₂ had a high soil AP and low soil C/P or N/P ratios, suggesting that the supply of soil mineralized P was sufficient, and the N availability in cotton soil was low. Hence, WSOC had significant positive correlations with soil AP, and negative correlations with the C/P and N/P ratios (Table 2), suggesting that soil microorganisms were likely to be restricted by P.

Soil enzymes. Soil enzymes are often involved in nutrient cycling and availability, SOC decomposition and synthesis, and soil fertility and quality, as indicators for microbiological and biochemical processes (Burns et al. 2013, Moghimian et al. 2017). In this study, the activities of invertase, cellulase and urease in cotton soil decreased significantly in the P₀. It is possible that the P deficient in the P₀ would induce the growth of soil microorganisms, stimulate the activity of soil invertase, cellulase and urease, and increase the C and N utilization in soil, which is consistent with the previous studies (Moghimian et al. 2017). As positive relationships were observed between invertase and cellulase activities with OC in cotton soil, and inorganic-N (mainly NH₄⁺-N); soil C/N ratio is negatively correlated with soil AP (Table 2). Although inorganic-P is given to have a feedback inhibition for ALP (Olander and Vitousek 2000), it was not observed to inhibit soil ALP activity with increasing soil AP (Figure 4). One possible reason is that most of the ALPs were not restrained by high P because they have stable interactions with colloids (Dilly and Nannipieri 2001).

ALPs can hydrolyse phosphate monoesters to obtain orthophosphate, and are only produced by soil microorganisms (Spohn and Kuzyakov 2013, Fraser et al. 2017). In addition, soil microorganisms, produced by ALPs, are more susceptible to P fertilization than other phosphatases, which is evident from the raised ALP activity due to P fertilization (Chen et al. 2014); this corresponds with the results of our study, which observed significant positive correlations between soil

AP and ALP (Table 2). Hence, the soil C/P ratio had a significant negative correlation with the ALP activity.

Urease is mainly derived from plants and microorganisms; it has a crucial role in the N cycling process (Li et al. 2014). When NH₄⁺ is the preferred N source, the synthesis of urease in cells is blocked (Geisseler et al. 2010). Nevertheless, an alternative N source or urea activates urease production (Mobley et al. 1995). In this study, the activity of urease had a significant positive correlation with soil NH₄⁺-N, N/P and C/P, and a negative relationship with soil C/N, suggesting that the high soil urease activity for the P₀ indicates a larger N cycle potential due to a lower C/N ratio. In addition, soil NH₄⁺-N and NO₃⁻-N for the P₁ and P₂ were lower than those for the P₀, possibly due to an influence of high AP on soil N availability.

Acknowledgement

We thank all members of the Experimental Station in Baibi town at Institute of Cotton Research of Chinese Academy of Agricultural Sciences for their help with cotton management. We also thank the reviewers for their valuable suggestions and comments on the manuscript.

REFERENCES

- Adetunji A.T., Lewu F.B., Mulidzi R., Ncube B. (2017): The biological activities of β -glucosidase, phosphatase and urease as soil quality indicators: A review. *Journal of Soil Science and Plant Nutrition*, 17: 794–807.
- Bowles T.M., Acosta-Martínez V., Calderón F., Jackson L.E. (2014): Soil enzyme activities, microbial communities, and carbon and nitrogen availability in organic agroecosystems across an intensively-managed agricultural landscape. *Soil Biology and Biochemistry*, 68: 252–262.
- Burns R.G., DeForest J.L., Marxsen J., Sinsabaugh R.L., Stromberger M.E., Wallenstein M.D., Weintraub M.N., Zoppini A. (2013): Soil enzymes in a changing environment: Current knowledge and future directions. *Soil Biology and Biochemistry*, 58: 216–234.
- Chen B.L., Jiang P.A., Sheng J.D. (2014): Effect of phosphate fertilizers on soil available phosphorus and soil enzyme activities in cotton field. *Chinese Journal of Soil Science*, 45: 185–188. (In Chinese)
- Cordell D., White S.B. (2013): Sustainable phosphorus measures: Strategies and technologies for achieving phosphorus security. *Agronomy*, 3: 86.
- DeForest J.L., Smemo K.A., Burke D.J., Elliott H.L., Becker J.C. (2012): Soil microbial responses to elevated phosphorus and pH in acidic temperate deciduous forests. *Biogeochemistry*, 109: 189–202.

- Dilly O., Nannipieri P. (2001): Response of ATP content, respiration rate and enzyme activities in an arable and a forest soil to nutrient additions. *Biology and Fertility of Soils*, 34: 64–72.
- Djodjic F., Mattsson L. (2013): Changes in plant-available and easily soluble phosphorus within 1 year after P amendment. *Soil Use and Management*, 29: 45–54.
- Fraser T.D., Lynch D.H., Gaiero J., Khosla K., Dunfield K.E. (2017): Quantification of bacterial non-specific acid (*phoC*) and alkaline (*phoD*) phosphatase genes in bulk and rhizosphere soil from organically managed soybean fields. *Applied Soil Ecology*, 111: 48–56.
- Geisseler D., Horwath W.R., Joergensen R.G., Ludwig B. (2010): Pathways of nitrogen utilization by soil microorganisms – A review. *Soil Biology and Biochemistry*, 42: 2058–2067.
- Griffiths B.S., Spillies A., Bonkowski M. (2012): C:N:P stoichiometry and nutrient limitation of the soil microbial biomass in a grazed grassland site under experimental P limitation or excess. *Ecological Processes*, 1: 6.
- Gu Y.J., Han C.L., Kong M., Shi X.Y., Zdruli P., Li F.M. (2018): Plastic film mulch promotes high alfalfa production with phosphorus-saving and low risk of soil nitrogen loss. *Field Crops Research*, 229: 44–54.
- Hai L., Li X.G., Li F.M., Suo D.R., Guggenberger G. (2010): Long-term fertilization and manuring effects on physically-separated soil organic matter pools under a wheat-wheat-maize cropping system in an arid region of China. *Soil Biology and Biochemistry*, 42: 253–259.
- Li Q., Liang J.H., He Y.Y., Hu Q.J., Yu S. (2014): Effect of land use on soil enzyme activities at karst area in Nanchuan, Chongqing, Southwest China. *Plant, Soil and Environment*, 60: 15–20.
- Liu Z.F., Fu B.J., Zheng X.X., Liu G.H. (2010): Plant biomass, soil water content and soil N:P ratio regulating soil microbial functional diversity in a temperate steppe: A regional scale study. *Soil Biology and Biochemistry*, 42: 445–450.
- Mühlbachová G., Čermák P., Vavera R., Káš M., Pechová M., Marková K., Hlušek J., Lošák T. (2018): Phosphorus availability and spring barley yields under graded P-doses in a pot experiment. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 66: 111–118.
- Mi J.Z., Gregorich E.G., Xu S.T., McLaughlin N.B., Liu J.H. (2018): Effects of a one-time application of bentonite on soil enzymes in a semi-arid region. *Canadian Journal of Soil Science*, 98: 542–555.
- Mobley H.L., Island M.D., Hausinger R.P. (1995): Molecular biology of microbial ureases. *Microbiological Reviews*, 59: 451–480.
- Moghimian N., Hosseini S.M., Kooch Y., Darki B.Z. (2017): Impacts of changes in land use/cover on soil microbial and enzyme activities. *Catena*, 157: 407–414.
- Niu Y.F., Chai R.S., Jin G.L., Wang H., Tang C.X., Zhang Y.S. (2013): Responses of root architecture development to low phosphorus availability: A review. *Annals of Botany*, 112: 391–408.
- Olander L.P., Vitousek P.M. (2000): Regulation of soil phosphatase and chitinase activity by N and P availability. *Biogeochemistry*, 49: 175–191.
- Pospíšilová L., Formánek P., Kučerek J., Liptaj T., Lošák T., Martensson A. (2011): Land use effects on carbon quality and soil biological properties in Eutric Cambisol. *Acta Agriculturae Scandinavica, Section B – Soil and Plant Science*, 61: 661–669.
- Powlson D.S., Gregory P.J., Whalley W.R., Quinton J.N., Hopkins D.W., Whitmore A.P., Hirsch P.R., Goulding K.W.T. (2011): Soil management in relation to sustainable agriculture and ecosystem services. *Food Policy*, 36: S72–S87.
- Schilling G., Gransee A., Deuhel A., Ležovič G., Ruppel S. (1998): Phosphorus availability, root exudates, and microbial activity in the rhizosphere. *Zeitschrift für Pflanzenernährung und Bodenkunde*, 161: 465–478.
- Shen J.B., Yuan L.X., Zhang J.L., Li H.G., Bai Z.H., Chen X.P., Zhang W.F., Zhang F.S. (2011): Phosphorus dynamics: From soil to plant. *Plant Physiology*, 156: 997–1005.
- Spohn M., Kuzyakov Y. (2013): Distribution of microbial- and root-derived phosphatase activities in the rhizosphere depending on P availability and C allocation – Coupling soil zymography with ¹⁴C imaging. *Soil Biology and Biochemistry*, 67: 106–113.
- Wu H. (2013): Study on the Cultivation Techniques of Sweet Sorghum in the Saline Soil of Yellow River Delta. Ji'nan, Shandong Normal University.
- Yang J.E., Jacobsen J.S. (1990): Soil inorganic phosphorus fractions and their uptake relationships in calcareous soils. *Soil Science Society of America Journal*, 54: 1666–1669.
- Zhang L., Ding X.D., Chen S.F., He X.H., Zhang F.S., Feng G. (2014): Reducing carbon: Phosphorus ratio can enhance microbial phytin mineralization and lessen competition with maize for phosphorus. *Journal of Plant Interactions*, 9: 850–856.
- Zheng L., Huang Z., Huang C. (2015): The study of soil test and formula fertilization index system on sweet potato. *Journal of Agriculture*, 5: 19–24. (In Chinese)

Received on May 30, 2019

Accepted on July 17, 2019

Published online on July 29, 2019