

Rove Beetles (Coleoptera: Staphylinidae) in an Apple Orchard

ALOIS HONĚK¹, MATÚŠ KOČIAN² and ZDENKA MARTINKOVÁ³

¹Department of Entomology, Division of Plant Health and ³Department of Plant Ecology and Weed Science, Division of Agroecology, Crop Research Institute, Prague, Czech Republic;

²Department of Ecology, Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Prague, Czech Republic

Abstract

HONĚK A., KOČIAN M., MARTINKOVÁ Z. (2012): **Rove beetles (Coleoptera: Staphylinidae) in an apple orchard.** Plant Protect Sci., **48**: 116–122.

Many rove beetle (Coleoptera: Staphylinidae) species are carnivorous. Despite their positive role in the biological control of agricultural arthropod pests rove beetles are relatively poorly studied in general, and little is known about their habitat associations including their occurrence and seasonal activity in the apple-orchard environment. In 1994, abundance and composition of adult staphylinid taxocenosis was established in a 12-years-old apple orchard at Prague-Ruzyně, and compared with taxocenoses inhabiting nearby grassy ridge, winter wheat field and forest. The staphylinids were collected throughout the vegetation season using unbaited pitfall traps. Staphylinid activity was highest in the apple orchard where 28 species were established in the total sample of 1238 individuals. Dominant species *Drusilla canaliculata* (F.), *Ocyopus nero semilanatus* Müller, *Dinarea angustula* (Gyllenhal) and *Oxytelus insecatus* Gravenhorst together represented 93% of the total sample. The former two species dominated also the taxocenosis of grassy ridge and could disperse to the orchard from surrounding swards. Most established species are polyphagous predators or parasitoids that might contribute to the biological control of some orchard pests.

Keywords: community; grass sward; field; forest; taxocenosis; seasonal dynamics; abundance; activity; diversity; similarity

Rove beetles (Staphylinidae) are the most species rich family of beetles (Coleoptera) in the Czech Republic (JELÍNEK 1993) and are an important part of the arthropod fauna of terrestrial habitats (BOHAC 1999). A large number of staphylinid species are predators consuming insects, arachnids, molluscs, and nematodes. Other food specialisations include ectoparasitism on insects, pollen and spore eating and scavenging (BALDUF 1935; CLAUSEN 1940). Therefore the family takes part in the complex of biological regulators of crop pests.

Since rove beetles have been relatively poorly studied in general, it is no wonder that few studies concern the abundance and composition of their taxocenoses (total of the species of the fam-

ily Staphylinidae occurring together in the same association, LINCOLN *et al.* 1998) in agricultural landscape. This becomes particularly evident in comparison with another “established” group of predators, ground beetles (Coleoptera: Carabidae). A flood of publications concerning different aspects of biology of carabids appears each year, most of them concerning agroecosystems (KOTZE *et al.* 2011). Sampling of staphylinids on the ground surface is less easy than in carabids. Pitfall traps used to sample the ground surface fauna catch disproportionately more carabids than staphylinids and the latter thus appear less numerous than they in fact are (LANG 2000; SHAH *et al.* 2003; VOLKMAR *et al.* 2004; HOLLAND *et al.* 2007). An important difficulty

Supported by the Ministry of Agriculture of the Czech Republic, Project No. 0002700604.

of community studies is also uneasy identification of staphylinid species. As a result, the study of staphylinid taxocenoses in agroecosystems was frequently neglected (ELLIOTT *et al.* 2006).

Most of the available studies of agroecosystems concern staphylinid taxocenoses of field habitats and/or abandoned or marginal sites important for staphylinid overwintering (LYS & NENTWIG 1994; HONEK 1997; KROOSS & SCHAEFER 1998a; GEIGER *et al.* 2009; EYRE & LEIFERT 2011). Orchards with only a few recent studies are a typical example of neglected habitats (MAJZLAN & HOLECOVA 1993; BALOG & MARKO 2007; BALOG *et al.* 2008, 2009). In this study we report on pitfall trap sampling of a taxocenosis of staphylinid beetles in a 12-years-old orchard abandoned for six years. The results were compared with a parallel study at three habitats typical of the surrounding agricultural landscape – winter wheat crop, a grassy ridge on its margin, and oak forest. The scope of the work was to establish typical species of orchard fauna and possible sources of their origin in agricultural landscape.

MATERIAL AND METHODS

Area and sampling. The occurrence and seasonal dynamics of rove beetles were established in parallel at four sites: in an apple orchard, grassy ridge, winter wheat crop and adjacent oak forest. This study concerns beetles collected in a 12-years-old apple orchard at Prague-Ruzyně (centred around 50°05'16.5"N, 14°17'58.9"E, altitude of 340 m a.s.l.). The rectangular 100 × 250 m orchard was situated on a mild south-facing slope and bordered by a 5–10 m wide grassy strip, bordered outside by a 2 m wide weed overgrown strip (*Arctium tomentosum*, *Echinops sphaerocephalus*, *Urtica dioica* and other perennial weeds). The garden was surrounded by fields and, on the southern side, by buildings. The ground surface arthropods were sampled using pitfall traps, plastic cups 8 cm in diameter (50 cm² outlet area) and 11 cm deep. The cups were dug into the soil, with the rim flush with the soil surface. No bait was used. A few pieces of soil on the bottom of the cups provided shelter for the trapped arthropods. Five pitfall traps spaced 50 m were placed along a 230 m E–W transect situated in the centre of the orchard parallelly to its long side. The traps were operated from April 12 to October 14, 1994 and emptied in 7–10 d intervals. In the parallel study (HONĚK

& KOCIAN 2003) rove beetles were sampled in a suburban agricultural area ca 1000 m away from the orchard, along a 400 m E–W transect crossing a grassy ridge (8 m wide, centred at 50°05'57.4"N, 14°17'51.6"E), winter wheat field (transect 310 m long, 50°05'57.2"N, 14°17'58.8"E) and oak forest (transect 80 m long, 50°05'58.0"N, 14°18'10.1"E). The traps were operated from April 15 to September 11, 1996 and emptied in 2–3 d intervals.

Data processing. Since the catches were low, numbers of individuals of particular species in each trap were summed over the entire catching period. At each sampling site (apple orchard, grassy ridge, wheat field, and oak forest) we listed numbers of individuals belonging to each species. From this data we established species richness (number of species at each site) and calculated their diversity using Shannon-Wiener index H' as follows:

$$H' = -\sum p_i \times \log p_i$$

where:

p_i – proportion of the total number of individuals captured at a site composed of species i

The value of H' increases with the increasing number of species and decreasing differences in their abundance. Similarity between samples caught at particular sampling sites was calculated using Renkonen coefficient Re as follows:

$$Re = \frac{\sum \min(x_{ip}, x_{jp})}{\sum x_{ip}}$$

where:

x_{ip} , x_{jp} – proportions of species p in the samples of sites i and j

The coefficient ranges between 0 when there are no species common to both samples and 1 when the species composition and proportions in samples are identical.

Chi-square test was used to establish an association between taxocenosis composition and sampling site, with species catches as rows and sites as columns of the contingency table. For each pair of taxocenoses (orchard vs. ridge, orchard vs. field, orchard vs. forest) the species were ranked by abundance and the analysis was performed of as many species as to meet the condition that no expected value in the contingency table is < 1 and $< 20\%$ is less than 5.

To establish seasonal trends in abundance the average catch of abundant species per trap/day was calculated for periods between successive trap

servicing. Average annual catch per trap was calculated to compare abundance (total number of staphylinids collected over the vegetation season) at particular sites.

RESULTS

In the apple orchard, a total of 1236 individuals of 28 species were captured over the sampling




Figure 1. Seasonal activity of abundant staphylinid species in the apple orchard at Prague-Ruzyně in 1994 (A) *Ocyptus nero semilanatus*, (B) *Oxytelus insecatus*, (C) *Dinarea angustula*, (D) *Drusilla canaliculata*. The bars indicate the activity of species (mean number of individuals/trap/day) captured over the period terminated by Julian day of emptying the trap




Figure 2. Species diversity (Shannon-Wiener index) and numbers of species (in brackets) in adult staphylinid taxocenoses of apple orchard (sampled in 1994) and taxocenoses of grassy ridge, winter wheat field and oak forest (sampled in 1996) at Prague-Ruzyně, and their similarity (Renkonen coefficient, bold numbers at the connecting lines between the labels of taxocenoses)

period (Table 1). Four dominant species, *Drusilla canaliculata* (F.), *Dinarea angustula* (Gyllenhal), *Oxytelus insecatus* Gravenhorst and *Ocyptus nero semilanatus* Müller, represented 93% of the total sample. The majority of species was represented by one (43% of the total established) or two individuals (21%). The staphylinid beetles were active through the whole vegetation period but their seasonal activity differed between species (Figure 1). *Ocyptus nero* was captured in spring and autumn, while other species have a unimodal pattern of activity peaking in June (*O. insecatus*) or July (*D. angustula*, *D. canaliculata*). The activity density (annual catches) was greater in two marginal traps placed near the grassy strip bordering the orchard (327.5 ± 17.50 individuals) than in the traps placed in the centre of the orchard (194.3 ± 8.82 individuals).

The staphylinid taxocenosis of apple orchard differed from the taxocenoses of other sites situated along the nearby transect crossing agricultural landscape (Table 1). The composition of orchard taxocenosis was significantly different from that of grassy ridge ($df = 5$, $\chi^2 = 239.0$, $P < 0.001$), winter wheat field ($df = 11$, $\chi^2 = 1156.3$, $P < 0.001$) and oak forest ($df = 11$, $\chi^2 = 1481.3$, $P < 0.001$). The annual catches per trap at the latter sites were by 38–77% smaller than in the orchard. Species diversity and number were greater in the forest and field taxocenosis but lower at the grassy ridge. The taxocenosis of apple orchard was similar to the taxocenosis of grassy ridge, more than the

Table 1 Staphylinid species (number of individuals in the annual sample) captured in the apple orchard and near habitats of agricultural landscape, grassy ridge, winter wheat field and oak forest, total sample of particular habitats (Total N), number of traps run at the habitat (N traps), and annual catch per trap (N/trap)

Species	Orchard	Ridge	Field	Forest
<i>Aleochara bipustulata</i> (L.)	5			1
<i>Aleochara curtula</i> (Goeze)			2	
<i>Aleochara haematodes ripicola</i> Mulsant et Rey	19		4	5
<i>Aleochara inconspicua</i> Aubé	1			
<i>Aleochara laevigata</i> Gyllenhal			1	
<i>Aleochara sparsa</i> Heer				1
<i>Aloconota gregaria</i> (Erichson)			4	
<i>Amischa analis</i> (Gravenhorst)			1	
<i>Amischa cavifrons</i> Sharp			1	1
<i>Amischa soror</i> (Kraatz)			2	
<i>Atheta amplicollis</i> (Mulsant)			1	1
<i>Atheta crassicornis</i> (F.)				20
<i>Atheta fungi</i> (Gravenhorst)			17	39
<i>Atheta gagatina</i> Baudi				6
<i>Atheta laticollis</i> (Stephens)			2	3
<i>Atheta livida</i> Mulsant et Rey				3
<i>Atheta pittionii</i> Sheerpeltz			1	3
<i>Atheta sodalis</i> (Erichson)				1
<i>Atheta subtilis</i> (Scriba)			11	1
<i>Atheta triangulum</i> (Kraatz)		3	5	1
<i>Bolitobius castaneus</i> (Stephens)	2			
<i>Bolitobius formosus</i> (Gravenhorst)		1	1	2
<i>Dinarea angustula</i> (Gyllenhal)	418		2	
<i>Dinarea linearis</i> (Gravenhorst)			1	1
<i>Drusilla canaliculata</i> (F.)	473	229	133	2
<i>Enalodroma hepatica</i> (Erichson)				3
<i>Gabrius osseticus</i> (Kolenati)	1			
<i>Ilyobates subopacus</i> Palm			5	
<i>Lathrimeum athrocephalum</i> (Gyllenhal)				1
<i>Lathrobium fulvipenne</i> Gravenhorst	1		8	
<i>Lathrobium pallidum</i> Nordmann	2			
<i>Leptacinus sulcifrons</i> (Stephens)			1	
<i>Mniobates forticornis</i> (Boisduval)	2		2	
<i>Ocypus fulvipennis</i> Erichson	1			
<i>Ocypus fuscatus</i> (Gravenhorst)		1	1	
<i>Ocypus melanarius</i> Heer	17			2
<i>Ocypus nero semilanatus</i> J. Müller	100	37	7	1
<i>Omaliium caesum</i> Gravenhorst		10	4	130
<i>Omaliium rivulare</i> (Paykull)				4
<i>Ontholestes haroldi</i> (Eppelsheim)	1			
<i>Othius punctulatus</i> (Goeze)				1
<i>Oxypoda abdominalis</i> Mannerheim				2
<i>Oxypoda haemorrhoea</i> Mannerheim		1		1
<i>Oxypoda lividipennis</i> Mannerheim			1	5
<i>Oxypoda longipes</i> Mulsant et Rey		5	2	13
<i>Oxypoda spectabilis</i> Märkel				1
<i>Oxypoda vicina</i> Kraatz			1	
<i>Oxytelus insecatus</i> Gravenhorst	163	1	8	

Table 1 to be continued

Species	Orchard	Ridge	Field	Forest
<i>Oxytelus mutator</i> Lohse			1	3
<i>Oxytelus rugosus</i> (F.)	2		2	
<i>Parabemus fossor</i> (Scopoli)		1		1
<i>Philonthus carbonarius</i> (Gravenhorst)	2		4	
<i>Philonthus chalceus</i> Stephens			1	1
<i>Philonthus cognatus</i> Stephens	1	5	323	
<i>Philonthus decorus</i> (Gravenhorst)		1	2	118
<i>Philonthus laminatus</i> (Creutzer)			5	
<i>Platarea dubiosa</i> (Benick)		7	4	2
<i>Platarea nigriceps</i> (Marsham)	13			
<i>Quedius limbatus</i> (Heer)				1
<i>Rugilus orbiculatus</i> (Paykull)	1		1	
<i>Rugilus subtilis</i> (Erichson)	1	1		
<i>Sepedophilus marshami</i> (Stephens)		1	1	1
<i>Sepedophilus testaceus</i> (F.)				1
<i>Staphylinus stercorarius</i> Olivier			1	
<i>Sunius melanocephalus</i> (F.)		1	1	
<i>Tachinus rufipes</i> (DeGeer)		2	23	
<i>Tachyporus chrysomelinus</i> (L.)	1			
<i>Tachyporus hypnorum</i> (F.)	2		12	
<i>Tachyporus nitidulus</i> (F.)	1			
<i>Tachyporus obtusus</i> (L.)			1	
<i>Tachyporus pusillus</i> Gravenhorst			1	
<i>Tachyporus solutus</i> (Erichson)	1		1	
<i>Xantholinus linearis</i> (Olivier)	3		1	
<i>Xantholinus longiventris</i> Heer	1			
<i>Zyras humeralis</i> (Gravenhorst)				2
<i>Zyras limbatus</i> (Paykull)	1	1		
Total N	1236	308	614	385
N traps	5	2	11	3
N/trap	247.2	154.0	55.8	128.3

taxocenoses of the field and the forest (Figure 2). This was mainly caused by sharing two dominant species, *D. canaliculata* and *O. nero semilanatus*. Other dominant species (*D. angustula*, *O. insecatus*) were typical of the orchard. In the orchard the common species shared with the grassy ridge were particularly abundant in marginal traps placed near the grassy strip bordering the orchard compared to the central traps. Consequently, the similarity of grassy ridge taxocenosis and taxocenosis of marginal orchard traps (Re = 0.532) was greater than similarity to the taxocenosis of central orchard traps (Re = 0.399). Diversity of staphylinid taxocenoses of the field and forest was, despite the lower total activity density of staphylinids at these sites, greater than in the orchard (Figure 2),

because of the greater species richness of these taxocenoses (Table 1).

DISCUSSION

The staphylinid activity in the apple orchard was higher compared to field and forest sites representing typical habitats of the surrounding agricultural landscape. This contrasted with higher species diversity and richness at the latter sites. Low diversity in the orchard was caused by dominance of a few species, two of which, *D. angustula* and *O. insecatus*, were typical of the orchard. The other dominant orchard species, *D. canaliculata* and *O. nero semilanatus*, were also typical of the

taxocenosis of grassy ridge and their common occurrence at both sites was responsible for greater similarity, in terms of Renkonen coefficient, between grassy ridge and orchard taxocenoses than between any other pair of taxocenoses. This similarity was, however, rather apparent because rare species were mostly different at both sites. *Drusilla canaliculata* is a generalist of non-forest areas while *O. nero semilanatus* probably disperses into the orchard from the marginal grassy strip similarly as it disperses from the grassy ridge into the marginal area of wheat stand (HONĚK & KOČIAN 2003).

The species composition of staphylinid taxocenoses of the Czech Republic partly matched the taxocenoses established in other studies of central Europe. The richness of staphylinid taxocenosis in Hungarian apple and pear orchards (BALOG & MARKO 2007; BALOG *et al.* 2008, 2009) was about 10 times greater than in our study, but dominant species *D. canaliculata* and *D. angustula* and subdominant *Xantholinus linearis* (Olivier) were similar to our study. *Omalius caesus* Gravenhorst dominant in Hungary was captured in our study in the grassy ridge and was very abundant in the oak forest. Staphylinid taxocenoses established in orchards of remote countries or in other habitats were not similar to taxocenoses of our study (PIETRASZKO & DECLERCQ 1978; GOOD & GILLER 1991; SOBOLEVA-DOKUCHAEVA *et al.* 2002; JUEN *et al.* 2003; BALOG *et al.* 2011). This was caused by the presence of species not established in this study or by preference of common species to different habitats.

Most species established in the orchard are predators. Predation was demonstrated for *D. canaliculata* (NOVÁK 1958; BABENKO 1985), *D. angustula* (BALDUF 1935) and congeners of *Ocypus* (KROOSS & SCHAEFER 1998b; BONACCI *et al.* 2006), *Oxytelus* (ACHIANO & GILIOME 2006), *Philonthus* (SEYMOUR & CAMPBELL 1993), *Tachyporus* (BALOG *et al.* 2011), and *Xantholinus* (BALDUF 1935) while *Atheta* spp. was shown to switch between adult predation and larval ectoparasitism (CLAUSEN 1940). Because of wide polyphagy, it is difficult to estimate the effect of staphylinid predation on orchard pests. Food items of staphylinids captured in the apple orchard or their above-mentioned congeners included ants, carabid larvae, aphids, dipteran eggs and pupae, mites, millipedes, slugs and other prey. Which species of orchard pests are eaten by staphylinids and in what quantities remains to be studied. Nevertheless, staphylinids are certainly an important component in the web of

relations that maintain balance among phytophagans, zoophagans and scavengers of production orchards. Because of their abundance the role of staphylinids in orchards may be even more important than at other habitats of the agroecosystem included in this study. The importance of staphylinid beetles for the control of pests in agroecosystems should be further studied.

References

- ACHIANO K.A., GILIOME J.H. (2006): Feeding behaviour of the potential predators of the house flies, *Musca domestica* L. and *Fannia canicularis* (L.) (Diptera: Muscidae). *African Entomology*, **14**: 69–75.
- BABENKO A.S. (1985): The biology of *Astilbus canaliculatus* (Coleoptera, Staphylinidae) in the south of West Siberia. *Zoologicheskii Zhurnal*, **64**: 993–996.
- BALDUF W.V. (1935): The Bionomics of Entomophagous Coleoptera. John S. Swift Co., St. Louis.
- BALOG A., MARKO V. (2007): Chemical disturbances effects on community structure of rove beetles (Coleoptera: Staphylinidae) in Hungarian agricultural fields. *North-Western Journal of Zoology*, **3**: 67–74.
- BALOG A., MARKO V., SZARVAS P. (2008): Dominance, activity density and prey preferences of rove beetles (Coleoptera: Staphylinidae) in conventionally treated Hungarian agro-ecosystems. *Bulletin of Entomological Research*, **98**: 343–353.
- BALOG A., MARKO V., IMRE A. (2009): Farming system and habitat structure effects on rove beetles (Coleoptera: Staphylinidae) assembly in Central European apple and pear orchards. *Biologia (Bratislava)*, **64**: 343–349.
- BALOG A., SZENASI A., SZEKERES D., PALINKAS Z. (2011): Analysis of soil dwelling rove beetles (Coleoptera: Staphylinidae) in cultivated maize fields containing the Bt toxins, Cry34/35Ab1 and Cry1F × Cry34/35Ab1. *Bio-control Science and Technology*, **21**: 293–297.
- BOHAC J. (1999): Staphylinid beetles as bioindicators. *Agriculture Ecosystems and Environment*, **74**: 357–372.
- BONACCI T., MASSOLO A., BRANDMAYR P., BRANDMAYR T.Z. (2006): Predatory behaviour on ground beetles (Coleoptera: Carabidae) by *Ocypus olens* (Muller) (Coleoptera: Staphylinidae) under laboratory conditions. *Entomological News*, **117**: 545–551.
- CLAUSEN C.P. (1940): *Entomophagous insects*. McGraw-Hill, New York, London
- ELLIOTT N., TAO F.L., GILES K.L., ROYER T.A., GREENSTONE M.H., SHUFRAN K.A. (2006): First quantitative study of rove beetles in Oklahoma winter wheat fields. *BioControl*, **51**: 79–87.

- EYRE M.D., LEIFERT C. (2011): Crop and field boundary influences on the activity of a wide range of beneficial invertebrate groups on a split conventional/organic farm in northern England. *Bulletin of Entomological Research*, **101**: 135–144.
- GEIGER F., WAECKERS F.L., BIANCHI F.J.J.A. (2009): Hibernation of predatory arthropods in semi-natural habitats. *BioControl*, **54**: 529–535.
- GOOD J.A., GILLER P.S. (1991): The effect of cereal and grass management on staphylinid (Coleoptera) assemblages in south-west Ireland. *Journal of Applied Ecology*, **28**: 810–826.
- HOLLAND J.M., THOMAS C.F.G., BIRKETT M., SOUTHWAY S. (2007): Spatio-temporal distribution and emergence of beetles in arable fields in relation to soil moisture. *Bulletin of Entomological Research*, **97**: 89–100.
- HONEK A. (1997): The effect of plant cover and weather on the activity density of ground surface arthropods in a fallow field. *Biological Agriculture and Horticulture*, **15**: 203–210.
- HONĚK A., KOČIAN M. (2003): Importance of woody and grassy areas as refugia for field Carabidae and Staphylinidae (Coleoptera). *Acta Societatis Zoologicae Bohemicae*, **67**: 71–81.
- JELÍNEK J. (ed.) (1993): Check-list of Czechoslovak insects IV (Coleoptera) – Seznam československých brouků. *Folia Heyrovskyana*, Suppl. 1: 3–172. (in English and Czech).
- JUEN A., STEINBERGER K.H., TRAUGOTT M. (2003): Seasonal change in species composition and size distribution of epigeic predators in a small field. *Entomologia Generalis*, **26**: 259–275.
- KOTZE D.J., BRANDMAYR P., CASALE A., DAUFFY-RICHARD E., DEKONINCK W., KOIVULA M.J., LÖVEI G.L., MOSAKOWSKI D., NOORDIJK J., PAARMAN W., PIZZOLOTTO R., SASKA P., SCHWERK A., SERRANO J., SZYSZKO J., TABOADA A., TURIN H., VENN S., VERMEULEN S., ZETTO T. (2011): Forty years of carabid beetle research in Europe – from taxonomy, biology, ecology and population studies to bioindication, habitat assessment and conservation. *Zookeys*, **100**: 55–148.
- KROOSS S., SCHAEFER M. (1998a): The effect of different farming systems on epigeic arthropods: a five-year study on the rove beetle fauna (Coleoptera: Staphylinidae) of winter wheat. *Agriculture Ecosystems and Environment*, **69**: 121–133.
- KROOSS S., SCHAEFER M. (1998b): How predacious are predators? A study on *Ocypus similis*, a rove beetle of cereal fields. *Annals of Applied Biology*, **133**: 1–16.
- LANG A. (2000): The pitfalls of pitfalls: a comparison of pitfall trap catches and absolute density estimates of epigeal invertebrate predators in arable land. *Anzeiger für Schädlingskunde*, **73**: 99–106.
- LINCOLN R., BOXSHALL G., CLARK P. (1998): *A Dictionary of Acology, Evolution and Systematics*. 2nd Ed. Cambridge University Press, Cambridge.
- LYS J.A., NENTWIG W. (1994): Improvement of the overwintering sites for Carabidae, Staphylinidae and Araneae by strip-management in a cereal field. *Pedobiologia*, **38**: 238–242.
- MAJZLAN O., HOLECOVA M. (1993): Arthropodocenoses of an orchards ecosystem in urban agglomeration. *Ekologia (Bratislava)*, **12**: 121–129.
- NOVÁK B. (1958): Ein Beitrag zur Kenntnis des mütterlichen Instinktes von *Astilbus canaliculatus* F. (mit einigen weiteren ökologischen Bemerkungen). *Sborník Vysoké Školy Pedagogické v Olomouci. Přírodní Vědy*, **5**: 173–186.
- PIETRASZKO R., CLERCQ R. DE (1978): Study of the Staphylinidae fauna in fields of winter wheat in Belgium. *Parasitica*, **34**: 191–198.
- SEYMOUR R.C., CAMPBELL J.B. (1993): Predators and parasitoids of house flies and stable flies (Diptera, Muscidae) in cattle confinements in west central Nebraska. *Environmental Entomology*, **22**: 212–219.
- SHAH P.A., BROOKS D.R., ASHBY J.E., PERRY J.N., WOIWOD I.P. (2003): Diversity and abundance of the coleopteran fauna from organic and conventional management systems in southern England. *Agricultural and Forest Entomology*, **5**: 51–60.
- SOBOLEVA-DOKUCHAEVA I.I., TSHERNYSHEV W.B., AFONINA V.A., TIMOKHOV A.V. (2002): Seasonal dynamics of spatial distribution of mass staphylinid-beetles (Coleoptera, Staphylinidae) in agroecosystems. *Zoologicheskii Zhurnal*, **81**: 451–456.
- VOLKMAR C., HUSSEIN M.L.A., WETZEL T. (2004): Ecological field studies in transgenic maize at Friemar (Thuringia). *Journal of Plant Diseases and Protection (Special Issue)*, **19**: 1017–1024.

Received for publication January 5, 2012

Accepted after corrections May 2, 2012

Corresponding author:

Doc. RNDr. ALOIS HONĚK, CSc., Výzkumný ústav rostlinné výroby, v.v.r., Odbor rostlinolékařství, oddělení entomologie, Drnovská 507, 161 06 Praha 6-Ruzyně, Česká republika
tel. + 420 233 022 269, e-mail: honek@vurv.cz
