

Developmental patterns of *FASN* and *LIPE* mRNA expression in adipose tissue of growing Jinhua and Landrace gilts

Z. MIAO¹, F. ZHU², H. ZHANG¹, X. CHANG¹, H. XIE¹, J. ZHANG¹, Z. XU²

¹College of Animal Science, Henan Institute of Science and Technology, XinXiang, P.R. China

²Key Laboratory for Molecular Animal Nutrition of Ministry of Education, Institute of Feed Science, Zhejiang University, HangZhou, P.R. China

ABSTRACT: The present study was aimed to investigate the developmental patterns of *FASN* (fatty acid synthase) and *LIPE* (lipase, hormone-sensitive) mRNA in adipose tissue in pigs of different breeds and the relation with carcass fat content. Subcutaneous adipose tissue was sampled and total RNA was extracted to determine *FASN* and *LIPE* mRNA levels by semi-quantitative RT-PCR. The results showed that the *FASN* mRNA level increased with age ($P < 0.05$) and Jinhua gilts expressed higher *FASN* mRNA compared with Landrace gilts at 80 and 125 days of age ($P < 0.05$). In addition, Jinhua gilts expressed lower *LIPE* mRNA compared with Landrace gilts at 80 days of age ($P < 0.01$). Furthermore, the ratio of *FASN*/*LIPE* mRNA had a similar model in the two breeds, and was higher in Jinhua gilts than that in Landrace gilts at 80 and 125 days of age ($P < 0.05$). The *FASN* mRNA level was positively related to carcass fat content in Jinhua and Landrace gilts ($r = 0.802$, $P = 0.01$; $r = 0.734$, $P = 0.02$; respectively), and the ratio of *FASN*/*LIPE* expression exhibited significantly positively related carcass fat content ($r = 0.804$, $P = 0.01$; $r = 0.749$, $P = 0.02$; respectively).

Keywords: gilt; carcass fat content; *FASN*; *LIPE*; gene expression; RT-PCR

The Jinhua pig is one of the most important Chinese local breeds characterized by low growth rate and high adiposity (Xu, 1994), while Landrace pigs are known by high growth rate, but low fat deposition. Therefore, these two breeds of pigs can serve as ideal animal models for studying adipose tissue deposition in pigs. Fat deposition in adipose tissue represents a balance between synthesis and degradation (Chilliard, 1993). *FASN* (fatty acid synthase), which is one of the key enzymes in the conversion of acetyl-CoA and malonyl-CoA to triglycerol, plays an important role in *de novo* lipogenesis in mammals (Semenkovich, 1997; Yan et al., 2002; Smith et al., 2003). *FASN* in pig adipose tissues is extremely positively correlated with the weight of carcass fat (Xiong et al.,

2001). *LIPE* (lipase, hormone-sensitive) is the key enzyme catalyzing the hydrolysis of stored triglycerol in adipose tissue into free fatty acid and glycerol, and one of the most important factors for controlling the lipolysis and fat accumulation in animals (Haemmerle et al., 2003; Kazala et al., 2003). Although it is clear that *FASN* and *LIPE* are involved in the regulation of fat deposition by balancing the lipogenesis and lipolysis, reports about the developmental changes of *FASN* and *LIPE* expression in adipose tissue of Jinhua pigs and reports about the breed differences between Jinhua and Landrace pigs are missing. Therefore, Jinhua and Landrace pigs with significant differences in fat deposition were used in the present study to investigate the developmental changes of *FASN*

and *LIPE* expression levels in adipose tissue and their effects on the adipose tissue deposition.

MATERIAL AND METHODS

Animals

This study was approved by the Institutional Animal Care and Use Committee of Zhejiang University. Purebred Jinhua and Landrace piglets (9 gilts of each breed, weaned at 28 days of age) were used in this experiment. Gilts were allocated to two groups according to breed. The feeding experiment lasted 90 days after a 7-day adaptation period. Jinhua and Landrace gilts were reared in the same conditions. All pigs had *ad libitum* access to an experimental diet and water via nipple drinkers. The experimental diets were formulated to meet the NRC (1998) nutrient requirements. Three gilts of each breed were slaughtered at 35, 80 and 125 days of age. The subcutaneous adipose tissue was quickly dissected and frozen in liquid nitrogen, then stored at -70°C until extraction for the total RNA. Left half-carcasses without head, legs and guts (except kidney) were weighed. Adipose tissue in the left half-carcass was dissected and weighed; the carcass fat content was calculated. All the animal experiments were done according to the guidelines of the National Institute of Animal Health (P.R. China) for animal experiments.

Primer design

Primer sequences for *FASN*, *LIPE* and *ACTB* encoding for β -actin were designed using Primer 5.0 software (Lalitha, 2000) on the basis of known sequences deposited in GeneBank (Table 1). The mRNA of *ACTB* was used as an internal standard for the determination of mRNA targeted levels.

Total RNA extraction

Total RNA of subcutaneous adipose tissue was isolated using TRIzol Reagent (Gibco BRL) as described by the manufacturer (Sigma, USA). The extracted RNA was dissolved in DEPC-treated water. The concentration and purity of RNA were checked using a spectrophotometer at 260/280 nm.

cDNA cloning

The synthesis of first strand cDNA was performed using Reverse Transcription System Kit (First Strand cDNA-synthesis Kit, Promega, USA) as described by the manufacturer with oligo-dT primer and using approximately 1 μg of total RNA as template.

Polymerase chain reaction (PCR)

The reverse-transcribed cDNA was amplified with *Taq* DNA polymerase (Promega, USA) by a PCR in a thermocycler (Gene Amp PCR System 9600, Pharmacia, Japan) using paired sense and antisense primers (Table 1). The PCR conditions for *FASN* were as follows: denaturation at 94°C for 2 min, followed by 31 cycles of amplification at 94°C for 50 s, 57°C for 50 s, and 72°C for 1 min, and followed by final extension cycle at 72°C for 10 min. The PCR conditions for *LIPE* were as follows: denaturation at 94°C for 2 min, followed by 31 cycles of amplification at 94°C for 50 s, 59°C for 50 s, and 72°C for 1 min, and followed by final extension cycle at 72°C for 10 min. The PCR conditions for *ACTB* were the same as those for *FASN*.

DNA sequencing and sequence analysis

The PCR products were electrophoresed in 1% agarose gel and selected electrophoresis bands

Table 1. Parameters of gene-specific primers for *FASN*, *LIPE* and *ACTB* genes

Target genes	Accession number	Primer sequence	Product size (bp)
<i>FASN</i>	EF589048	F: 5'-ACCGCCTCTGCAGTGTCTATTAC-3' R: 5'-ACGCCGCCCGAGCCCGAGTG-3'	358
<i>LIPE</i>	AY686758	F: 5'-CCCCGGGCCCCGTCCTCGTC-3' R: 5'-GTTCCCGCCTGCGCTGTCTCCTG-3'	397
<i>ACTB</i>	DQ845171	F: 5'-CCGCACCACTGGCATTGTTCAT-3' R: 5'-CAGCACCGTGTGGCGTAGAGG-3'	460

were purified using QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). The purified amplified products were directly ligated into pUCm-T Vector (Promega, USA) and transformed into *E. coli* JM109. Plasmids were then isolated and purified for DNA sequencing using Wizard Miniprep Kit (Promega, USA) and analysed on ABI PRISM 310 Genetic Analyser (Perkin Elmer, USA). The sequences of amplified fragments were aligned using the BLAST sequence analysis program (<http://www.ncbi.nlm.nih.gov/BLAST>) with the corresponding reported sequences according to which the gene-specific primers were designed. Amplified DNA fragments were 99% identical to the known sequences of *FASN*, *LIPE* and *ACTB* in GenBank. The results indicated that the amplified cDNA fragments of the three genes were gene-specific products.

mRNA expression analysis

The expression of pig *FASN* and *LIPE* mRNA was determined by semi-quantitative RT-PCR (Kousteni et al., 1999) using the housekeeping gene *ACTB* as internal control. The PCR products were electrophoresed on 1% agarose gel. Electrophoresis band intensities of the PCR products were quantified by NIH Image Version 1.62 software (Pharmacia, Japan).

Statistical analyses

All data were analyzed by the ANOVA procedure of SPSS 11.5. Comparison among age groups with the same breed uses one-way analysis of variance with age as the main effect. Differences among

means were tested by Duncan's comparisons. Comparison between breed groups with the same age was determined by *t*-test. Bivariate correlations were used to evaluate the correlation between carcass fat content and gene expression level. All data are presented as mean \pm SEM.

RESULTS

Carcass fat deposition

As shown in Figure 1, the carcass fat percentage showed an age-dependent pattern. It was low at 35 days of age, followed by a significant stepwise increase in the two breeds from 35 to 125 days of age. Jinhua gilts had a higher carcass fat percentage than Landrace gilts during the whole experimental period ($P < 0.05$).

Developmental patterns of *FASN* and *LIPE* mRNA expression

As shown in Figure 2, the developmental changes in *FASN* mRNA expression of adipose tissue in growing Jinhua and Landrace gilts were evaluated by semi-quantitative RT-PCR. The *FASN* mRNA expression level in subcutaneous adipose tissue showed an age-dependent pattern. It was low at 35 days of age, followed by a significant stepwise increase with age in the two breeds ($P < 0.05$). In addition, the *FASN* mRNA expression was higher in Jinhua gilts than that in Landrace gilts at 80 and 125 days of age ($P < 0.05$).

As shown in Figure 3, the *LIPE* mRNA expression in subcutaneous adipose tissue did not dif-




Figure 1. Developmental changes in carcass fat content in growing Jinhua and Landrace gilts

LD = Landrace gilts; JH = Jinhua gilts means without a common letter (small letters for Jinhua pigs and capital letters for Landraces) differ significantly between ages, $P < 0.05$

Single asterisk * indicates significant differences ($P < 0.05$) between breeds at the same age




Figure 2. Expression of *FASN* mRNA in adipose tissue of pigs from 35 to 125 days of age; (A) electrophoresis of RT-PCR products for *FASN* and *ACTB* genes in the subcutaneous adipose tissue of Jinhua gilts; (B) electrophoresis of RT-PCR products for *FASN* and *ACTB* genes in the subcutaneous adipose tissue of Landrace gilts; (1), (2), (3): products from the first pig, second pig and third pig in each age group, respectively; (C) the developmental patterns of *FASN* mRNA in Jinhua and Landrace gilts, significant differences between ages ($P < 0.05$) are designated by letters and means without common superscript (capital letters for Landrace gilts and small letters for Jinhua gilts); single asterisk * indicates significant differences ($P < 0.05$) between breeds at the same age

fer from 35 to 125 days of age in the two breeds ($P > 0.05$). Whereas, the *LIPE* mRNA expression level in Jinhua gilts was lower than that in Landrace gilts at 80 days of age ($P < 0.01$).

Developmental patterns of *FASN* and *LIPE* mRNA expression

As shown in Figure 4, the ratio of *FASN* to *LIPE* mRNA in subcutaneous adipose tissue showed an age-dependent pattern. It was low at 35 days of age, followed by a significant increase with age in the two breeds ($P < 0.05$). The *FASN/LIPE* mRNA expres-

sion was higher in Jinhua gilts than that in Landrace gilts at 80 and 125 days of age ($P < 0.05$).

The relationship between *FASN* and *LIPE* mRNA expression and carcass fat content

The relationship between the expression of the two genes and carcass fat content was analyzed by bivariate correlations. As shown in Table 2, the *FASN* expression was positively correlated with carcass fat content in Jinhua gilts (*FASN*/carcass fat content: $r = 0.802$, $P = 0.01$). A similar result was observed in Landrace gilts (*FASN*/carcass fat




Figure 3. The expression of *LIPE* mRNA in adipose tissue of pigs from 35 to 125 days of age; (A) electrophoresis of RT-PCR products for *LIPE* and *ACTB* genes in the subcutaneous adipose tissue of Jinhua gilts; (B) electrophoresis of RT-PCR products for *LIPE* and *ACTB* genes in the subcutaneous adipose tissue of Landrace gilts; (1), (2), (3): products from the first pig, second pig and third pig in each age group respectively; (C) the developmental patterns of *LIPE* mRNA in Jinhua and Landrace gilts, significant differences between ages ($P < 0.05$) are designated by letters and means without common superscript (capital letters for Landrace gilts and small letters for Jinhua gilts); double asterisk ** indicates extreme differences ($P < 0.01$) between breeds at the same age

content: $r = 0.734$, $P = 0.02$). Furthermore, the *FASN* per *LIPE* mRNA and carcass fat content showed a significantly positive relationship in the

two breeds (in Jinhua gilts: $r = 0.804$, $P = 0.01$; in Landrace gilts: $r = 0.749$, $P = 0.02$), whereas the mRNA level of *LIPE* had no significant rela-




Figure 4. The developmental patterns of *FASN/LIPE* mRNA levels in adipose tissue of growing Jinhua and Landrace gilts, significant differences between ages ($P < 0.05$) are designated by letters and means without common superscript (capital letters for Landrace gilts and small letters for Jinhua gilts); single star * indicate significant differences ($P < 0.05$), between breeds at the same age

Table 2. Correlations between *FASN* and *LIPE* mRNA expression levels and carcass fat content in growing Jinhua and Landrace gilts

	Breed	<i>FASN</i> mRNA levels	<i>LIPE</i> mRNA levels	<i>FASN/LIPE</i>
Carcass fat	JH	$r = 0.802$ ($P = 0.01$)	$r = -0.226$ ($P = 0.56$)	$r = 0.804$ ($P = 0.01$)
content (%)	LD	$r = 0.734$ ($P = 0.02$)	$r = -0.443$ ($P = 0.24$)	$r = 0.749$ ($P = 0.02$)

JH = Jinhua gilts; LD = Landrace gilts

tion with carcass fat content in the two breeds ($P > 0.05$).

DISCUSSION

There are many factors affecting fat deposition in adipose tissue such as breed, age and nutrition (Mersmann et al., 1973; Yang, 2005). Jinhua gilts contained more carcass fat compared with Landrace gilts during growth. These results are in agreement with previous reports which indicate that carcass fat content displayed breed differences (Wanger et al., 1999). This difference may be attributed to different phenotypes in the fat deposition of Jinhua and Landrace gilts (Jinhua gilts had a greater ability of fat deposition). The higher lipid accretion in Jinhua pigs is probably related to an increase in extra energy available for lipid synthesis. Similar results were reported when Meishan (White et al., 1995), Iberian pigs (Morales et al., 2003) or Creole pigs (Renaudeau and Mourot, 2007) were compared to conventional lean pigs. In addition, carcass fat content increased with age in the two breeds. These results are in accordance with the results of some studies which indicate that carcass fat content displayed age-dependent patterns (Carr et al., 1978; Friesen et al., 1994; D'Souza et al., 2004; Correa et al., 2006).

Fat accumulation in adipose tissue represents a balance between synthesis and degradation (Chilliard, 1993). In pigs, the majority of lipids in the body is derived from *de novo* fatty acid synthesis (O'Hea and Leveille, 1969) and adipose tissue is considered to be the primary site of *de novo* fatty acid synthesis (O'Hea and Leveille, 1969; Bauman, 1976). *FASN* and *LIPE* play the rate-limiting role in the homeostasis of accumulation and metabolism of triglycerides in adipose tissue, which include both the fatty acid synthesis (lipogenesis) and fat degradation (lipolysis). The *FASN* is highly expressed

in pig adipose tissue and to a much lesser extent in liver. In this study, a positive correlation between the *FASN* mRNA level and carcass fat content was observed in Jinhua and Landrace gilts. These results are in accordance with previous reports. The increased *FASN* expression could lead to obesity through the accumulation of triglycerides in humans (Semenkovich, 1997). In pigs, there was a significant correlation between the *FASN* potential enzyme activity and its mRNA expression level, the decreased *FASN* expression reduced the capacity of adipose tissue to synthesize lipids and subsequently led to decreased fat accumulation (Huang et al., 2008). However, there was no obvious relationship between the *FASN* expression and intramuscular fat (IMF) content in the porcine longissimus muscle (Chen et al., 2004) and there was a negative correlation between the *FASN* mRNA level and IMF content in Kazak sheep (Qiao et al., 2007). It suggested that the *FASN* function as an enzyme of fat storage has a diverse influence in subcutaneous adipose tissue and intramuscular fat tissue. In this study, the *FASN* mRNA level showed an age-dependent pattern, which increased with age in the two breeds. These results are in accordance with previous studies which indicate that adipose tissue may have different activity of related enzymes at different growth periods (Shan et al., 2006). All Jinhua and Landrace gilts were studied in the period of growth in which adipose tissue may be in the process of differentiation, which could cause developmental patterns of *FASN* expression levels in adipose tissue. In addition, Jinhua gilts had higher *FASN* expression levels compared with Landrace gilts, suggesting the differential expression levels of *FASN* in different breeds of pigs.

LIPE is an adipocyte enzyme that cleaves fatty acids from intracellular triacylglycerol (Mersmann, 1998), and an increase in *LIPE* could depress the accumulation of triglycerol in adipose cells (Thompson et al., 1993; Sztalryd et al., 1995). In

the present study, there was no obvious relationship between *LIPE* expression and carcass fat content in the two breeds, but there was a significant positive relationship between carcass fat content and the ratio of *FASN* to *LIPE* mRNA. The results confirmed that fat accumulation in adipose tissue represents a balance between synthesis and degradation (Chilliard, 1993). There was no significant correlation between IMF content and *LIPE* mRNA expression, whereas the ratio of *FASN* to *LIPE* mRNA was found to be positively correlated with IMF content (Chen et al., 2004). However, there was a negative relationship between *LIPE* expression or the *FASN/LIPE* mRNA and IMF content in Kazak sheep (Qiao et al., 2007). This discrepancy may be caused by different characteristics of fat deposition in the species. In this study, Jinhua gilts had lower *LIPE* expression compared with Landrace gilts, which implies that in the course of fat deposition pigs may accumulate body fat through a depression of lipid utilization and therefore Jinhua gilts need lower *LIPE* expression due to a decrease in lipid utilization. In addition, Jinhua gilts had higher the *FASN/LIPE* mRNA compared with Landrace gilts. This may be explained by the model difference in carcass fat accumulation in different pig breeds.

REFERENCES

- Bauman D.E. (1976): Intermediary metabolism of adipose tissue. *Federation Proceedings*, 35, 2308–2313.
- Carr T.R., Walters L.E., Whiteman J.V. (1978): Carcass composition changes in growing and finishing swine. *Journal of Animal Science*, 47, 615–621.
- Chen J., Yang X.J., Dong H., Chen J., Zhao R.Q. (2004): Expression of FAS and HSL mRNA in *longissimus dorsi* muscle and their relation to intramuscular fat contents in pigs. *Journal of Agricultural Biotechnology*, 12, 422–426. (in Chinese)
- Chilliard Y. (1993): Dietary fat and adipose tissue metabolism in ruminants, pigs and rodents: a review. *Journal of Dairy Science*, 76, 3897–3931.
- Correa J.A., Faucitano L., Laforest J.P., Rivest J., Marcoux M., Gariépy C. (2006): Effects of slaughter weight on carcass composition and meat quality in pigs of two different growth rates. *Meat Science*, 72, 91–99.
- D'Souza D., Pethick D., Dunshea F., Suster D., Pluske J., Mullan B. (2004): The pattern of fat and lean muscle tissue deposition differs in the different pork primal cuts of female pigs during the finisher growth phase. *Livestock Production Science*, 91, 1–8.
- Friesen K.G., Nelssen J.L., Unruh J.A., Goodband R.D., Tokach M.D. (1994): Effects of the interrelationship between genotype, sex, and dietary lysine on growth performance and carcass composition in finishing pigs fed to either 104 or 127 kilograms. *Journal of Animal Science*, 72, 946–954.
- Haemmerle G., Zimmermann R., Zechner R. (2003): Letting lipids go: Hormone-sensitive lipase. *Current Opinion in Lipidology*, 14, 289–297.
- Huang Q.C., Xu Z.R., Han X.Y., Li W.F. (2008): Effect of dietary betaine supplementation on lipogenic enzyme activities and fatty acid synthase mRNA expression in finishing pigs. *Animal Feed Science and Technology*, 140, 365–375.
- Kazala E.C., Petrak J.L., Lozeman F.J., Mir P.S., Laroche A., Deng J., Weselake R.J. (2003): Hormone-sensitive lipase activity in relation to fat content of muscle in Wagyu hybrid cattle. *Livestock Production Science*, 79, 87–96.
- Kousteni S., Tura-Kockar F., Ramji D.P. (1999): Sequence and expression analysis of a novel Xenopus Laevis cDNA that encodes a protein similar to bacterial and chloroplast ribosomal protein L24. *Gene*, 235, 13–18.
- Lalitha S. (2000): Primer Primer 5.0. *Biotech and Internet Report*, 1, 270–272.
- Mersmann H.J. (1998): Lipoprotein and hormone-sensitive lipases in porcine adipose tissue. *Journal of Animal Science*, 76, 1396–1404.
- Mersmann H.J., Houk J.M., Phinney G., Underwood M.C. (1973): Effect of diet and weaning age on *in vitro* lipogenesis in young swine. *Journal of Nutrition*, 103, 821–828.
- Morales J., Baudet J.J., Pérez J.F., Mourot J., Gasa J. (2003): Body fat content, composition and distribution in Landrace and Iberian finishing pigs given ad libitum maize and acorn-sorghum-maize-based diets. *Animal Science*, 77, 215–224.
- O'Hea E.K., Leveille G.A. (1969): Significance of adipose tissue and liver as sites of fatty acid synthesis in the pig and efficiency of utilization of various substrates for lipogenesis. *Journal of Nutrition*, 99, 338–344.
- Qiao Y., Huang Z., Li Q., Liu Z., Hao C., Shi G., Dai R., Xie Z. (2007): Developmental changes of the FAS and HSL mRNA expression and their effects on the content of intramuscular fat in Kazak and Xinjiang sheep. *Journal of Genetics and Genomics*, 34, 909–917.
- Renaudeau D., Mourot I. (2007): A comparison of carcass and meat quality characteristics of Creole and Large White pigs slaughtered at 90 kg BW. *Meat Science*, 76, 165–171.
- Semenkovich C.F. (1997): Regulation of fatty acid synthase (*FAS*). *Progress in Lipid Research*, 36, 43–53.

- Shan T.Z., Wang Y.Z., Liu J.X., Xu Z.R., Feng J. (2006): Development gene expression of fatty acid synthase (*FAS*) in abdominal adipose of swine. *Acta Veterinaria et Zootechnica Sinica*, 37, 662–666. (in Chinese)
- Smith S., Witkowski A., Joshi A.K. (2003): Structural and functional organization of the animal fatty acid synthase. *Progress in Lipid Research*, 42, 289–317.
- Sztalryd C., Komaromy M.C., Kraemer F.B. (1995): Overexpression of hormone-sensitive lipase prevents triglyceride accumulation in adipocytes. *The Journal of Clinical Investigation*, 95, 2652–2661.
- Thompson M.P., Cooper S.T., Parry B.R., Tuckey J.A. (1993): Increased expression of the mRNA for hormone-sensitive lipase in adipose tissue of cancer patients. *Biochimica et Biophysica Acta*, 1180, 236–242.
- Wanger J.R., Schinckel A.P., Chen W., Forrest J.C., Coe B.L. (1999): Analysis of body composition changes of swine during growth and development. *Journal of Animal Science*, 77, 1442–1466.
- White B.R., Lan Y.H., McKeith F.K., Novakofski J., Wheeler M.B., McLaren D.G. (1995): Growth and body composition of Meishan and Yorkshire barrows and gilts. *Journal of Animal Science*, 73, 738–749.
- Xiong W.Z., Yang F., Zhou A.G. (2001): Study of regulation of exogenous recombinant somatotropin on fat metabolism in different cross-finishing pigs. *Chinese Journal of Animal and Veterinary Science*, 32, 1–4. (in Chinese)
- Xu Z.R. (1994): *Modern Nutrition of Swine*. Zhejiang University Press, Hangzhou, 57–72. (in Chinese)
- Yan X.C., Wang Y.Z., Xu Z.R. (2002): Regulation of fatty acid synthase (*FAS*) gene expression in animals. *Acta Zoonutritional Sinica*, 14, 1–4. (in Chinese)
- Yang H.L. (2005): Study on the biochemistry and histomorphology of intramuscular fat deposition in Laiwu black pigs. MA [Thesis]. Shandong Agricultural University, Shandong, China. (in Chinese)

Received: 2009–05–07

Accepted after corrections: 2010–04–23

Corresponding author

Dr. Zirong Xu, Key Laboratory for Molecular Animal Nutrition of Ministry of Education, Institute of Feed Science, Zhejiang University, 310029 Hangzhou, P.R. China
Tel. +86 373 3693 019, fax +86 373 3040 718, e-mail: miaozhiguo2000@yahoo.com.cn
