

<https://doi.org/10.17221/340/2018-PSE>

Effects of pre-preceding leguminous crops on yield and chemical composition of winter wheat grain

AGNIESZKA PSZCZÓŁKOWSKA*, ADAM OKORSKI, JACEK OLSZEWSKI,
GABRIEL FORDOŃSKI, SŁAWOMIR KRZEBIETKE, ALINA CHAREŃSKA

Faculty of Environmental Management and Agriculture, University of Warmia and Mazury in Olsztyn, Olsztyn, Poland

*Corresponding author: agnieszka.pszczolkowska@uwm.edu.pl

ABSTRACT

Pszczółkowska A., Okorski A., Olszewski J., Fordoński G., Krzebietke S., Chareńska A. (2018): Effects of pre-preceding leguminous crops on yield and chemical composition of winter wheat grain. *Plant Soil Environ.*, 64: 592–596.

The after-effects of pre-preceding crops (second year), i.e. legumes and spring wheat, and nitrogen fertilization rate (0, 60, 120 and 180 kg N/ha) on the yield and chemical composition of winter wheat grain were analysed in a field experiment conducted in 2013–2015. Winter wheat was characterized by higher yield when sown after blue lupine (increase of 0.23 t/ha) and faba beans with a determinate growth habit (increase of 0.37 t/ha) than after spring wheat. Grain yield increased significantly with a rise in nitrogen fertilization rate (by 2.03, 3.47 and 4.02 t/ha, respectively). The species of pre-preceding crops had no significant effect on the phosphorus, potassium, magnesium and calcium content of winter wheat grain. Winter wheat grown after faba beans with an indeterminate growth habit was most abundant in nitrogen. The applied nitrogen fertilizer rates did not modify the concentrations of phosphorus, magnesium and calcium in winter wheat grain. The nitrogen content of grain increased significantly with a rise in nitrogen fertilization rates. A significant increase in manganese and zinc levels was observed when spring wheat was the pre-preceding crop and the iron content of grain increased significantly when winter wheat was grown after peas and blue lupine.

Keywords: cereals; *Triticum aestivum* L.; nutrition; Fabaceae

Winter wheat yields are influenced by both environmental and agronomic factors (Sugár et al. 2016). Preceding crop and nitrogen fertilization are the key agronomic determinants of grain yield. Optimal growing conditions contribute to an increase in the technological efficiency of grain production.

The influence of preceding crops on winter wheat grain yield was widely researched (Lehocká and Klimeková 2009, Houšť et al. 2012, Švančárková and Žák 2015, Fordoński et al. 2016). The results of published studies indicate that legumes, perennial crops of the family Fabaceae and winter oilseed rape are the optimal preceding crops for winter wheat. Cereals are less suitable preceding crops because

they strongly degrade soil and increase the risk of pests and disease, in particular fungal infections.

Nitrogen fertilization significantly influences the yield and quality of winter wheat grain. Nitrogen is a mineral nutrient with the greatest yield-forming effect in plant production. Plants that are well supplied with nitrogen produce more tillers and grains but excessive rates of nitrogen fertilization can lead to lower yields (Kutman et al. 2011, Klikocka et al. 2016).

High nitrogen rates increase grain yield and the content of protein and gluten in grain, but they can also compromise grain quality. When applied in the late growth stages of wheat, nitrogen can

Supported by the Polish Ministry of Agriculture and Rural Development, Project No. HOR. 3.3.3/2011-2015.

increase the percentage of low-molecular-weight gliadin subunits in grain protein, which deteriorates the properties of gluten (Johansson et al. 2001, Litke et al. 2018).

The aim of the study was to assess the effects of pre-preceding leguminous crops and nitrogen fertilization rate on yield and chemical composition of winter wheat grain.

MATERIAL AND METHODS

A field experiment was conducted in the growing seasons of 2012/2013–2014/2015 in the Agricultural Experiment Station in Bałcyny (53°35'49.7"N, 19°51'17.3"E). The experiment had a randomized block design with two factors and 4 replications. The first experimental factor was the pre-preceding crop for winter wheat: semi-leafless peas cv. Tarchalska, blue lupine cv. Zeus (with an indeterminate growth habit), faba beans cv. Olga (with an indeterminate growth habit), faba beans cv. Granit (with a determinate growth habit) and spring wheat cv. Trappe. The second experimental factor was the rate of nitrogen fertilization: 0 (control); 60 kg N/ha (single application in early spring before the beginning of the growing season); 120 kg N/ha (two applications – 60 kg in early spring and 60 kg two weeks later); 180 kg N/ha (three applications – 60 kg in early spring, 60 kg two weeks later and 60 kg during heading). Nitrogen was applied as ammonium nitrate (34% N).

Winter oilseed rape was the preceding crop for winter wheat cv. Muszelka and spring wheat and selected legume species were the preceding crops for winter oilseed rape. The experimental plots had an area of 30 m² each. Each year, winter wheat cv. Muszelka was sown between 10 and 20 September.

The yield and the macronutrient and micronutrient content of winter wheat grain were determined at a certified laboratory of the District Chemical and Agricultural Station in Olsztyn, Poland. The phosphorus content of grain was determined by the vanadium-molybdenum method, calcium and potassium content by atomic emission spectrometry (AES), magnesium content by flame atomic absorption spectrometry (FAAS), and total nitrogen content by the hypochlorite method. The micronutrient content of grain was analysed by flame atomic absorption spectrometry (FAAS). Grain was mineralized in a mixture of nitric acid and hypochloric acid, and micronutrient concentra-

tions were determined by flame atomic absorption spectrometry (<http://oschr.olsztyn.pl/wp-content/uploads/2018/02/AB-277.pdf>).

The grain yields of winter wheat and the results of chemical analyses were verified statistically. The least significant difference was calculated at $\alpha \leq 0.05$ for the yield of winter wheat grain and at $\alpha \leq 0.01$ for macronutrient and micronutrient content using the Tukey's test. All calculations were performed in Statistica v. 10 software (Dell Inc. 2016, Tulsa, USA).

RESULTS AND DISCUSSION

The grain yield of winter wheat was high when the main crop was grown after legumes and spring wheat as pre-preceding crops (Table 1). In all treatments, the grain yield of winter wheat was highest in 2014 and lowest in 2013 (Table 1). Winter wheat produced significantly higher yields after blue lupine and the determinate cultivar of faba beans (increase of 0.23 t/ha and 0.37 t/ha, respectively) than after spring wheat. The grain yields in treatments where winter wheat was grown after peas and the indeterminate cultivar of faba beans did not differ significantly from those noted in treatments where spring wheat was the pre-preceding crop.

The combined effects of nitrogen fertilization and pre-preceding crops on winter wheat yield were visible only in treatments with fertilizer rates of 0 and 60 kg N/ha (Table 1). In the control treatments (0 kg N/ha), grain yield was highest when winter wheat was grown after determinate faba beans (6.72 t/ha) and lowest after indeterminate faba beans (6.21 t/ha) and spring wheat (6.34 t/ha). In treatments fertilized with 60 kg N/ha, the lowest yield was observed after spring wheat, and the highest yield was noted after determinate faba beans. In the remaining fertilization treatments, pre-preceding crops did not exert a significant influence on winter wheat yields. An analysis of the effect of nitrogen fertilization across years revealed that the grain yield of winter wheat was high in the control treatments, but it was significantly lowest in comparison with the experimental treatments. Grain yield increased significantly with a rise in the rate of nitrogen fertilization except in 2014 at nitrogen rates of 60 and 120 kg N/ha. Nitrogen fertilizer applied in treatments with pre-preceding crops at 180 kg N/ha produced the highest grain yields (Table 1). In the experimental treat-

<https://doi.org/10.17221/340/2018-PSE>

Table 1. Effects of nitrogen (N) fertilization rates and pre-preceding crops on the yield of winter wheat grain harvested in 2013–2015 (t/ha) (preceding crop – winter oilseed rape)

Pre-preceding crop	N rate (kg/ha)					Mean	N rate (kg/ha)				
	0	60	120	180	0		60	120	180	Mean	
	2013						2015				
Pea	5.24 ^e	7.46 ^c	8.74 ^{ab}	8.85 ^a	7.57 ^b	5.94 ^d	7.42 ^c	9.60 ^{ab}	10.37 ^a	8.33 ^a	
Blue lupine	5.23 ^e	7.43 ^c	8.71 ^{ab}	9.07 ^a	7.61 ^b	5.53 ^d	7.60 ^c	9.49 ^{ab}	10.12 ^a	8.19 ^a	
Determinate cultivar of faba bean	5.90 ^d	7.78 ^c	8.67 ^b	9.28 ^a	7.91 ^a	5.68 ^d	7.90 ^c	8.63 ^b	10.03 ^a	8.06 ^a	
Indeterminate cultivar of faba bean	5.19 ^e	7.48 ^c	8.96 ^a	8.86 ^a	7.62 ^b	5.13 ^{de}	7.63 ^c	9.07 ^{ab}	9.93 ^a	7.94 ^a	
Spring wheat	5.29 ^e	7.45 ^c	8.67 ^b	8.81 ^a	7.56 ^b	5.61 ^d	7.53 ^c	9.35 ^{ab}	9.98 ^a	8.12 ^a	
Mean	5.37 ^b	7.52 ^b	8.75 ^a	8.97 ^a	–	5.58 ^d	7.62 ^c	9.23 ^b	10.09 ^a	–	
	2014						2013–2015				
Pea	7.93 ^l	11.93 ^b	12.38 ^a	10.11 ^g	10.59 ^c	6.37 ^h	8.33 ^e	10.09 ^{bc}	10.53 ^a	8.83 ^{bc}	
Blue lupine	9.10 ⁱ	11.61 ^c	12.17 ^{ab}	10.56 ^f	10.86 ^b	6.62 ^{gh}	8.53 ^e	9.94 ^{bc}	10.45 ^a	8.88 ^{ab}	
Determinate cultivar of faba bean	8.58 ^{jk}	12.20 ^{ab}	12.65 ^a	11.00 ^e	11.11 ^a	6.72 ^g	8.89 ^d	9.83 ^{bc}	10.65 ^a	9.02 ^a	
Indeterminate cultivar of faba bean	8.30 ^k	11.83 ^{bc}	12.44 ^a	10.23 ^{fg}	10.70 ^{bc}	6.21 ^h	8.45 ^e	9.95 ^{bc}	10.41 ^{ab}	8.75 ^{bc}	
Spring wheat	8.12 ^{kl}	11.35 ^{cd}	12.08 ^{ab}	9.57 ^h	10.28 ^d	6.34 ^h	8.18 ^{ef}	9.79 ^c	10.29 ^{ab}	8.65 ^c	
Mean	8.41 ^d	11.78 ^b	12.34 ^a	10.29 ^c	–	6.45 ^d	8.48 ^c	9.92 ^b	10.47 ^a	–	

^{a,b,c}Means with the same letter are not significantly different at $\alpha \leq 0.05$

ments, the grain yield of winter wheat increased by 4.02 t/ha on average, relative to control.

According to many authors (Lehocká and Klimeková 2009, Houšť et al. 2012, Švančárková and Žák 2015), preceding crops significantly influence winter wheat yields. Kozak and Kotecki (2006) analysed the effect of peas on the development and yield of winter wheat cv. Kobra. Winter wheat plants grown in a treatment where the post-harvest residues of peas (roots, stubble and straw) were incorporated into the soil by ploughing were significantly taller and were characterized by significantly longer flag leaves, higher 1000 kernel weight, highest grain yield, highest straw biomass and highest total protein content, relative to treatments without pea residues. In a study by Švančárková and Žák (2015), winter wheat yields were higher after peas (5.9 t/ha) than after red clover (5.4 t/ha). Peas not only significantly increased yield, but also improved grain quality parameters. In an experiment by Lehocká and Klimeková (2009), winter wheat yields reached 6.5 t/ha after alfalfa and were 0.6 t/ha lower when spring barley was the preceding crop. The lowest yields were observed after peas (5.6 t/ha). The grain of winter wheat grown after spring barley was characterized by the highest processing suitability and optimal technological properties. Houšť et al. (2012) did not report significant differences in winter

wheat yields in three crop rotations (alfalfa, peas, maize grown for silage) with different percentages of cereals (33.3, 50 and 66.6%). In a study by Prusiński et al. (2016), the average yield and protein content of winter triticale grain were higher after leguminous forecrops than after spring barley.

Recent research indicates that the grain yield of selected wheat cultivars continues to increase up to a nitrogen rate of 200 kg N/ha. Similar results were reported by Campillo et al. (2010) who found that a fertilizer rate of 200 kg N/ha produced the highest profits in the production of winter wheat. In the work of Tayebbeh et al. (2011), winter wheat responded most favourably to a nitrogen rate of 240 kg N/ha. A fertilizer rate of 360 kg N/ha led to a further increase in grain yield, but the difference between treatments fertilized with 360 and 240 kg N/ha was not statistically significant. In the experiments conducted by Sugár et al. (2016), wheat yields peaked in response to 80 and 160 kg N/ha, subject to cultivar. Prusiński et al. (2016) observed that mineral nitrogen fertilizers applied at a rate of 60 to 180 kg N/ha did not induce significant differences in the average grain yield of winter triticale cv. Tulus. In a study by Dogan and Bilgili (2010), the optimal nitrogen rate for winter wheat was 150 kg N/ha, and an increase to 200 kg N/ha did not lead to a significant increase in grain

<https://doi.org/10.17221/340/2018-PSE>

Table 2. Effects of pre-preceding crops on the macronutrient (g/kg dry matter (DM)) and micronutrient (mg/kg DM) content of winter wheat grain (preceding crop – winter oilseed rape)

Pre-preceding crop	Macronutrient					Micronutrient			
	N	P	K	Mg	Ca	Mn	Zn	Cu	Fe
Pea	12.90 ^D	2.65 ^A	4.82 ^A	0.95 ^A	0.40 ^A	17.55 ^B	18.47 ^{AB}	2.10 ^B	39.00 ^A
Blue lupine	13.50 ^B	2.62 ^A	4.72 ^A	0.90 ^A	0.33 ^A	17.97 ^B	17.90 ^B	2.37 ^A	38.20 ^A
Determinate cultivar of faba bean	13.20 ^C	2.67 ^A	4.80 ^A	0.95 ^A	0.33 ^A	17.55 ^B	16.95 ^C	2.27 ^{AB}	35.80 ^B
Indeterminate cultivar of faba bean	14.20 ^A	2.63 ^A	4.65 ^A	0.85 ^A	0.30 ^A	17.00 ^B	17.90 ^B	2.30 ^{AB}	33.80 ^C
Spring wheat	13.50 ^B	2.77 ^A	4.87 ^A	0.97 ^A	0.35 ^A	20.22 ^A	19.10 ^A	2.07 ^B	33.20 ^C

^{A,B,C}Means with the same letter are not significantly different at $\alpha \leq 0.01$

yield. Litke et al. (2018) found that the grain yield of winter wheat increased significantly with a rise in nitrogen rate. In cv. Skagen, grain yield increased significantly up to nitrogen rate of 180 kg N/ha. The values of grain quality parameters also increased considerably when nitrogen fertilizer was applied at 150 to 210 kg N/ha.

Podolska et al. (2005) analysed the profitability of winter wheat production and concluded that nitrogen fertilization rates higher than 120 kg N/ha were not economically justified because they decreased the productivity of land, labour and capital. The fertilizer rate of 120 kg N/kg delivered optimal yields, grain quality and profitability.

The evaluated experimental factors induced minor variations in the macronutrient content of winter wheat grain (Tables 2 and 3). The analysed pre-preceding crops did not significantly differentiate the concentrations of phosphorus, potassium, magnesium and calcium in grain (Table 2). The highest nitrogen content of grain (14.2 g N/kg DM (dry matter)) was observed in the treatment where the indeterminate cultivar of faba beans was the pre-preceding crop. The grain of winter wheat grown after peas had the lowest nitrogen content

(12.9 g N/kg DM). The observed differences were similar in the remaining treatments.

The nitrogen content of winter wheat grain increased significantly with a rise in the rate of nitrogen fertilization (Table 3). The tested fertilizer rates did not contribute to significant differences in the phosphorus, magnesium and calcium content of grain (Table 3). In contrast, Wojtkowiak et al. (2015) demonstrated that a rise in nitrogen rate from 80 to 120 kg N/ha increased the phosphorus content of spring triticale grain cv. Milewo. Potassium levels were lowest in treatments fertilized with 180 kg N/ha. The grain of winter wheat fertilized with 60 kg N/ha was characterized by the highest potassium content (increase of 0.28 g K/kg DM) (Table 3).

The manganese and zinc content of grain increased significantly when winter wheat was grown after spring wheat. The highest iron concentration in grain was observed when peas and blue lupine were the pre-preceding crops (Table 2). The grain of winter wheat grown after spring wheat and peas was characterized by the lowest copper content, and the grain of winter wheat grown after blue lupine by the highest copper content.

Table 3. Effects of nitrogen (N) fertilization rates on the macronutrient (g/kg dry matter (DM)) and micronutrient (mg/kg DM) content of winter wheat grain

N rate (kg/ha)	Macronutrient					Micronutrient			
	N	P	K	Mg	Ca	Mn	Zn	Cu	Fe
0	13.10 ^E	2.68 ^A	4.78 ^C	0.90 ^A	0.34 ^A	17.42 ^A	18.40 ^A	2.36 ^A	35.01 ^C
60	13.20 ^C	2.70 ^A	4.92 ^A	0.96 ^A	0.44 ^A	18.04 ^A	17.98 ^A	2.26 ^A	31.93 ^D
120	13.30 ^B	2.70 ^A	4.84 ^B	0.88 ^A	0.30 ^A	18.22 ^A	17.84 ^A	2.16 ^B	39.90 ^A
180	14.30 ^A	2.62 ^A	4.64 ^D	0.90 ^A	0.36 ^A	18.56 ^A	18.04 ^A	2.12 ^B	37.30 ^B

^{A,B,C}Means with the same letter are not significantly different at $\alpha \leq 0.01$

<https://doi.org/10.17221/340/2018-PSE>

Manganese and zinc concentrations in winter wheat grain did not differ significantly in treatments with various rates of nitrogen fertilization (Table 3). Copper content was highest in treatments without nitrogen fertilization, whereas iron content peaked in response to 120 kg N/ha. The macronutrient and micronutrient content of grain was within the reference range for winter wheat grown in the Polish climate (Nogalska et al. 2012).

In conclusion, this study demonstrated that pre-preceding crops had a significant effect on the grain yield of winter wheat. Grain yields increased when winter wheat was grown after blue lupine and determinate faba beans. An increase in the rate of nitrogen fertilization significantly increased winter wheat yields. When applied at the rate of 180 kg N/ha in treatments with pre-preceding crops, nitrogen fertilizer increased wheat yields to 10.47 t/ha on average. The nitrogen content of grain increased with a rise in the rate of nitrogen fertilizer. The grain of winter wheat grown after indeterminate faba beans was characterized by the highest nitrogen content. The evaluated pre-preceding crops did not induce significant variations in the phosphorus, potassium, magnesium and calcium content of winter wheat grain. The manganese and zinc content of grain increased significantly when winter wheat was grown after spring wheat, and the iron content of grain peaked when peas and blue lupine were the pre-preceding crops. Winter wheat grain was the least abundant in copper when grown after spring wheat and peas. The applied rates of nitrogen fertilization did not contribute to significant differences in phosphorus, magnesium, calcium, manganese and zinc concentrations in grain. Winter wheat grain was characterized by the highest copper content in treatments without nitrogen fertilization, and iron concentrations peaked in response to the nitrogen fertilizer rate of 120 kg N/ha.

REFERENCES

- Campillo R., Jobet C., Undurraga P. (2010): Effects of nitrogen on productivity, grain quality, and optimal nitrogen rates in winter wheat cv. Kumpa-INIA in Andisols of southern Chile. *Chilean Journal of Agricultural Research*, 70: 122–131.
- Dogan R., Bilgili U. (2010): Effects of previous crop and N-fertilization on seed yield of winter wheat (*Triticum aestivum* L.) under rain-fed Mediterranean conditions. *Bulgarian Journal of Agricultural Science*, 16: 733–739.
- Fordoński G., Pszczółkowska A., Okorski A., Olszewski J., Załuski D., Gorzkowska A. (2016): The yield and chemical composition of winter oilseed rape seeds depending on different nitrogen fertilization rates and the preceding crop. *Journal of Elementology*, 21: 1225–1234.
- Houšť M., Procházková B., Hledík P. (2012): Effect of different tillage intensity on yields and yield-forming factors in winter wheat. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis*, 60: 89–96.
- Johansson E., Prieto-Linde M.L., Jonsson J.Ö. (2001): Effects of wheat cultivar and nitrogen application on storage protein composition and breadmaking quality. *Cereal Chemistry*, 78: 19–25.
- Klikocka H., Cybulska M., Barczak B., Narolski B., Szostak B., Kobiałka A., Nowak A., Wójcik E. (2016): The effect of sulphur and nitrogen fertilization on grain yield and technological quality of spring wheat. *Plant, Soil and Environment*, 62: 230–236.
- Kozak M., Kotecki A. (2006): The residual effect of common pea cultivars on growth and yielding of winter wheat. Part III Growth and yielding of winter wheat. *Zeszyty Naukowe Uniwersytetu Przyrodniczego we Wrocławiu*, 546: 159–174. (In Polish)
- Kutman U.B., Yildiz B., Cakmak I. (2011): Effect of nitrogen on uptake, remobilization and partitioning of zinc and iron throughout the development of durum wheat. *Plant and Soil*, 342: 149–164.
- Lehocká Z., Klimeková M. (2009): Yields and selected quality parameters of winter wheat (*Triticum aestivum* L.) in organic cropping system as affected by forecrops in the years 2006–2008. *Agricultura – Stiinta si Practica*, 69–701: 13–18.
- Litke L., Gaile Z., Ruža A. (2018): Effect of nitrogen fertilization of winter wheat yield and yield quality. *Agronomy Research*, 16: 500–509.
- Nogalska A., Sienkiewicz S., Czaplá J., Skwierawska M. (2012): The effect of multi-component fertilizers on the yield and mineral composition of winter wheat and macronutrient uptake. *Journal of Elementology*, 17: 629–638.
- Podolska G., Krasowicz S., Sułek A. (2005): Economic and quality evaluation of winter wheat cultivation in relation to different nitrogen fertilization levels. *Pamiętnik Puławski*, 139: 175–188. (In Polish)
- Prusiński J., Borowska M., Kaszkowiak E., Olszak G. (2016): The after-effect of chosen Fabaceae forecrops on the yield of grain and protein in winter triticale (*Triticosecale* sp. Wittmack ex A. Camus 1927) fertilized with mineral nitrogen. *Plant, Soil and Environment*, 62: 571–576.
- Sugár E., Berzsenyi Z., Árendás T., Bónis P. (2016): Effect of nitrogen fertilization and genotype on the yield and yield components of winter wheat. *Die Bodenkultur: Journal of Land Management, Food and Environment*, 67: 25–34.
- Švančárková M., Žák Š. (2015): The grain quality of winter wheat in organic and conventional farming. *Acta Fytotechnica et Zootechnica*, 18 (special issue): 22–24.
- Tayebeh A., Alemzadeh A., Kazemeini S.A. (2011): Wheat yield and grain protein response to nitrogen amount and timing. *Australian Journal of Crop Science*, 5: 330–336.
- Wojtkowiak K., Stępień A., Warechowska M., Markowska A. (2015): Effect of nitrogen fertilization method on the yield and quality of Milewo variety spring triticale grain. *Polish Journal of Natural Science*, 30: 173–184.

Received on May 21, 2018

Accepted on August 23, 2018

Published online on November 9, 2018