

## The influence of land use practices on earthworm communities in saline agriculture soils of the west coast region of China's Bohai Bay

Y. Tao<sup>1,3</sup>, W. Gu<sup>1</sup>, J. Chen<sup>1,2</sup>, J. Tao<sup>1</sup>, Y.J. Xu<sup>1</sup>, H. Zhang<sup>1</sup>

<sup>1</sup>State Key Laboratory of Earth Surface Process and Resource Ecology,  
Beijing Normal University, Beijing, P.R. China

<sup>2</sup>Academy of Disaster Reduction and Emergency Management, Ministry of Civil Affairs  
and Ministry of Education, Beijing Normal University, Beijing, P.R. China

<sup>3</sup>College of Urban and Environmental Science, Northeast Normal University,  
Changchun, P.R. China

### ABSTRACT

The effects of land use practices on soil fauna, especially earthworms, are poorly known in coastal saline agricultural soils. Here we compare earthworm communities in six types of land use practice in the coastal region of China's Bohai Bay, namely uncultivated saline soil, two orchard (pear and winter jujube) lands, man-made forests (chinese ash), vegetable land and cropped land (maize). In addition, we recorded selected physicochemical properties of the soil. Soil organic matter content and total N were significantly higher under pear orchard and vegetable land than under the other land use practices, and their lowest values were observed from uncultivated saline soil. Vegetable land and pear orchard land showed a significantly higher abundance of earthworms than the other land use practices, whereas no earthworm was found in uncultivated saline soil. The sites under individual practices supported one to three earthworm species. *Aporrectodea trapezoides* species was present under four types of land use practice, and the biomass of this species accounted for more than 60% of the community. Vegetable land and pear orchard land supported richer earthworm community than the other land use practices, dominated by *Aporrectodea trapezoides* and *Drawida japonica*. These preliminary results indicated that land use practices have substantial effects on the abundance and composition of earthworm communities in saline soils.

**Keywords:** soil fauna; community structure; biological environment; salt-affected agricultural soils

The conversion of natural tropical ecosystems to agricultural ecosystems may accelerate the changes of the soil. This implies not only a change in soil chemical properties, but also a change in biodiversity (Hairiah et al. 2001, Wu et al. 2001). Adequate management of land use may favor environmental conditions required by certain groups of soil flora and fauna (Geissen et al. 2009). The studies of Emmerling (1995) showed that land use

and soil organic matter content were the main factors affecting the distribution of soil macrofauna. Dlamini and Haynes (2004) also found agricultural land use impacted on the size, composition and diversity of earthworm communities in agricultural soils. In return, management of earthworm populations is becoming more important for sustaining soil productivity and fertility in agro-ecosystems (Whalen et al. 1998).

---

Supported by the National Natural Science Foundation of China, Project No. 41201237, and by the Free Exploring Foundation from State Key Laboratory of Earth Surface Processes and Resource Ecology, Project No. 2012-ZY-04, and by the National '863', Project No. 2011AA100505. Tao Yan was funded by the Open Foundation from State Key Laboratory of Earth Surface Processes and Resource Ecology, Beijing Normal University, Project No. 2011-KF-13.

However, the research on the effects of land use practices (forest, orchard, cropped land, etc.) on soil faunal communities is still lacking in saline agricultural soils, where land use practices may substantially affect physical, chemical and biological properties of the soils (Qiao et al. 2001). Changes in environmental characteristics of soil could affect the community of soil fauna in soils, especially earthworms (Ammer et al. 2006). Earthworms are involved in key soil processes such as soil organic matter decomposition, soil nutrient turnover, aggregate formation and soil pore water dynamics through their burrowing activities (Edwards et al. 1990, Blair et al. 1995, Wessells et al. 1997, Brown and Doube 2004) and act as so called 'ecosystem engineers' (Lavell et al. 1997). Moreover, the relationships between earthworm abundance and soil properties could be used as indicators of the performance of agricultural systems in terms of soil quality (Clapperton et al. 2004).

Two main research questions were examined in the present study: (1) did the abundance of earthworm communities vary among land use practices in saline agricultural soils of the west coastal region of China's Bohai Bay; (2) What land use practice optimizes the use of soil in driving the improvement of ecosystem environments in a region of saline agricultural soils?

## MATERIAL AND METHODS

**Sites and soils.** The study area was located within the Zhongjie friendship farm (117°37'E, 38°25'N) of Hebei province in west coast of China's Bohai Bay. The area belongs to semi-humid continental monsoon climate zone with average annual temperature of 12.3°C. Average annual precipitation is 590.6 mm, of which most occurs from late June to September. However, precipitation is highly variable. Average mean annual evaporation is 2117.4 mm. Most sites have gone through several land uses and known history of fields was: > 8 years pear orchard (*Pyrus ussuriensis* Maxim), > 10 years winter jujube orchard (*Zizyphus jujube* Mill. cv. Dongzao), > 6 years chinese ash forest (*Fraxinus chinensis* Roxb), > 11 years vegetable land (*Brassica rapa* L. and *Allium fistulosum* L.), > 15 years cropped land (*Zea mays* L.). The uncultivated saline soil was used as a control. In cropped land, conventional farming was practiced by monoculture of maize (to a lesser extent of wheat), using mechanical

tillage, removal of crop straw and application of inorganic fertilizer. In winter jujube land, fertilizer, pesticides and herbicides were applied each year, whereas in pear land maize straw was covered onto the soil surface (besides fertilizer application). In Chinese ash forest, weeding and pest management operation were timely performed in those first few years. In vegetable land, cabbage seeds and scallion were cultivated each year. Large amount of manure were applied as basal fertilizer for vegetative soil. The soils in the area are classified as salinization tidal soils, composed of clay (27.1%), silt (30.1%) and sand (42.8%).

**Earthworm sampling and identification.** A total of 18 fields (three replicates of each type of land practice) were sampled in October, 2010. Within each field area (approximately 0.2 ha), three plots (about 15 m<sup>2</sup> in area) were randomly chosen. At each plot, a 50 cm × 50 cm area was excavated to the depth of 30 cm and the soil block was then hand-sorted to collect earthworms. Additionally, formalin was added to the bottom of the sampling holes (Raw 1959) but no additional deep burrowing earthworms were collected. The earthworms collected from each sampling point were counted in the field, placed in vials containing 70% ethanol and transferred to the laboratory for identification. All worms were washed clean of adhering soil, patted dry with a paper towel and weighed for total fresh biomass. Earthworm species were identified using the keys by Yin (1992).

**Soil sampling and analysis.** Adjacent to each earthworm sampling point, three soil cores of area of 2.5 cm<sup>2</sup> to depth of 30 cm were taken randomly using a steel core and pooled together as one composite sample. Soil samples were air-dried and ground. The soil analysis followed the 'Analysis of soil characteristics' (Lu 2000). Bulk density (BD) of each soil core was calculated as the total mass of oven dried soil divided by the core volume. Organic matter content was analyzed on ground soil by the Walkley and Black dichromate oxidation method (Blakemore et al. 1972). The total N content was determined using the Kjeldahl method. Soil pH was measured in a 1:5 soil:water (distilled water) slurry using a glass electrode. Available P was extracted with 0.5 mol/L NaHCO<sub>3</sub> by the Olsen method. Available K was extracted with 1 mol/L NH<sub>4</sub>OAc and was determined in all pot soil samples. Soil electrical conductivity was extracted using a 1:5 soil:water suspension and were determined with conductivity meter.

**Statistical analysis.** An analysis of variance (ANOVA) was performed to test for significant differences between the study sites, when the normality assumption (Kolmogorov-Smirnov test) was met. Non-normal distribution of the data was first log-transformed and then was tested with an analysis of variance (ANOVA). The standard error of the mean for values for each land practice was calculated using the three replicated sampling fields, three points in each. All differences were assessed using one-way ANOVA. Differences between means were tested with the *LSD* test ( $P < 0.05$ ). The relationships between the number and biomass of earthworms and specific soil parameters were expressed by linear correlation coefficient. All statistical analyses was done using the software package SPSS 16.0 (SPSS Inc., Chicago, USA).

## RESULTS AND DISCUSSION

The measured physical/chemical properties of the soils are presented in Table 1. Both organic matter content and total N were significantly higher under pear orchard and vegetable land than the other land use practices, and their lowest values were observed on uncultivated saline soil. In vegetable land, chinese cabbage residues was return into soil annually and inputs of manure. Pear orchards are grassed-down so organic matter inputs occur mainly through turnover of grass root material and irregular inputs of maize straw.

Qiao et al. (2001) found that vegetable land use practice showed a higher organic matter content than other land use practices, which was closely related to the inputs of fertilizer.

Land management practices applied to the saline soil are likely to have some positive effects on habitation of soil fauna (Emmerling and Paulsch 2001). The present study showed that the abundance of earthworms under vegetable land ( $> 340$  individuals/m<sup>2</sup>) and pear orchard land use practices ( $> 120$  individuals/m<sup>2</sup>) was significantly higher than those under the other land use practices (Figure 1), which may be due to the inputs of organic residues. Lowe and Butt (2002) found that organic matter management is important in the development of sustainable earthworm populations and their role in soil amelioration at restored sites. According to Sumner (1995), soils are generally classified as saline when they have an EC of 4 dS/m or more. Vegetable lands had a higher soil salinity concentrations considered to be 'unsafe' for many plant (Schoeneberger et al. 2002), but the soil salinity concentration had no effect on the population of earthworms. We speculated that higher organic matter content under vegetable land, which provided substrates or food resources for the soil organisms, probably reduces the negative effects of soil salinity on soil organisms (Pathak and Rao 1998, Wichern et al. 2006). The previous study of Tao et al. (2012) suggested that maize residue application alleviated the negative effects of soil salinity on the growth and

Table 1. Mean values ( $\pm$  standard error,  $n = 3$ ) of soil physicochemical parameters in different types of land use

	Organic matter content (g/kg)	Total N	Available K (mg/kg)	Available P (mg/kg)	Soil electrical conductivity (dS/m)	pH	Bulk density (g/cm <sup>3</sup> )	Soil water content (%)
Uncultivated saline soil	10.75 $\pm 0.12^c$	0.47 $\pm 0.02^c$	195.00 $\pm 7.64^c$	12.51 $\pm 0.54^b$	20.64 $\pm 3.01^a$	7.65 $\pm 0.05^a$	1.35 $\pm 0.01^a$	25.55 $\pm 0.50^b$
Pear orchard	13.18 $\pm 0.47^b$	0.94 $\pm 0.05^a$	336.67 $\pm 54.87^b$	13.46 $\pm 0.97^b$	1.86 $\pm 0.29^c$	7.68 $\pm 0.04^a$	1.16 $\pm 0.03^c$	26.31 $\pm 1.34^{ab}$
Winter jujube	10.98 $\pm 0.03^c$	0.55 $\pm 0.01^{bc}$	226.67 $\pm 8.82^{bc}$	13.99 $\pm 0.57^b$	1.32 $\pm 0.25^c$	7.81 $\pm 0.03^a$	1.28 $\pm 0.02^{ab}$	22.54 $\pm 1.25^c$
Chinese ash forest	10.70 $\pm 0.14^c$	0.48 $\pm 0.44^c$	228.33 $\pm 24.55^{bc}$	13.50 $\pm 0.61^b$	1.48 $\pm 0.50^c$	7.73 $\pm 0.06^a$	1.37 $\pm 0.01^a$	16.06 $\pm 0.50^d$
Vegetable land	16.61 $\pm 0.52^a$	0.92 $\pm 0.49^a$	600 $\pm 70.24^a$	18.85 $\pm 0.78^a$	4.49 $\pm 1.17^b$	7.84 $\pm 0.09^a$	1.20 $\pm 0.05^{bc}$	28.86 $\pm 0.87^a$
Cropped field	11.05 $\pm 0.14^c$	0.63 $\pm 0.47^b$	203.33 $\pm 17.63^c$	14.14 $\pm 0.81^b$	2.43 $\pm 0.37^c$	7.78 $\pm 0.06^a$	1.20 $\pm 0.06^{bc}$	22.06 $\pm 0.57^c$

Different letters indicate significant ( $P < 0.05$ ) differences for each parameter among different land uses (by *LSD* test)




Figure 1. Mean earthworm numbers and biomass ( $n = 3$ ) under different land uses. All numbers and biomass differences are respectively compared at  $P < 0.05$  for total earthworm (\*) or species (+); ne – no earthworms

reproduction of *A. trapezoides*. Additionally, under vegetable land and pear orchard land the farmers brought in organic manures or material from outside each year, which is the possible source of earthworms. The high abundance of earthworms under vegetable land and pear orchards land, in turn, would improve soil quality and enhance soil nutrients (Baker 2007, Eriksen-Hamel and Whalen 2008). However, under uncultivated saline soil, the higher soil salinity concentration or lower organic matter maybe limited production or invasion of earthworms from other lands.

Both cropped land and winter jujube land showed a low abundance of earthworms (Figure 1), which may be due to the types of land use practices. In cropped land, besides ordinary tillage, maize straw was removed from the field and burned annually. In winter jujube land, because of herbicide application soil surface is essentially bare apart from a thin layer of winter jujube leaves. These soil management practices adversely affected the population of soil fauna. But inorganic fertilizer application in both land uses, in order to increase their yields, could improve root growth of crop to keep a certain number of earthworms (Leroy et al. 2008). In two orchard land, pear land showed

a significantly higher abundance of earthworms than winter jujube land, maybe because pear land is covered with organic residues and grass. In Chinese ash forest, there was a layer of ash leaves and inorganic fertilizer was not added into the soil. Earthworms are less abundant in Chinese ash forest ( $< 50$  individuals/m<sup>2</sup>) (Figure 1), which could be due to lack of inorganic fertilizer. Inorganic fertilizer can promote decomposition of organic litter and growth of plant that affect soil microorganisms and fauna in low-level nutrient soils (Edwards et al. 1995). The uncultivated saline soil floor is essentially bare. As a result, organic matter content and total N were lower under uncultivated saline soil than the other land use practices. Earthworms do not occur under uncultivated land. Earthworm numbers and biomass were positively correlated with soil organic matter content and total N, in contrast with soil bulk density which showed a negative correlation (Table 2). These results were consistent with the general view (Reeleder et al. 2006, Huerta et al. 2007, Ouellet et al. 2008).

In the present study, different land use practices were supported. Saline agricultural soils except for uncultivated land (Figure 1). Improved vegetable land and pear orchard land had a larger

Table 2. Linear correlation coefficients between measures of selected soil properties and earthworm total numbers and biomass

Measures of earthworms	Organic matter content	Total N	Available K	Available P	Soil electrical conductivity	pH	Bulk density	Soil water content
Total numbers	0.938**	0.943**	0.771	0.657	0.029	0.600	−0.841*	0.600
Total biomass	0.886*	0.847*	0.829*	0.600	−0.143	0.657	−0.754	0.657

\* $P < 0.05$ ; \*\* $P < 0.01$

earthworm community and was dominated by endogeic *Aporrectodea trapezoides* living in a mineral soil layer and feeding on mineral particles of soil (Bonkowski and Schaefer 1997). The second predominant species was *Drawida japonica*. The number of *Drawida japonica* was higher under pear orchard than winter jujube, which may be due to roots of weed under pear orchard. This species may be likely to live around roots of plant. *Eisenia fetida* is an epigeic species and used as composting worm, preferring to live near the surface of the soil. This earthworm specie was present only under vegetable land, which was due to large amount of manure application for growing vegetables. In summary, different land use practices affected earthworm community characteristics in saline soil of the west coast region of China's Bohai Bay. Both vegetable land and pear orchard land use practices maybe optimizers the use of soil in driving the improvement of ecosystem environments in saline soils of the west coast region of China's Bohai Bay.

## Acknowledgement

I thank Dr. Zhang Yufeng for assistance with the identification of the earthworm species.

## REFERENCES

- Ammer S., Weber K., Abs C., Ammer C., Prietzel J. (2006): Factors influencing the distribution and abundance of earthworm communities in pure and converted Scots pine stands. *Applied Soil Ecology*, 33: 10–21.
- Baker G. (2007): Differences in nitrogen release from surface and incorporated plant residues by two endogeic species of earthworms (Lumbricidae) in a red-brown earth soil in southern Australia. *European Journal of Soil Biology*, 43: S165–S170.
- Blair J.M., Parmelee R.W., Lavelle P. (1995): Influences of earthworms on biogeochemistry. In: Hendrix P.F. (ed.): *Earthworm Ecology and Biogeography in North America*. CRC Press, Boca Raton, 127–158.
- Blakemore L.C., Searle P.L., Daly B.K. (1972): *Methods for Chemical Analysis of Soils*. New Zealand Soil Bureau Report 10A. Government Printer, Wellington.
- Bonkowski M., Schaefer M. (1997): Interactions between earthworms and soil protozoa: A trophic component in the soil food web. *Soil Biology and Biochemistry*, 29: 499–502.
- Brown G.G., Doube B.M. (2004): The functional interactions between earthworms, microorganisms, organic matter, and plant. In: Edwards C.A. (ed.): *Earthworm Ecology*. 2<sup>nd</sup> Edition. CRC Press, Boca Raton, 213–240.
- Clapperton J., Chan K.Y., Larney F.J. (2004): Managing the soil habitat for enhanced biological fertility. In: Abbott L., Murphy D. (eds.): *Soil Biological Fertility – A Key to Sustainable Land Use in Agriculture*. Kluwer Academic Publishers, Dordrecht, 203–224.
- Dlamini T.C., Haynes R.J. (2004): Influence of agricultural land use on the size and composition of earthworm communities in northern KwaZulu-Natal, South Africa. *Applied Soil Ecology*, 27: 77–88.
- Edwards C.A., Bohlen P.J., Linden D.R., Subler S. (1995): Earthworms in agroecosystems. In: Hendrix P.F. (ed.): *Earthworm Ecology and Biogeography in North America*. Lewis, Boca Raton, 185–213.
- Edwards W.M., Shipitalo M.J., Owens L.B., Norton L.D. (1990): Effect of *Lumbricus terrestris* L. burrows on hydrology of continuous no-till corn fields. *Geoderma*, 46: 73–84.
- Emmerling C. (1995): Long-term effects of inundation dynamics and agricultural land-use on the distribution of soil macrofauna in fluvisols. *Biology and Fertility of Soils*, 20: 130–136.
- Emmerling C., Paulsch D. (2001): Improvement of earthworm (Lumbricidae) community and activity in mine soils from open-cast coal mining by the application of different organic waste materials. *Pedobiologia*, 45: 396–407.
- Eriksen-Hamel N.S., Whalen J.K. (2008): Earthworms, soil mineral nitrogen and forage production in grass-based hayfields. *Soil Biology and Biochemistry*, 40: 1004–1010.
- Geissen V., Peña-Peña K., Huerta E. (2009): Effects of different land use on soil chemical properties, decomposition rate and earthworm communities in tropical Mexico. *Pedobiologia*, 53: 75–86.
- Hairiah K., Williams S.E., Bignell D., Swift M., Van Noordwijk M. (2001): Effects of land use change on belowground biodiversity.

- ASB Lecture Note 6A. International Centre for Research in Agroforestry, Bogor.
- Huerta E., Rodriguez-Olan J., Evia-Castillo I., Montejo-Meneses E., De la Cruz-Mondragón M., Garcia-Hernandez R., Uribe S. (2007): Earthworms and soil properties in Tabasco, Mexico. *European Journal of Soil Biology*, 43: S190–S195.
- Lavell P., Bignell D., Lepage M. (1997): Soil function in a changing world: The role of invertebrate ecosystem engineers. *European Journal of Soil Biology*, 33: 159–193.
- Leroy B.L.M., Schmidt O., Van den Bossche A., Reheul D., Moens M. (2008): Earthworm population dynamics as influenced by the quality of exogenous organic matter. *Pedobiologia*, 52: 139–150.
- Lowe C.N., Butt K.R. (2002): Influence of organic matter on earthworm production and behaviour: A laboratory-based approach with applications for soil restoration. *European Journal of Soil Biology*, 38: 173–176.
- Lu R.K. (2000): *Analysis of Soil Characteristics*. China Agricultural Sciencetech Press, Beijing.
- Ouellet G., Lapen D.R., Topp E., Sawada M., Edwards M. (2008): A heuristic model to predict earthworm biomass in agroecosystems based on selected management and soil properties. *Applied Soil Ecology*, 39: 35–45.
- Pathak H., Rao D.L.N. (1998): Carbon and nitrogen mineralization from added organic matter in saline and alkali soils. *Soil Biology and Biochemistry*, 30: 695–702.
- Qiao Y.H., Wu W.L., Xu Q., Chen K. (2001): Relationship between production input and secondary succession of earthworm population in salinity transforming region of North China – A case study in Quzhou county. *Chinese Journal of Applied Ecology*, 12: 414–416.
- Raw F. (1959): Estimating earthworm populations by using formalin. *Nature*, 184: 1661–1662.
- Reeleder R.D., Miller J.J., Ball Coelho B.R., Roy R.C. (2006): Impacts of tillage, cover crop, and nitrogen on populations of earthworms, microarthropods, and soil fungi in a cultivated fragile soil. *Applied Soil Ecology*, 33: 243–257.
- Schoeneberger P.J., Wysocki D.A., Benham E.C., Broderson W.D. (2002): *Field Book for Describing and Sampling Soils*. Version 1.1. National Soil Survey Center, Natural Resources Conservation Services. US Department of Agriculture Lincoln, Nebraska.
- Sumner M.E. (1995): Sodic soils: new perspectives. In: Naidu R., Sumner M.E., Rengasamy P. (eds.): *Australian Sodic Soils: Distribution, Properties and Management*. CSIRO, Melbourne, 1–34.
- Tao J., Gu W., Griffiths B., Liu X.J., Xu Y.J., Zhang H. (2012): Maize residue application reduces negative effects of soil salinity on the growth and reproduction of the earthworm *Aporrectodea trapezoides*, in a soil mesocosm experiment. *Soil Biology and Biochemistry*, 49: 46–51.
- Wessells M.L.S., Bohlen P.J., McCartney D.A., Subler S., Edwards C.A. (1997): Earthworm effects on soil respiration in corn agroecosystems receiving different nutrient inputs. *Soil Biology and Biochemistry*, 29: 409–412.
- Whalen J.K., Parmelee R.W., Edwards C.A. (1998): Population dynamics of earthworm communities in corn agroecosystems receiving organic or inorganic fertilizer amendments. *Biology and Fertility of Soils*, 27: 400–407.
- Wichern J., Wichern F., Joergensen R.G. (2006): Impact of salinity on soil microbial communities and the decomposition of maize in acidic soils. *Geoderma*, 137: 100–108.
- Wu W.J., Qiao Y.H., Xu Q., Chen K. (2001): An ecological study on earthworm in farmland of salinity transforming area in North China Plain. *Acta Ecologica Sinica*, 21: 1109–1113.
- Yin W.Y. (1992): *China Subtropical Soil Animal*. Science Press, Beijing.

Received on June 5, 2012

Accepted on November 20, 2012

---

*Corresponding author:*

Dr. Tao Jun, Beijing Normal University, State Key Laboratory of Earth Surface Process and Resource Ecology, Beijing, P.R. China  
 phone: + 8610 5880 9998, fax: + 8610 5880 9998, e-mail: juntao@bnu.edu.cn

---