

# Physiological response of *Monimopetalum chinense* to light stress under habitat fragmentation

C.Y. Hao<sup>1,2</sup>, R. Fan<sup>1</sup>, H.S. Wu<sup>1</sup>, X.P. Zhang<sup>2,3</sup>, L. Wang<sup>2,3</sup>, W.L. Chen<sup>2,3</sup>, Z.L. Chen<sup>4</sup>

<sup>1</sup>*Spice and Beverage Research Institute, Chinese Academy of Tropical Agricultural Sciences, Wanning, P.R. China*

<sup>2</sup>*Provincial Key Laboratory of Conservation and Employment of Biological Resources, Anhui Normal University, Wuhu, P.R. China*

<sup>3</sup>*Environment and Ecological Safety of Colleges in Anhui Province, Anhui Normal University, Wuhu, P.R. China*

<sup>4</sup>*The Administration Bureau of Dapanshan National Natural Reserve, Pan'an, P.R. China*

## ABSTRACT

To determine the effect of light stress under fragmental habitat on the physiology, this paper investigated the physiological responses of *Monimopetalum chinense* with different light intensities in the Xianyu Mountains (Anhui, China). The study showed that both weak and intense light brought about by habitat fragmentation could improve antioxidant enzymes activities, and promote electrical conductivity and malondialdehyde content of *M. chinense* leaves. However, too strong light could inhibit photosynthesis rates, superoxide dismutase, catalase, and ascorbate peroxidase activities. In addition, the characteristics of leaves were affected by light intensity at the fragmental habitat. Specifically, intense light was disadvantageous to photosynthesis and antioxidant enzymes of the species. Our results suggest that the biodiversity conservation of *M. chinense* is necessary, and that light intensity should be considered carefully when implementing conservation efforts.

**Keywords:** endangered; photosynthesis rate; antioxidant enzymes; electrical conductivity; lipid peroxidation

Human alteration of the global environment has triggered the 'sixth major extinction event' in the history of life, as well as it caused widespread changes in the abundance and distribution of organisms. Most forest ecosystems worldwide were fragmented by logging and other human disturbance, resulting in highly dissected landscape patterns (Turner 1996). One notable consequence of fragmentation is the increase in forest edge/interior ratio in remaining forest patches. An altered microclimatic environment around forest edges can directly influence plant reproduction, recruitment, and many others (Tomimatsu and Ohara 2004). Light is one of the main factors influencing the establishment and development of many forest species; it also plays a major role in the ecology of intrinsic species. However, exces-

sive light energy can photoinhibit photosynthesis, leading to reduced photochemical efficiency and conductance, increased temperature of leaves, impairment of CO<sub>2</sub> assimilation rate, and photo-oxidative destruction of photosynthetic apparatus through the formation of reactive oxygen species (Valladares and Pearcy 1997).

*Monimopetalum chinense*, a woody vine member of the Celastraceae, grows in a limited mountain area on the board between Anhui, Jiangxi, and Hubei Provinces in China (Anonymous 1999). As for its capsular fruit and isomerous flowers, this new Celastraceous species belongs to the Celastroideae-Euonymaceae; it is related closely to *Euonymus*, but with many differing aspects. As a monotypic genus, *Monimopetalum* contributes significantly in the evolutionary history of

---

Supported by the Natural Science Foundation of Anhui Province, Project No. 050430501.

Celastraceae species (Sytsma et al. 2002). However, due to anthropogenic disturbance and environmental deterioration, its habitat has been destroyed and is fragmented continuously, causing the species to be classified as endangered in the official list of Chinese rare species. Several studies were carried out, focusing on the biological and ecological factors associated with *M. chinense* decline (Ren 1989, Xie and Tan 1998, Hao et al. 2008, Zhang et al. 2008). However, studies on its physiological response to microclimate change under habitat fragmentation were not extensive. In the present study, the physiological responses of *M. chinense* under fragmental habitat were investigated to determine the impact of light on physiology, as well as to explore the most favorable light range and endangering mechanisms.

## MATERIALS AND METHODS

**Site description.** The study region (30°00'N, 117°18'E) is located in the Shangkeng Village at the Xianyu Mountains of Anhui Province. The two communities (A and B) of *M. chinense* are isolated by a highway and by some buildings constructed as recent as 10 years ago, with more than 50 m apart. Community A corresponds to an area of more than 5000 m<sup>2</sup>; it is undisturbed and in good condition. Meanwhile, community B was seriously destroyed by logging in 1998, and continues suffering from human disturbance. The light intensity in community B is much higher than in community A.

Seven sites were selected for sampling, specifically those from the middle to the edge of communities A and B. Light intensity was measured in each site at 10:30 a.m. on July 15, 2007 and July 13, 2008. The mean values for the two years were 40 (site I), 150 (site II) and 230 (site III)  $\mu\text{mol}/\text{m}^2/\text{s}$  for community A, and 460 (site IV), 720 (site V), 1190 (site VI) and 1650 (site VII)  $\mu\text{mol}/\text{m}^2/\text{s}$  for community B. At each site, healthy and adult individuals with root collar diameters between 10 to 15 mm were selected for sampling. The third pairs of leaves on the two-year-old branches were used to determine all physiological parameters at the closest time as possible. Net photosynthesis rate and stomatal conductance were measured directly in-situ. Other parameters were sampled into an ice pot and measured in a temporary laboratory. This laboratory was built in the Shangkeng Village.

**Analyses.** Light intensity, photosynthetic rate, and stomatal conductance were measured with a CI-340 portable photosynthesis system (CID Inc.,

Washington, USA). The leaf areas were measured with a Li-Cor 3100 (Li-Cor, Lincoln, Nebraska, NE) area meter and then dried at 68°C for 72 h and weighed.

The estimation of lipid peroxidation (i.e., malondialdehyde or MDA) was conducted according to the method of Velikova et al. (2000). Leaf membrane permeability was expressed as percentage of total electrolyte leakage and measured by electrical conductivity (Sukumaran and Weiser 1972). Total chlorophyll content was estimated according to the equations proposed by Wellburn (1994). The activity of superoxide dismutase (SOD) (EC 1.15.1.1) was determined by the photochemical method described by Giannopolitis and Ries (1977). Catalase (CAT) (EC 1.11.1.6) activity was assayed in a reaction mixture containing the enzyme and 25 mmol/l H<sub>2</sub>O<sub>2</sub>. Next, H<sub>2</sub>O<sub>2</sub> decomposition was conducted at 240 nm (Cakmak and Marschner 1992). Determination of ascorbate peroxidase (APX) (EC 1.11.1.11) activity was performed as described by Nakano and Asada (1981). Peroxidase (POD) (EC 1.11.1.7) activity was assayed using the method of Luck (1963). The measurements were made in five replicates.

Statistical analyses were carried out by one-way analysis of variance (ANOVA) using Student's *t*-test for the evaluation of significance of differences. A correlation analysis was applied to test the relationship of physiological characteristics in the SPSS software.

## RESULTS

**Leaf parameters in different habitats.** Leaf parameters of *M. chinense* varied under different light intensities, and significant differences ( $P < 0.05$ ) were observed (Table 1). With light intensity increase, the fresh weight per leaf area, water content per leaf area, and ratio of chlorophyll a/b first decreased, then finally increased. Chlorophyll content increased gradually with decrease in light intensity, but it was reduced when light intensity was too weak; the opposite was observed for fresh weight per leaf area and chlorophyll a/b. The stomatal conductance increased with light intensity at about 40–460  $\mu\text{mol}/\text{m}^2/\text{s}$ , but decreased when it was over 460  $\mu\text{mol}/\text{m}^2/\text{s}$ .

**Photosynthesis rate in different habitats.** Net photosynthesis rate increased linearly with light intensity at about 40–460  $\mu\text{mol}/\text{m}^2/\text{s}$ , but decreased rapidly when light intensity was over 460  $\mu\text{mol}/\text{m}^2/\text{s}$  (Figure 1). Moreover, it was significantly ( $P < 0.05$ ) enhanced by 641% from 40 to 460  $\mu\text{mol}/\text{m}^2/\text{s}$ ,

Table 1. The characteristics of *Monimopetalum chinense* leaves with different light intensities under fragmental habitats

Site	Light intensity ( $\mu\text{mol}/\text{m}^2/\text{s}$ )	Fresh weight	Water content	Stomatal conductance ( $\mu\text{mol H}_2\text{O}/\text{m}^2/\text{s}$ )	Content of chlorophyll (mg/g DW)	Chlorophyll a/b
		per leaf area	per leaf area			
		(mg/cm <sup>2</sup> )				
I	40	13.49 ± 1.15 <sup>a</sup>	9.96 ± 0.52 <sup>cde</sup>	65.85 ± 7.36 <sup>c</sup>	2.63 ± 0.20 <sup>a</sup>	2.60 ± 0.14 <sup>d</sup>
II	150	13.97 ± 0.87 <sup>a</sup>	9.51 ± 0.38 <sup>e</sup>	84.42 ± 12.92 <sup>bc</sup>	2.83 ± 0.19 <sup>b</sup>	2.75 ± 0.12 <sup>cd</sup>
III	230	14.18 ± 0.67 <sup>a</sup>	10.22 ± 0.47 <sup>cde</sup>	126.47 ± 19.20 <sup>ab</sup>	2.61 ± 0.15 <sup>bc</sup>	2.82 ± 0.09 <sup>cd</sup>
IV	460	14.11 ± 1.22 <sup>a</sup>	11.53 ± 0.44 <sup>bcd</sup>	145.87 ± 23.68 <sup>a</sup>	2.31 ± 0.06 <sup>cd</sup>	2.91 ± 0.15 <sup>bcd</sup>
V	720	15.35 ± 1.22 <sup>ab</sup>	11.77 ± 1.10 <sup>bc</sup>	133.84 ± 10.29 <sup>a</sup>	1.95 ± 0.21 <sup>de</sup>	3.01 ± 0.09 <sup>bc</sup>
VI	1190	16.14 ± 0.84 <sup>ab</sup>	12.50 ± 0.53 <sup>ab</sup>	105.00 ± 17.99 <sup>abc</sup>	1.70 ± 0.12 <sup>ef</sup>	3.22 ± 0.15 <sup>ab</sup>
VII	1650	17.61 ± 1.25 <sup>b</sup>	13.87 ± 0.77 <sup>a</sup>	84.29 ± 7.73 <sup>bc</sup>	1.24 ± 0.11 <sup>f</sup>	3.51 ± 0.08 <sup>a</sup>

All values are means ± S.E. ( $n = 5$ ). Different letters after S.E. indicate a significant difference ( $P < 0.05$ )

but significantly ( $P < 0.05$ ) declined by 46% from 460 to 1650  $\mu\text{mol}/\text{m}^2/\text{s}$ , indicating that intensive light is not favorable for the photosynthesis rate of *M. chinense*.

**Electrical conductivity and MDA content.** Electrical conductivity proportionally increased with the increase in light intensity; it was lowest at 40  $\mu\text{mol}/\text{m}^2/\text{s}$  and highest at 1650  $\mu\text{mol}/\text{m}^2/\text{s}$  (Figure 2). MDA content decreased gradually with the decline in light intensity. These results indicate that the peroxidation of *M. chinense* membrane lipids is intensified when exposed to intense light. Moreover, correlation analysis also showed that there was a very significant positive correlation between electrical conductivity and MDA content ( $r = 0.75^*$ ,  $n = 8$ ).

**Antioxidant enzymes.** SOD (Figure 3a) increased to a maximum at 460  $\mu\text{mol}/\text{m}^2/\text{s}$  and then decreased.

The trends of CAT activity (Figure 3b) and APX activity (Figure 3c) were quite similar, with activities first decreasing, then increasing, and finally decreasing with light intensity increase. Correlation analysis showed that there was a significant positive correlation among the two enzymes ( $P < 0.01$ ). However, POD presented a different trend (Figure 3d). When light intensity was lower than 460  $\mu\text{mol}/\text{m}^2/\text{s}$ , antioxidant enzymes activities declined first and then increased; this indicates that both weak and intense lights can improve the metabolism of ROS. When light intensity was more than 460  $\mu\text{mol}/\text{m}^2/\text{s}$ , the activities of SOD, CAT, and APX began to decrease, indicating that the light stress exceeded its tolerance. The result was similar to the related studies (Sofa et al. 2003).




Figure 1. Net photosynthesis rate of *M. chinense* leaves with different light intensities under fragmental habitats. Data correspond to mean ± S.E. ( $n = 5$ ). Different letters after S.E. indicate a significant difference ( $P < 0.05$ )




Figure 2. Electrical conductivity (a) and MDA content (b) of *M. chinense* leaves under different light intensity in fragmental habitats. Data correspond to mean  $\pm$  S.E. ( $n = 5$ ). Different letters above the bars indicate a significant difference ( $P < 0.05$ )

## DISCUSSION

Light is the only energy source required for photosynthesis in higher plants; it is a basic requirement for plant growth and development. However, light can also harm the so-called sun plants if it is too weak to satisfy photosynthesis requirements, or if the light energy absorbed by leaves of shade

plants exceeds the capability of photosynthetic apparatus (Valladares and Pearcy 1997). Our results revealed that *M. chinense* demonstrated similar characteristics as shade plants (Niinemets et al. 1998), indicating that this species can endure shady habitat to a certain extent. Studies on its stem and leaf anatomy proved that this species prefer shaded habitats (Ren 1989). The results




Figure 3. Antioxidant enzymes, SOD (a), CAT (b), APX (c), and POD (d) of *M. chinense* leaves with different light intensities under fragmental habitats. Data correspond to mean  $\pm$  S.E. ( $n = 5$ ). Different letters above the bars indicate a significant difference ( $P < 0.05$ )

presented by Xie and Zhang (1999) showed that this species is suitable for habitats with medium canopy density. Based on the analysis, *M. chinense* belongs to the shade-living plant group, but it cannot endure light that is too weak.

Intense light induces severe lipid peroxidation due to the removal of hydrogen by reactive oxygen species from unsaturated fatty acid, thus leading to lipid radical formation (Sofo et al. 2003). Dimerization and polymerization of proteins also occur, and these are considered as most damaging to the membranes (Bor et al. 2003). In relation, lipid peroxidation corresponds to the basic cell membrane reactive damage in cellular mechanisms. The electrical conductivity and MDA content under 1650  $\mu\text{mol}/\text{m}^2/\text{s}$  increased by 48% and by 151%, respectively, compared with their lowest values. These indicate that lipid peroxidation in *M. chinense* leaves is very sensitive to changes in light intensity. In addition to carotenoids, antioxidative systems play significant roles in protecting plants from the negative effects of reactive oxygen species resulting from light energy excess (Demmig-Adams 1998). The antioxidant enzymes, SOD, CAT, POD, and APX, play vital roles in scavenging these destructive oxidant species. By catalyzing the detoxification of  $\text{O}_2^- \cdot$  to  $\text{O}_2$  and  $\text{H}_2\text{O}_2$ , SOD can prevent  $\text{O}_2^- \cdot$ -driven cell damage; meanwhile, CAT, POD, and APX can remove  $\text{H}_2\text{O}_2$  (Cakmak and Marschner 1992, Aravind and Prasad 2004). Our current study also showed that SOD, CAT, POD, and APX activities changed with light intensity, indicating a close correlation between light intensity and the physiological behavior of the species.

In general, low levels of stress can lead to an increase of antioxidant enzymes activities and remove  $\text{O}_2^- \cdot$  (Yao et al. 2008). However, when accumulation of reactive oxygen species under more severe stress condition exceeds the removal capacity of the antioxidant system, oxidative damage occurs, including peroxidation of membrane lipids, destruction of photosynthetic and inactivation pigments, and inactivation of photosynthetic enzymes (Foyer et al. 1994, Sofo et al. 2003). Nevertheless, light intensity above 460  $\mu\text{mol}/\text{m}^2/\text{s}$  might lead to an imbalance between antioxidant defense and amount of active oxygen species, thereby resulting in more severe stress in the physiology of *M. chinense*.

*M. chinense* prefers to live at the foot or slope of hills of low latitudes, making it vulnerable to the effects of human activities. With excessive tree cutting and serious destruction of the eco-environment, more and more *M. chinense* populations are now located

in fragmental habitats. Small population size, isolation of populations from each other, and habitat light disturbance could be the three key factors associated with increased risks of species extinction. Based on this study, *M. chinense* has adapted to low light intensity, as high levels of light intensity is disadvantageous to its photosynthesis and antioxidant enzymes. The change in solar radiation under habitat fragmentation has influenced the physiology and ecology of *M. chinense*. According to the reports gathered from our field work, the seeds and seedlings of *M. chinense* are seldom found in fragmented habitats.

Based on the above analysis, the present work demonstrated that the change in light intensity under fragmental habitat could markedly affect the leaves characteristics, photosynthesis rates, and antioxidant enzymes activities of *M. chinense*. Although adaptable to shady habitat, both intense and too weak light are disadvantageous to this species. Therefore, light intensity should be considered carefully when implementing biodiversity conservation for *M. chinense*, especially during transplant conservation. The results obtained in this study can improve our knowledge on the effects of habitat fragmentation on *M. chinense* and impending efforts to implement biodiversity conservation.

## Acknowledgements

Authors would like to thank Yu Zhang, Wenliang Li and several other anonymous for their assistance.

## REFERENCES

- Anonymous (1999): Flora of China. Vol 45. No. 3. Science Press, Beijing, 96. (In Chinese)
- Aravind P, Prasad M.N.V. (2004): Zinc alleviates cadmium-induced oxidative stress in *Ceratophyllum demersum* L.: a free floating freshwater macrophyte. *Plant Physiology and Biochemistry*, 41: 391–397.
- Bor M., Özdemir F., Türkan I. (2003): The effect of salt stress on lipid peroxidation and antioxidants in leaves of sugarbeet *Beta vulgaris* L. and wild beet *Beta maritima* L. *Plant Science*, 164: 77–84.
- Cakmak I, Marschner H. (1992): Magnesium deficiency and high light intensity enhance activities of superoxide dismutase, ascorbate, peroxidase, and glutathione reductase in bean leaves. *Plant Physiology*, 98: 1222–1227.
- Demmig-Adams B. (1998): Survey of thermal energy dissipation and pigment composition in sun and shade leaves. *Plant and Cell Physiology*, 39: 474–482.

- Foyer C.H., Descouvières P., Kunert K. (1994): Protection against oxygen radicals: an important defence mechanism studied in transgenic plants. *Plant, Cell and Environment*, *17*: 507–523.
- Giannopolitis N., Ries S.K. (1977): Superoxide dismutase I: occurrence in higher plants. *Plant Physiology*, *59*: 309–314.
- Hao C.Y., Zhang X.P., Li W.L., Zhang Y. (2008): Spatial distribution of *Monimopetalum chinense* populations in different forest types. *Acta Ecologica Sinica*, *28*: 2900–2908. (In Chinese)
- Luck H. (1963): Peroxidase. In: Bergmeyer H.U. (ed.): *Methods of Enzymic Analysis*. Academic Press, New York, 895–897.
- Nakano Y., Asada K. (1981): Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant and Cell Physiology*, *22*: 867–880.
- Niinemets Ü., Kull O., Tenhunen J.D. (1998): An analysis of light effects on foliar morphology, physiology and light interception in temperate deciduous woody species of contrasting shade tolerance. *Tree Physiology*, *18*: 681–696.
- Ren X.F. (1989): The study of stem and leaf anatomy of an endemic species of China – *Monimopetalum chinense* Rehd. *Journal of Anhui Normal University*, *4*: 22–30. (In Chinese)
- Sofa A., Dichio B., Xiloyannis C., Masia A. (2003): Effects of different irradiance levels on some antioxidant enzymes and on malondialdehyde content during rewatering in olive tree. *Plant Science*, *166*: 293–306.
- Sukumaran N.P., Weiser C.J. (1972): An excised leaflet test for evaluating potato frost tolerance. *HortScience*, *7*: 467–468.
- Sytsma K.J., Morawetz J., Pires J.C., Nepokroeff M., Conti E., Zjhra M., Hall J.C., Chase M.W. (2002): Urticalean rosids: circumscription, rosid ancestry, and phylogenetics based on *rbcL*, *trnL-F*, and *ndhF* sequences. *American Journal of Botany*, *89*: 1531–1546.
- Tomimatsu H., Ohara M. (2004): Edge effects on recruitment of *Trillium camschatcense* in small forest fragments. *Biological Conservation*, *17*: 509–519.
- Turner I.M. (1996): Species loss in fragments of tropical rain forests: a review of the evidence. *Journal of Applied Ecology*, *33*: 200–209.
- Valladares F., Pearcy R.W. (1997): Interactions between water stress, sun-shade acclimation, heat tolerance and photoinhibition in the sclerophyll *Heteromeles arbutifolia*. *Plant, Cell and Environment*, *20*: 25–36.
- Velikova V., Yordanov I., Edreva A. (2000): Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Science*, *151*: 59–66.
- Wellburn A.R. (1994): The spectral determination of chlorophyll *a* and *b*, as well as total carotenoids, using various solvents with spectrophotometers of different resolutions. *Journal of Plant Physiology*, *144*: 307–313.
- Xie G.W., Tan C.M. (1998): On the population existing state and conservation of *Monimopetalum* Rehd. of a new recorded genus in Hubei. *Journal of Plant Resources and Environment*, *7*: 38–42. (In Chinese)
- Xie G.W., Zhang Z.Y. (1999): The geographical distribution and population spatial pattern of *Monimopetalum chinense*. *Ecologic Science*, *18*: 7–11. (In Chinese)
- Yao Y.N., Yang Y.Q., Li Y., Lutts S. (2008): Intraspecific responses of *Fagopyrum esculentum* to enhanced ultraviolet B radiation. *Plant Growth Regulation*, *56*: 297–306.
- Zhang X.P., Hao C.Y., Fan R., Li W.L., Zhang Y., Kan X.B. (2008): Population structure of *Monimopetalum chinense* and its relationship with environmental factors. *Chinese Journal of Applied Ecology*, *19*: 474–480. (In Chinese)

Received on March 4, 2009

---

*Corresponding author:*

Prof. X.P. Zhang, Anhui Normal University, College of Life Science, Wuhu 241000, P.R. China  
phone: + 86 553 3883 539, e-mail: pinghenxu@sina.com.cn

---