

<https://doi.org/10.17221/453/2018-PSE>

Fertilization effects on CH₄, N₂O and CO₂ fluxes from a subtropical double rice cropping system

YE YUAN^{1,2}, XIAOQIN DAI^{1*}, HUIMIN WANG¹

¹Qianyanzhou Ecological Research Station, Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing, P.R. China

²Anhui Provincial Key Laboratory of the Conservation and Exploitation of Biological Resources, College of Life Sciences, Anhui Normal University, Wuhu, P.R. China

*Corresponding author: daiqx@igsnr.ac.cn

Citation: Yuan Y., Dai X.Q., Wang H.M. (2019): Fertilization effects on CH₄, N₂O and CO₂ fluxes from a subtropical double rice cropping system. *Plant Soil Environ.*, 65: 189–197.

Abstract: A 2-year field study was conducted in a double rice cropping system in southern China to examine the effect of fertilization on CH₄, N₂O and CO₂ fluxes using static opaque chambers and gas chromatographs. Two treatments were set up including conventional fertilization with a rate of 358 kg N/ha per year in forms of compound fertilizer and urea, and a control with no fertilizer application. The results indicated that fertilization did not have a significant effect on CH₄ fluxes and led to a significantly higher cumulative N₂O emission in the two years of observation period. Fertilization promoted CO₂ fluxes by increasing the autotrophic respiration instead of heterotrophic respiration. By combining the global warming impact of CH₄, N₂O and CO₂, it was found that fertilization increased the area-scaled but not the yield-scaled global warming impact. These results indicated that, according to the current amount of nitrogen applied, fertilization may increase the global warming effect of paddy fields in this region. However, the appropriate dose of nitrogen fertilizer application is still a reasonable agricultural management due to the comprehensive consideration of production and environmental impacts.

Keywords: nitrogen addition; rice paddy; greenhouse gases; greenhouse gas intensity

Carbon dioxide (CO₂), methane (CH₄) and nitrous oxide (N₂O) are the most important greenhouse gases (IPCC 2013). Rice paddies have been identified as a major CH₄ source, emitting between 33 and 40 Tg CH₄/year (IPCC 2013). N₂O emissions could also be pronounced in paddy fields with frequent drying-wetting alternation (Kudo et al. 2014). Fertilization has been commonly adopted to maintain rice yields, and the application of fertilizer may regulate greenhouse gas emissions (Jassal et al. 2011, Abalos et al. 2014). Therefore, a study on the effect of fertilization on greenhouse gas emissions is imperative for estimation of greenhouse gas emissions and designing possible mitigation strategies in the paddy soils.

The addition of nitrogen fertilizer can raise crop yield and biomass, which may have a positive effect

on soil carbon (C) sequestration as a result of the biomass input into the soil from crop residues and roots (Liu and Greaver 2009). Soil carbon decomposition might be retarded by fertilization when the litter quality and the growth rate of decomposers are both changed. Experimental additions of nitrogen to cropland usually result in increased N₂O emission (Wu et al. 2017). For CH₄, different studies got the converse conclusion that fertilization could promote (Shang et al. 2011), inhibit (Yao et al. 2012) or have no significant effect (Dan et al. 2001).

Most of the field measurements of greenhouse gas fluxes were conducted in temperate regions (Shang et al. 2011). However, over 60% of all nitrogen fertilizer will be used in the tropics and subtropics by 2020 (Galloway et al. 2003). China has a large area of paddy fields with

Supported by the National Natural Science Foundation of China, Grants No. 31700415 and 41001179, and by the Natural Science Foundation of Anhui Province, Grant No. 1808085QC60.

90% of them being located in the subtropics (Wang et al. 2015). The environmental conditions in subtropical regions are different from those in temperate regions not only in climate but also highly acidic with low base cation concentrations (Mo et al. 2008). Therefore, it is necessary to do more research on the effect of fertilization especially in subtropical region where greenhouse gas emissions remain to be uncertain.

In the present study, two years of measurements were conducted on CH₄, N₂O and CO₂ fluxes from paddy fields in southern China. It is hypothesized that fertilization would significantly increase CO₂ and N₂O fluxes mainly through promoting autotrophic respiration and applying substrate for the formation of N₂O.

MATERIAL AND METHODS

Site description. The experimental fields are located at the Qianyanzhou Ecological Research Station (QYZ, 26°44'46"N, 115°04'05"E) in the Jiangxi province, southern China. The site consists of a typical red soil hilly region with a subtropical monsoon climate. The mean air temperature is 18.0°C, and the coldest and warmest months are January and July, respectively. The region receives an average total annual precipitation of 1509 mm. The soils (equivalent to Plinthudults in the US Soil Taxonomy) are predominately sand and loam and were formed from alluvial fans of nearby rivers. The soil had 580, 310 and 110 g/kg sand, silt and clay, respectively. The topsoil (0–10 cm) had an organic carbon content of 9.33 g/kg and a total nitrogen content of 1.00 g/kg, and the soil pH was 4.98 and the bulk density was 1.32 g/cm³ before the experiment. Double cropping of rice in a year is the primary cropping system in this area.

Experimental design. The experimental design consisted of a completely randomized block with four replicates. Each plot had an area of 120 m². Two fertilization levels (conventional fertilization (CF) and no fertilization (NF)) was adopted in this study. Compound fertilizer (N:P:K = 15%:6.6%:12.5%) and urea were applied at a rate of 358 kg N/ha per year to the fertilized plots according to the local typical agricultural management. The fertilizer was spread into rice fields evenly. The time and rates of nitrogen applied to fertilized plots are shown in Table 1. Besides, a non-vegetated subplot of 2 m × 2 m was set up in each plot to determine heterotrophic respiration. The subplots received the same fertilization and agricultural management practices as the other part of the plots.

Measurements of CH₄, N₂O and CO₂ fluxes. Static chamber and gas chromatograph (GC) method was used to simultaneously measure CH₄, N₂O and CO₂ fluxes. The design of the static chamber was described in our previous article (Yuan et al. 2016). In July 2012, two stainless steel bases were installed in each plot. One base was installed in a place with normal vegetation with five seedlings of rice in it and the other was installed in the middle of the non-vegetated subplot of 2 m × 2 m. Greenhouse gases fluxes were measured in the same frequency in the two static chambers. Five gas samples were collected from each chamber using 100 mL plastic syringes at 10-min interval twice each week between 8:00 AM and 12:00 PM throughout the growing season. In fallow seasons, samples were collected once a week. The gas samples were analysed using a gas chromatograph (GC System, 7890A, Agilent Technologies, USA). The CH₄, N₂O and CO₂ fluxes were calculated from linear or nonlinear changes in the gas concentrations over time (Yuan et al. 2016).

Table 1. Field management practices of the fertilized treatment

Rice growing year	Crop type	Growing season	Fertilization date	Nitrogen application rate (kg N/ha)
The first year (2012–2013)	late rice	2012-7-30–2012-11-14	2012-7-30 2012-8-10	71.7 (compound fertilizer) 107.5 (urea)
	fallow	2012-11-15–2013-4-23		
	early rice	2013-4-24–2013-7-23	2013-4-24 2013-5-3	71.7 (compound fertilizer) 107.5 (urea)
	late rice	2013-7-24–2013-11-2	2013-7-26 2013-8-12	71.7 (compound fertilizer) 107.5 (urea)
The second year (2013–2014)	fallow	2013-11-3–2014-4-24		
	early rice	2014-4-25–2014-7-25	2014-4-19 2014-4-29	71.7 (compound fertilizer) 107.5 (urea)

<https://doi.org/10.17221/453/2018-PSE>

Auxiliary measurements. The air temperature and precipitation data were collected from an on-site automatic meteorological station adjacent to the experimental plots. When the fields were flooded, the standing water depth was monitored using a steel ruler. To determine soil chemical properties, soil samples were taken in May 2012, July 2013 and July 2014 from five points of each plot at a depth of 0–10 cm and then combined and mixed thoroughly. Soil pH was measured by a calibrated pH meter (Mettler Toledo, Greifensee, Switzerland) after shaking a soil-water (1:2.5 w:v) suspension for 30 min. Soil nitrate (NO_3^- -N) and ammonium (NH_4^+ -N) were extracted with 1 mol/L KCl (1:5 soil:water ratio) and measured using a continuous flow analyser (Skakar, Breda, the Netherlands). Total carbon (TC) and total nitrogen (TN) were measured with an elemental analyser (Elementar, Vario Max, Hanau, Germany). The results of soil chemical properties can be found in Table 2. To determine biomass and yields, the weights of the rice grain, straw and root were recorded from the average of three 1 m² areas at harvest. The grain, straw or root samples, which had masses of approximately 100 g, were oven-dried to a constant weight at 65°C for 24 h after determining the wet weight.

Data analysis. The seasonal or annual cumulative emissions of CH_4 , N_2O and CO_2 were computed as the sum of all daily CH_4 , N_2O and CO_2 fluxes, respectively. For those days when measurements were not conducted, the emissions were calculated by linear interpolation using the data from the nearest days. The annual cumulative greenhouse gas emissions were the sum of the two growing seasons (late rice and early rice) and the fallow season. CO_2 emissions of the vegetated plots were considered to be the ecosystem respiration including autotrophic

respiration (plant-derived CO_2 efflux from soil) and heterotrophic respiration (microbial respiration). CO_2 emissions of the non-vegetated subplots of 2 m × 2 m in each plot were considered to be heterotrophic respiration. Autotrophic respiration was calculated by CO_2 emissions of the vegetated plots minus CO_2 emissions of the non-vegetated plots.

The combined global warming impact of the three greenhouse gases (GWI) ($\text{g CO}_{2\text{eq}}/\text{m}^2$) = CH_4 emissions × 25 + N_2O emissions × 298 + CO_2 emissions (Jiang et al. 2010) as the global warming potentials of 25 for CH_4 and 298 for N_2O over 100-years (IPCC 2007). The greenhouse gas intensity (GHGI) is calculated by dividing GWI by rice grain yield (Mosier et al. 2006).

All statistical analyses were performed with the SPSS 19.0 (SPSS Inc., Chicago, USA) and Origin 7.0 (Origin Lab Corporation, Northampton, USA) software. The effect of fertilization and growing seasons on CH_4 , N_2O and CO_2 fluxes was determined to be statistically significant using repeated measures of the analysis of variance (ANOVA). An independent samples *t*-test was used to analyse the difference in biomass, yields, cumulative emissions of greenhouse gases, GWI and GHGI between paddy fields with conventional fertilization and no fertilization.

RESULTS

Air temperature, precipitation, flood water depth, rice biomass and yields. The average temperature was 18.5°C in the first year and 18.6°C in the second year. In the four rice growing seasons, the average temperature was 22.7, 26.0, 24.2 and 25.7°C, respectively. The precipitation of the second year (1397.6 mm) was lower than that of the first year (1518.7 mm), especially in the late rice growing season (Figure 1a).

Table 2. Soil chemical properties of paddy fields with conventional fertilization (CF) and no fertilization (NF) during the observation period

Time	Treatment	pH	NO_3^- -N	NH_4^+ -N	TC	TN
			(mg/kg)			
May (2012)	CF	4.92 ± 0.09 ^a	0.32 ± 0.16 ^a	5.78 ± 0.36 ^a	9.23 ± 0.31 ^a	1.01 ± 0.02 ^a
	NF	5.03 ± 0.07 ^a	0.22 ± 0.08 ^a	5.35 ± 0.27 ^a	9.42 ± 0.39 ^a	1.00 ± 0.03 ^a
July (2013)	CF	4.80 ± 0.04 ^a	0.88 ± 0.32 ^a	39.53 ± 0.99 ^a	9.68 ± 0.43 ^a	0.94 ± 0.03 ^a
	NF	4.92 ± 0.06 ^a	0.82 ± 0.41 ^a	14.62 ± 1.05 ^b	9.28 ± 0.27 ^a	0.89 ± 0.03 ^a
July (2014)	CF	5.17 ± 0.04 ^a	0.51 ± 0.15 ^a	27.88 ± 8.61 ^a	11.13 ± 0.45 ^a	1.09 ± 0.03 ^a
	NF	5.34 ± 0.07 ^a	0.22 ± 0.04 ^a	9.39 ± 1.62 ^a	10.27 ± 0.47 ^a	1.04 ± 0.03 ^a

Different letters within the same column between different treatments indicate significant differences between CF and NF at $P < 0.05$. Data shown are means ± standard errors of four spatial replicates. TC – total carbon; TN – total nitrogen




Figure 1. Seasonal variations in daily air temperature and precipitation (a), and flood water depth (b) during the observation period

The water-logging period lasted for 37, 43, 35 and 43 days in the four rice growing seasons and the corresponding average flood water depth was 4.53, 3.42, 2.44 and 3.22 cm, respectively (Figure 1b). The application of fertilizer significantly increased biomass and rice yields in the second year but not in the first year (Figure 2a,b). In the second year, the rice biomass and yields in the late rice growing season were 7922 and 3084 kg/ha for NF treatment, while the values were 11 982 and 4316 kg/ha for CF treatment. In the early rice growing season of the second year, the biomass and yields also increased by 52.6% and 57.9% under fertilization. Biomass

yield after fertilization significantly increased CO_2 consumption during photosynthesis.

Greenhouse gas fluxes and overall global warming impact. CH_4 fluxes showed significant seasonal variations (Figure 3a). In the rice growing seasons when the fields were water logged, high CH_4 fluxes were observed. The peak value of CH_4 fluxes could reach up to 37.9 and 31.1 $\text{mg C/m}^2/\text{h}$ for CF and NF treatments, respectively. In the winter fallow when no rice was grown and standing water was absent, CH_4 fluxes remained at very low levels and even dropped to negative values sometimes. Among the four growing seasons, the late rice growing season in the second year had the




Figure 2. (a) Total biomass and (b) grain yields of paddy fields with conventional fertilization (CF) and no fertilization (NF). The data shown are means of the four replicates for individual treatments. Vertical bars indicate standard errors. Different letters indicate significant differences between CF and NF at $P < 0.05$

<https://doi.org/10.17221/453/2018-PSE>




Figure 3. Seasonal variations in (a) CH₄ fluxes; (b) N₂O fluxes and (c) CO₂ fluxes of paddy fields with conventional fertilization (CF) and no fertilization (NF) and the fertilization time and amount (b). The data shown are means of the four replicates for individual treatments. Vertical bars indicate standard errors

lowest CH₄ emissions both for CF and NF (Table 3). By comparing the cumulative emissions, fertilization was found to have no significant effect on CH₄ emission in any of the four growing seasons (Table 3).

N₂O fluxes of NF treatment stayed low throughout the entire observation period. N₂O fluxes of CF treatment were pulse-like and showed apparent seasonal variations (Figure 3b). The peaks were observed after fertilization except in the late rice growing season in the first year when no peak was ever found. The highest N₂O fluxes were observed in the late rice growing season in the second year with an average value of 105.74 μg N/m²/h. Fertilization significantly increased cumulative N₂O emissions by 2.16 times in the observation period of two years (Table 3). This finding can partially confirm our hypothesis.

CO₂ fluxes showed significant seasonal variations. They were high in the rice growing seasons and relatively low in the winter fallow period (Figure 3c). The average fluxes in the four growing seasons were 187, 190, 128 and 143 mg C/m²/h for CF and 132, 139, 98 and 24 for NF. Fertilization increased the cumulative CO₂ emissions by 34.5% and 21.0% in the first year and the second year (Table 3). By di-

viding the ecosystem respiration into autotrophic and heterotrophic respiration, it was found that the autotrophic respiration was significantly higher with fertilization except in the late rice growing season of the second year while the heterotrophic respiration, CO₂ fluxes measured from the non-vegetated subplots, was not affected by fertilization (Figure 4). This is consistent with our hypothesis that fertilization would significantly increase CO₂ fluxes mainly through promoting autotrophic respiration.

GWI was mainly contributed by CO₂ and CH₄ (Table 4). Fertilization significantly increased GWI of the whole observation period of two years. GHGI, the aggregate emissions of the three greenhouse gases expressed on grain yield bases, in two years of the observation period were 5.79 and 5.69 kg CO_{2eq} kg/yield for CF and NF treatments. GHGI was not significantly affected by fertilization (Table 4).

DISCUSSION

Seasonal and interannual variation of CH₄, N₂O and CO₂ fluxes. Our results showed that N₂O fluxes stayed quite low throughout the entire observation

Table 3. Cumulative greenhouse gas emissions of paddy fields with conventional fertilization (CF) and no fertilization (NF) during the observation period

	Treatment	CH ₄ (kg C/ha)	N ₂ O (kg N/ha)	CO ₂ (kg C/ha)
Late rice (2012)	CF	206 ± 58 ^a	0.32 ± 0.10 ^a	5346 ± 145 ^a
	NF	167 ± 32 ^a	0.21 ± 0.09 ^a	3655 ± 313 ^b
Fallow (2012–2013)	CF	0.34 ± 0.17 ^a	0.51 ± 0.14 ^a	2492 ± 179 ^a
	NF	0.85 ± 0.38 ^a	0.14 ± 0.04 ^a	2231 ± 299 ^a
Early rice (2013)	CF	152 ± 35 ^a	0.41 ± 0.03 ^a	4307 ± 258 ^a
	NF	154 ± 23 ^a	0.19 ± 0.02 ^a	3147 ± 391 ^b
Late rice (2013)	CF	43 ± 9 ^a	2.65 ± 0.87 ^a	3197 ± 246 ^a
	NF	88 ± 31 ^a	0.35 ± 0.01 ^b	2613 ± 237 ^a
Fallow (2013–2014)	CF	0.11 ± 0.49 ^a	0.38 ± 0.16 ^a	2487 ± 370 ^a
	NF	-0.20 ± 0.27 ^a	0.35 ± 0.18 ^a	2222 ± 200 ^a
Early rice (2014)	CF	137 ± 24 ^a	0.19 ± 0.05 ^a	3523 ± 164 ^a
	NF	86 ± 16 ^a	0.18 ± 0.10 ^a	2476 ± 312 ^b
The first year (2012~2013)	CF	358 ± 92 ^a	1.24 ± 0.18 ^a	12 146 ± 358 ^a
	NF	321 ± 51 ^a	0.54 ± 0.15 ^b	9033 ± 584 ^b
The second year (2013~2014)	CF	180 ± 31 ^a	3.22 ± 1.01 ^a	9207 ± 695 ^a
	NF	174 ± 30 ^a	0.88 ± 0.20 ^a	7311 ± 505 ^a
Two years	CF	538 ± 120 ^a	4.46 ± 0.89 ^a	21 352 ± 1013 ^a
	NF	495 ± 76 ^a	1.41 ± 0.22 ^b	16 344 ± 1070 ^b

Different letters within the same column between different treatments indicate significant differences between CF and NF at $P < 0.05$. Data shown are means ± standard errors of four spatial replicates

period except that several pulse-like emissions were observed after fertilization (Figure 3b). This is consistent with many previous studies (Bronson et al. 1997, Yao et al. 2012) and the low fluxes could be explained by reduction of N₂O to N₂ through the denitrification

process in the anaerobic environment (Yuan et al. 2016, Smith et al. 2018). The highest N₂O fluxes were observed in the late rice growing season of the second year when CH₄ fluxes were relatively low (Figure 3a,b). The discrepancy could be caused by the differences in




Figure 4. (a) Autotrophic respiration and (b) heterotrophic respiration of paddy fields with conventional fertilization (CF) and no fertilization (NF). The data shown are means of the four replicates for individual treatments. Vertical bars indicate standard errors. Different letters indicate significant differences between CF and NF at $P < 0.05$

<https://doi.org/10.17221/453/2018-PSE>

Table 4. The carbon dioxide equivalents of CH₄, N₂O, CO₂ and their global warming impact (GWI) and greenhouse gas intensity (GHGI) of paddy fields with conventional fertilization (CF) and no fertilization (NF) during the observation period

	Treatment	Carbon dioxide equivalents (kg CO _{2eq} /ha)			GWI (kg CO _{2eq} /ha)	GHGI (kg CO _{2eq} /kg yield)
		CH ₄	N ₂ O	CO ₂		
The first year (2012~2013)	CF	11 946 ± 3059 ^a	580 ± 84 ^a	44 534 ± 1314 ^a	57 061 ± 3379 ^a	6.76 ± 0.78 ^a
	NF	10 723 ± 1687 ^a	251 ± 69 ^b	33 122 ± 2142 ^b	44 096 ± 3737 ^b	5.68 ± 0.43 ^a
The second year (2013~2014)	CF	5989 ± 1032 ^a	1507 ± 475 ^a	33 758 ± 2547 ^a	41 254 ± 2413 ^a	5.03 ± 0.65 ^a
	NF	5806 ± 989 ^a	411 ± 94 ^a	26 806 ± 1853 ^a	33 023 ± 2355 ^a	5.75 ± 0.19 ^a
Two years	CF	17 936 ± 4004 ^a	2086 ± 417 ^a	78 292 ± 3714 ^a	98 314 ± 4620 ^a	5.79 ± 0.35 ^a
	NF	16 529 ± 2527 ^a	662 ± 102 ^b	59 928 ± 3923 ^b	77 119 ± 6046 ^b	5.69 ± 0.25 ^a

Different letters within the same column between different treatments indicate significant differences between CF and NF at $P < 0.05$. Data shown are means ± standard errors of four spatial replicates

water flooding and distribution of precipitation. In the late rice growing season of the second year, the precipitation was sparse (Figure 1a) and the duration of flooding was shorter with lower flood water depth (Figure 1b). The less flood water and precipitation might have caused the lower CH₄ fluxes and the higher N₂O fluxes. Water condition has been proved to play an important role in regulating CH₄ and N₂O fluxes (Li et al. 2005). Significant CH₄ emissions could occur when soil is submerged and redox potential stays low. When the flooded layer disappears and the soil becomes aerobic with elevated redox, potential N₂O emissions could be significant (Hou et al. 2000). Previous studies also found that paddies may have lower CH₄ emissions (Berger et al. 2013) while N₂O emissions would be pronounced when aerobic environments were formed (Kudo et al. 2014).

CO₂ fluxes were lower in the winter fallow than in rice growing seasons (Figure 3c). CO₂ fluxes included autotrophic respiration and heterotrophic respiration. However, autotrophic respiration was generally zero in the winter fallow since there was no vegetation growing. Another possible reason is temperature, which is one of the most important factors affecting soil respiration (Smith et al. 2018). The low temperature could partially explain the small CO₂ fluxes in winter. CO₂ fluxes were lower in the second year than the first year (Figures 3c and 4a,b). Temperature and plant biomass were good proxies for variations in both autotrophic and heterotrophic capacity for respiration (Flanagan and Johnson 2005, Smith et al. 2018). However, in the present study, both of them did not differ interannually (Figure 1a) and could not be the reason for the interannual variability of CO₂ fluxes. Soil moisture was another

important environmental factor that controlled seasonal and interannual variation in CO₂ fluxes besides temperature (Flanagan and Johnson 2005). In the present study, the precipitation and flood water of the second year were both less than the first year, and this might have caused the difference in CO₂ fluxes between the two years. Ecosystem respiration is controlled by the complex interaction of environmental and biotic factors (Han et al. 2007). Further study needs to be conducted to identify all of the other factors.

Effects of fertilization on CH₄, N₂O and CO₂ fluxes. The application of fertilizer did not significantly affect CH₄ fluxes (Table 3) and this is in line with previous studies (Dan et al. 2001) although other studies also found that the application of fertilizers would either promote (Liu and Greaver 2009) or decrease (Yao et al. 2012) CH₄ emissions in paddy fields. The effects of N fertilization on CH₄ production, oxidation and transport are complex, involving many biochemical processes (Schimel 2000). N fertilization promotes plant growth and provides more substrates for CH₄ production, but this will also relieve the methanotrophs and promote CH₄ oxidation in the intermittent flooded paddy fields, especially in the drainage period (Banger et al. 2012). Dan et al. (2001) found that N fertilization stimulated both CH₄ production and CH₄ oxidation and resulted in no significant effect on CH₄ emissions.

Our findings are in line with many other studies that fertilization would promote N₂O emission (Millar et al. 2010), and this might be caused by applying substrate for nitrification and denitrification (Azam et al. 2002, Yao et al. 2012). However, a significant promotion effect of fertilization on N₂O emissions was found only in the late rice growing season of the

second year with less flood water and precipitation. Soil might have a chance to be aerobic when the paddy fields were logged with lower water depth for shorter periods. As it was well known that N_2O could be reduced to N_2 through the denitrification process in the anaerobic environment in many previous studies (Bronson et al. 1997, Yao et al. 2012, Smith et al. 2018), this study speculated that most of the nitrogen fertilizer might have been transformed to N_2 by denitrification in the other three growing seasons as the fields were flooded (Hayatsu et al. 2008). This result suggests that water condition associated with fertilizer application, plays an important role in regulating N_2O fluxes in paddy fields.

As for CO_2 emitted by ecosystem respiration, fertilization showed a significant promotion effect through promoting autotrophic respiration other than heterotrophic respiration (Figure 4). This indicated that fertilization affected ecosystem respiration through promoting crop growth. Although the biomass of the fertilized treatment was not significantly higher in the first year, the root activity and respiration may have been promoted. A study by Xu and Wan (2008) also found that soil respiration was greater in the fertilized plots than in the unfertilized plots and this was attributable to stimulated plant growth, root activity and respiration. It should be noted that both phosphate and potash were also applied into the paddy fields, but their effects on greenhouse gas fluxes were not examined in the present study.

Effects of fertilization on rice biomass, yields and the integrated greenhouse effects of CH_4 , N_2O and CO_2 . Fertilization significantly increased rice yields of the second year but not those of the first year (Figure 2b). It was noticed that all the fields had been fertilized for years before the experiment was conducted. Previous studies found that most of the intensive agricultural production in China was over-fertilized (Vitousek et al. 2009). The strong retention of external nitrogen inputs could remain available for the subsequent crops (Ju et al. 2009). The retention of nitrogen might have maintained the high rice yields for NF in the first year. This indicates that it is feasible to reduce nitrogen application rates in the excessive nitrogen fertilization agro-ecosystem without sacrificing crop yield.

To assess the combined climatic impacts from the three greenhouse gases as affected by fertilization, the carbon dioxide equivalents of CH_4 , N_2O and CO_2 were calculated. It was found that fertilization increased the area-scaled global warming impact but did not change the yield-scaled global warming impact (Table 4). This

is because fertilization promoted greenhouse gas emissions and increased crop yields, simultaneously. Other factors affecting the environment, such as mineral nitrogen leaching and ammonia volatilization (Griggs et al. 2007, Zhao et al. 2009) should be included in the further research to provide a more comprehensive evaluation of fertilization in paddy fields.

Both N_2O and CO_2 emissions were significantly promoted by the application of fertilizer, while CH_4 had no difference between treatments with and without fertilization in a paddy field in southern China. Fertilization increased CO_2 fluxes by promoting the autotrophic respiration instead of heterotrophic respiration. Fertilization increased the area-scaled global warming impact of CH_4 , N_2O and CO_2 in the two years of observation period while the yield-scaled global warming impact did not differ significantly with fertilization. For NF, stopping fertilizer use did not reduce rice yields immediately in the first year as the retention of external nitrogen inputs in the previous growing seasons might have maintained the rice yields. An appropriate amount of nitrogen fertilization should be an effective way to maintain crop yields without increasing environmental impact.

REFERENCES

- Abalos D., Sanchez-Martin L., Garcia-Torres L., van Groenigen J.W., Vallejo A. (2014): Management of irrigation frequency and nitrogen fertilization to mitigate GHG and NO emissions from drip-fertigated crops. *Science of The Total Environment*, 490: 880–888.
- Azam F., Müller C., Weiske A., Benckiser G., Ottow J.C.G. (2002): Nitrification and denitrification as sources of atmospheric nitrous oxide – Role of oxidizable carbon and applied nitrogen. *Biology and Fertility of Soils*, 35: 54–61.
- Bronson K.F., Neue H.U., Singh U., Abao E.B.Jr. (1997): Automated chamber measurements of methane and nitrous oxide flux in a flooded rice soil: I. Residue, nitrogen, and water management. *Soil Science Society of America Journal*, 61: 981–987.
- Banger K., Tian H.Q., Lu C.Q. (2012): Do nitrogen fertilizers stimulate or inhibit methane emissions from rice fields? *Global Change Biology*, 18: 3259–3267.
- Berger S., Jang I., Seo J., Kang H., Gebauer G. (2013): A record of N_2O and CH_4 emissions and underlying soil processes of Korean rice paddies as affected by different water management practices. *Biogeochemistry*, 115: 317–332.
- Dan J.G., Krüger M., Frenzel P., Conrad R. (2001): Effect of a late season urea fertilization on methane emission from a rice field in Italy. *Agriculture, Ecosystems and Environment*, 83: 191–199.
- Flanagan L.B., Johnson B.G. (2005): Interacting effects of temperature, soil moisture and plant biomass production on ecosystem respiration in a northern temperate grassland. *Agricultural and Forest Meteorology*, 130: 237–253.

<https://doi.org/10.17221/453/2018-PSE>

- Galloway J.N., Aber J.D., Erisman J.W., Seitzinger S.P., Howarth R.W., Cowling E.B., Cosby B.J. (2003): The nitrogen cascade. *BioScience*, 53: 341–356.
- Griggs B.R., Norman R.J., Wilson C.E., Slaton N.A. (2007): Ammonia volatilization and nitrogen uptake for conventional and conservation tilled dry-seeded, delayed-flood rice. *Soil Science Society of America Journal*, 71: 745.
- Han G.X., Zhou G.S., Xu Z.Z., Yang Y., Liu J.L., Shi K.Q. (2007): Biotic and abiotic factors controlling the spatial and temporal variation of soil respiration in an agricultural ecosystem. *Soil Biology and Biochemistry*, 39: 418–425.
- Hayatsu M., Tago K., Saito M. (2008): Various players in the nitrogen cycle: Diversity and functions of the microorganisms involved in nitrification and denitrification. *Soil Science and Plant Nutrition*, 54: 33–45.
- Hou A.X., Chen G.X., Wang Z.P., Van Cleemput O., Patrick W.H. (2000): Methane and nitrous oxide emissions from a rice field in relation to soil redox and microbiological processes. *Soil Science Society of America Journal*, 64: 2180–2186.
- IPCC (2013): Climate change 2013: The physical science basis. In: Stocker T.F., Qin D., Plattner G.-K., Tignor M., Allen S.K., Boschung J., Nauels A., Xia Y., Bex V., Midgley P.M. (eds.): Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge, New York, Cambridge University Press, 1535.
- IPCC (2007): Climate change 2007: The physical science basis. In: Solomon S., Qin D., Manning M., Chen Z., Marquis M., Averyt K.B., Tignor M., Miller H.L. (eds): Contribution of Working Group I to the fourth assessment report of the Intergovernmental Panel on Climate Change. Cambridge, Cambridge University Press.
- Jassal R.S., Black T.A., Roy R., Ethier G. (2011): Effect of nitrogen fertilization on soil CH₄ and N₂O fluxes, and soil and bole respiration. *Geoderma*, 162: 182–186.
- Jiang C.M., Yu G.R., Fang H.J., Cao G.M., Li Y.N. (2010): Short-term effect of increasing nitrogen deposition on CO₂, CH₄ and N₂O fluxes in an alpine meadow on the Qinghai-Tibetan Plateau, China. *Atmospheric Environment*, 44: 2920–2926.
- Ju X.T., Xing G.X., Chen X.P., Zhang S.L., Zhang L.J., Liu X.J., Cui Z.L., Yin B., Christie P., Zhu Z.L., Zhang F.S. (2009): Reducing environmental risk by improving N management in intensive Chinese agricultural systems. *Proceedings of the National Academy of Sciences of the United States of America*, 106: 3041–3046.
- Kudo Y., Noborio K., Shimozono N., Kurihara R. (2014): The effective water management practice for mitigating greenhouse gas emissions and maintaining rice yield in central Japan. *Agriculture, Ecosystems and Environment*, 186: 77–85.
- Li C.S., Frohling S., Xiao X.M., Moore B., Boles S., Qiu J., Huang Y., Salas W., Sass R. (2005): Modeling impacts of farming management alternatives on CO₂, CH₄, and N₂O emissions: A case study for water management of rice agriculture of China. *Global Biogeochemical Cycles*, 19.
- Liu L.L., Greaver T.L. (2009): A review of nitrogen enrichment effects on three biogenic GHGs: The CO₂ sink may be largely offset by stimulated N₂O and CH₄ emission. *Ecology Letters*, 12: 1103–1117.
- Millar N., Robertson G.P., Grace P.R., Gehl R.J., Hoben J.P. (2010): Nitrogen fertilizer management for nitrous oxide (N₂O) mitigation in intensive corn (maize) production: An emissions reduction protocol for US Midwest agriculture. *Mitigation and Adaptation Strategies for Global Change*, 15: 185–204.
- Mo J.M., Zhang W., Zhu W.X., Gundersen P., Fang Y.T., Li D.J., Wang H. (2008): Nitrogen addition reduces soil respiration in a mature tropical forest in southern China. *Global Change Biology*, 14: 403–412.
- Mosier A.R., Halvorson A.D., Reule C.A., Liu X.J. (2006): Net global warming potential and greenhouse gas intensity in irrigated cropping systems in northeastern Colorado. *Journal of Environmental Quality*, 35: 1584–1598.
- Schimel J.P. (2000): Global change: Rice, microbes and methane. *Nature*, 403: 375–377.
- Shang Q.Y., Yang X.X., Gao C.M., Wu P.P., Liu J.J., Xu Y.C., Shen Q.R., Zou J.W., Guo S.W. (2011): Net annual global warming potential and greenhouse gas intensity in Chinese double rice-cropping systems: A 3-year field measurement in long-term fertilizer experiments. *Global Change Biology*, 17: 2196–2210.
- Smith K.A., Ball T., Conen F., Dobbie K.E., Massheder J., Rey A. (2018): Exchange of greenhouse gases between soil and atmosphere: Interactions of soil physical factors and biological processes. *European Journal of Soil Science*, 54: 10–20.
- Vitousek P.M., Naylor R., Crews T., David M.B., Drinkwater L.E., Holland E., Johnes P.J., Katzenberger J., Martinelli L.A., Matson P.A., Nziguheba G., Ojima D., Palm C.A., Robertson G.P., Sanchez P.A., Townsend A.R., Zhang F.S. (2009): Agriculture. Nutrient imbalances in agricultural development. *Science*, 324: 1519–1520.
- Wang W., Sardans J., Lai D.Y.F., Wang C., Zeng C., Tong C., Liang Y., Peñuelas J. (2015): Effects of steel slag application on greenhouse gas emissions and crop yield over multiple growing seasons in a subtropical paddy field in China. *Field Crops Research*, 171: 146–156.
- Wu X., Liu H.F., Fu B.J., Wang Q., Xu M., Wang H.M., Yang F.T., Liu G.H. (2017): Effects of land-use change and fertilization on N₂O and NO fluxes, the abundance of nitrifying and denitrifying microbial communities in a hilly red soil region of southern China. *Applied Soil Ecology*, 120: 111–120.
- Xu W.H., Wan S.Q. (2008): Water- and plant-mediated responses of soil respiration to topography, fire, and nitrogen fertilization in a semiarid grassland in northern China. *Soil Biology and Biochemistry*, 40: 679–687.
- Yao Z., Zheng X., Dong H., Wang R., Mei B., Zhu J. (2012): A 3-year record of N₂O and CH₄ emissions from a sandy loam paddy during rice seasons as affected by different nitrogen application rates. *Agriculture, Ecosystems and Environment*, 152: 1–9.
- Yuan Y., Dai X., Wang H., Xu M., Fu X., Yang F. (2016): Effects of land-use conversion from double rice cropping to vegetables on methane and nitrous oxide fluxes in southern China. *PLoS ONE* 11(5):e0155926.
- Zhao X., Xie Y.X., Xiong Z.Q., Yan X.Y., Xing G.X., Zhu Z.L. (2009): Nitrogen fate and environmental consequence in paddy soil under rice-wheat rotation in the Taihu lake region, China. *Plant and Soil*, 319: 225–234.

Received on June 26, 2018

Accepted on March 25, 2019

Published online on April 12, 2019