

<https://doi.org/10.17221/471/2021-PSE>

## Biochar-based fertiliser improved the yield, quality and fertiliser utilisation of open field tomato in karst mountainous area

MENG ZHANG, YANLING LIU, QUANQUAN WEI, LINGLING LIU, JIULAN GOU\*

*Institute of Soil and Fertiliser, Guizhou Academy of Agricultural Sciences, Guiyang, Guizhou, P.R. China*

\*Corresponding author: [gzgoujiulan@163.com](mailto:gzgoujiulan@163.com)

**Citation:** Zhang M., Liu Y.L., Wei Q.Q., Liu L.L., Gou J.L. (2022): Biochar-based fertiliser improved the yield, quality and fertiliser utilisation of open field tomato in karst mountainous area. *Plant Soil Environ.*, 68: 163–172.

**Abstract:** Biochar-based fertiliser (BF) is beneficial to improve yield and quality, but the effect of BF on open field tomato remains unclear, especially in karst mountainous areas. The objective of this study was to identify the application effect and optimum application rate of BF. A field experiment was carried out in Southwestern China from 2019 to 2020 to study the effects of different application amounts of BF on the yield, quality, nutrients accumulation and fertiliser utilisation of open field tomatoes. The results showed that compared with the traditional fertilisation practice, BF can significantly increase the yield of open field tomato by 5–9% (2019) and 12–23% (2020), and significantly reduce nitrate content and increase vitamin C content of fruits. Meanwhile, nutrient accumulations, agronomic efficiency, and recovery efficiency of BF treatments were all significantly improved. In conclusion, the BF rate of 2 326 kg/ha improves yield and fertiliser utilisation in open-field tomatoes and could be recommended for tomato production in karst mountainous areas.

**Keywords:** potential productivity; nutritive value; nutrient uptake; fertiliser management strategies

Karst landform is a kind of geological landscape with special hydrological conditions and geological characteristics. The distribution area of the karst landform accounts for about 13% of the global land area (Yan et al. 2021a). The karst ecosystem is fragile and prone to rocky desertification, the soil cultivation layer in the karst area is shallow, and nutrient leaching is also severe (Liu et al. 2016). These factors are very unfavourable to agricultural production and economic development in karst regions.

About 900 million tons of agricultural and forestry waste and 3.8 billion tons of livestock and poultry manure can be collected in China every year (Hong et al. 2016). However, some of the resources are used to feed livestock and returned to the field at present, and about 20% are dumped or directly burned, which not only causes a serious waste of biomass resources but also leads to serious environmental

pollution problems (Yin et al. 2018). In the current agricultural production, biochar is used to effectively reduce this waste and improve the utilisation efficiency of biomass resources.

As an important renewable resource, biochar is prepared through pyrolysis of biomass raw materials at 300–700 °C (Yan et al. 2021b), such as crop straw (Luo et al. 2016), livestock manure (Wang et al. 2019), and agricultural and forestry processing waste (Kong et al. 2021). Biochar is rich in carbon and is not easily decomposed by microorganisms due to the recalcitrant nature of carbon compounds (Dumroese et al. 2011). Pituello et al. (2018) showed that biochar increased soil porosity and reduced bulk density, thus improving soil physical structure. In addition, biochar can increase soil nutrient content, cation exchange capacity (CEC), and water retention capacity, and affects the diversity and structure of soil

---

Supported by the National Natural Science Foundation of China, Grants No. 31860594 and 32060302; by the Subsidy Project from NSFC of Guizhou Academy of Agricultural Sciences, Project No. [2021]32, and by the Science and Technology Planning Project of Guizhou Province, Project No. [2020]1Y087.

microbial communities directly (Kumar et al. 2022). Meanwhile, biochar can also participate in the soil nutrient cycle of farmland ecosystems (Zhang et al. 2021a) and has an important effect on the conversion, migration, and distribution of soil nutrients through their physicochemical properties or interaction with the soil (Sun et al. 2021). Furthermore, biochar's rich chemical-reaction-type functional groups also endow the biochar with extremely strong adsorption capacity, which makes biochar a good fertiliser synergist and makes it possible to compound with mineral fertilisers to improve its utilisation (Yan et al. 2021a). Therefore, biochar has enormous potential in improving soil physicochemical properties, soil fertility, and crop productivity. However, the global price of biochar was 192–621 USD/t, which was much higher than the average wage in some countries (Maroušek and Trakal 2021). This may make it difficult to realise high-dose application in current agricultural production.

Biochar-based fertiliser (BF) prepared by mixing biochar and fertilisers can not only increase the nutrients but also release the fertilisers slowly (Dong et al. 2021). This can help to improve soil quality and sequester carbon oxide while supplying nutrients to crops (Zheng et al. 2021). Hydrophilic groups in biochar can make the surfaces of BF absorb water in the soil, such as hydroxyl and carboxyl groups. In this case, the hydrophilicity and water retention capacity of biochar-based fertilisers can be improved by adjusting to the ratios of O/C and (N + O)/C (Pan et al. 2017). In addition, BF can directly increase the content of soil nutrients and affect the migration and transformation of original nutrients in the soil (Das and Ghosh 2021). Moreover, BF can enable nutrients to migrate toward plants and increase the ionic potential of plant root membranes, which can control plants to absorb nutrient cations and anions (Chew et al. 2020). In addition, BF increases the soil flux and the potential difference on the root membrane surface, which may help to save the energy required by plants to absorb nutrients (Joseph et al. 2015). Generally, biochar is mostly alkaline (pH 5–12) and can improve acidic soil (Liu et al. 2013). Therefore, BF can also adjust the structure of microbial communities in acidic soil to reduce the acidic functional groups, which is conducive to the growth and reproduction of microorganisms (Yang et al. 2021). Furthermore, BF can also inhibit the activity of nitrifying bacteria and reduce nitrogen loss by 11–12% in soil (Zhang et al. 2020). Therefore, we believe that BF may be conducive to managing agriculture in the karst mountains to increase crop yield and quality, fertiliser efficiency, and farmers' income.

Though the importance of biochar is increasingly understood, the existing research mainly focuses on greenhouse gas emission reduction and soil remediation. The influence of BF on yield and quality improvement is still poorly known, especially in the rare ecological environment. Therefore, strengthening the research on the macro and micromechanisms of BF will help to improve the comprehensive understanding level of BF and the global agricultural planting management technology. This study was conducted through a two-year field experiment to evaluate how BF potentially improve the biological effect and economic benefits of open field tomatoes in the karst mountainous areas. Therefore, the objectives of this study were to (i) research the effect of BF with different application amounts on the yield and quality of tomatoes; (ii) calculate nutrients accumulation of BF with different application amounts, and (iii) evaluate the fertiliser utilisation rates of BF application in open-field tomatoes. It is anticipated that the study results will be useful for formulating novel management ways for improving crop production in karst areas.

## MATERIAL AND METHODS

**Site description and sampling.** The field experiment was carried out in Southwest China in 2019 and 2020. This region is a typical karst landform, with an average annual rainfall of 1 168.3 mm, an average annual temperature of 15.6 °C, and an average frost-free period of 276 days. The planting crops in the experiment region were open field tomatoes from May to October each year. The tested soil was a typical zonal yellow soil, which as a Dystric Luvisol (Ultisols) is widely distributed in south karst China. The basic physicochemical properties of the surface soil (0–20 cm) in the experiment region were: pH 5.9, organic carbon of 27.1 g/kg, total nitrogen of 0.31 g/kg, available phosphorus of 32.1 mg/kg, and available potassium of 108.5 mg/kg. Due to the strong leaching capacity of yellow soil in karst regions, the tested soil is weakly acidic and the soil fertility is low.

The tomato cultivar used in the experiment was Anma 161 (Guizhou Jinnong Technology Co., Ltd., Guiyang, China). The raw material of biochar was distillers' grains which are biomass waste generated from the production process of distilled spirits (Kweichow Moutai (Group) Circular Economy Industrial Investment and Development Co., Ltd., Zunyi, China), and the carbonisation temperature was 550 °C using a biological carbonisation furnace (SSDP-

<https://doi.org/10.17221/471/2021-PSE>

5000, Huai'an Huadian Environmental Protection Machinery Manufacturing Co., Ltd., Huai'an, China). The basic physicochemical properties of biochar were: pH 8.8, organic carbon of 279.4 g/kg, total nitrogen of 7.6 g/kg, total phosphorus of 10.6 g/kg, and total potassium of 21.4 g/kg. Fertilisers used in the experiment included organic fertiliser (Guizhou Dibao Co., Ltd., Guiyang, China), compound fertiliser (N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O 15-15-15, Guizhou Xiyang Industrial Co., Ltd., Guiyang, China), and biochar-based fertilisers (BF, N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O 15-6-18, with a biochar content of 20%, Guizhou Institute of Soil and Fertiliser, Guiyang, China).

The BF was prepared using the SKJ – 120 flat-grinding extrusion granulator (Shanghai Jiale Electromechanical Group Co., Ltd., Shanghai, China) through the extrusion granulation method. The specification was in the shape of a cylinder of 1–2 cm. The raw materials used were urea (N 46%, Guizhou Chitianhua Group Co., Ltd., Zunyi, China), mono-ammonium phosphate (N-P<sub>2</sub>O<sub>5</sub> 10-50, Guizhou Chitianhua Group Co., Ltd., Zunyi, China), potassium sulfate (K<sub>2</sub>O 50%, Xinjiang Luobupo Hoevellite Co., Ltd., Korla, China), distillers' grains biochar and solid binder (Guizhou Jinnong Technology Co., Ltd., Guiyang, China). Firstly, the raw materials were crushed through a 0.38 mm standard screen and dried in a 45 °C constant temperature electric blast drying oven (DHG-9070A, Shanghai Jingqi Instrument Co., Ltd., Shanghai, China) for 24 h. Then, after all, kinds of raw materials were mixed evenly according to a certain proportion, fertiliser particles were formed by flat-grinding extrusion granulator. Finally, after the powder and small particles were removed through a 3.35 mm aperture screen, the fertiliser particles were baked in the drying oven at 45 °C for 6 h to obtain the finished BF. The surface of BF particles was flat

and the structure was uniform. The pH of BF was 6.8 and the average compressive strength was 257.2 N.

**Experimental design and management.** This experiment included six treatments, these being the control of no fertiliser (CK); compound fertiliser of 3 000 kg/ha (TFP, traditional fertilisation practice); BF of 1 875 (BF1); 2 250 (BF2); 2 625 (BF3) and 3 000 (BF4) kg/ha. Except for CK treatment, all fertilisation treatments were applied with organic fertiliser of 1 500 kg/ha at the same time. The specific application amount of the nutrients in the treatments were shown in Table 1. Before the tomato seedlings were transplanted, the fertilisers were spread evenly on the surface of the soil, and then a hoe was used to mix fertilisers and the soil evenly. Tomato seedlings were transplanted 10 days after the ridging and mulching. The planting density was  $2.7 \times 10^4$  plants/ha (with a plant spacing of 50 cm, row spacing of 70 cm). Each treatment was conducted in a random block and repeated three times, and the plot area was 33.6 m<sup>2</sup> (8.0 m × 4.2 m). The same fungicide, insecticide and herbicide were used in all treatments so that no diseases, pests or weeds occurred during the growing season. Other field management was consistent with the habits of local farmers.

**Sampling and measurement.** Soil samples (0–20 cm) were collected from 10 randomly selected spots of the main experimental area one day before fertilisation. The soil samples were composited and air-dried, ground and passed through 1 mm and 0.149 mm sieves for the determination of soil physicochemical characteristics. The physical and chemical properties of soil were determined according to Bao (2000), and the specific methods were as follows. Soil pH was measured in a 1:2.5 (soil:water ratio, w/v) extraction followed by reading using a pH meter (FE20K, Mettler

Table 1. Nutrients amounts of different treatments

Treatment	Basal dressing fertiliser (kg/ha)			Dressing fertiliser (kg/ha)		
	N	P	K	N	P	K
CK	–	–	–	–	–	–
TFP	225.00	98.24	186.70	225.00	98.24	186.70
BF1	281.25	49.12	280.05	–	–	–
BF2	337.50	58.94	336.05	–	–	–
BF3	393.75	68.77	392.07	–	–	–
BF4	450.00	78.59	448.09	–	–	–

N – nitrogen; P – phosphorus; K – potassium; CK – no fertiliser; TFP – traditional fertilisation practice; BF1 – 1 875 kg/ha of biochar-based fertiliser (BF); BF2 – 2 250 kg/ha of BF; BF3 – 2 625 kg/ha of BF; BF4 – 3 000 kg/ha of BF. N, P and K nutrients content in compound fertiliser were 15.0, 6.5 and 12.4%. N, P and K nutrients content in BF were 15.0, 2.6 and 14.9%

Toledo, Zurich, Switzerland). Soil organic carbon (SOC) was measured *via* the potassium dichromate external heating method. Total nitrogen was determined using the Kjeldahl method after digestion by a mixture of concentrated  $\text{HClO}_4$ - $\text{H}_2\text{SO}_4$ , without considering the influence of mineralisation. Available phosphorus (Olsen-P) was determined using the molybdenum blue method with a 0.5 mol/L  $\text{NaHCO}_3$  solution at a soil/solution ratio of 1:20, mainly in the form of calcium-based phosphates (Olsen et al. 1954). Available potassium was extracted using 1 mol/L  $\text{NH}_4\text{OAc}$  for 30 min and then determined with a flame photometer (FP640, Jingke, Shanghai, China).

Six plants from each experimental plot were sampled before the final harvest, which were used to test the plant nutrition and fruit quality. The tomato plants were divided into two parts: stems-leaf and fruit, which were dried to constant weight at 60 °C after a heat at 105 °C for 30 min. All dried samples were ground and passed through a 0.25 mm sieve and digested with a mixture of concentrated  $\text{H}_2\text{SO}_4$  and  $\text{H}_2\text{O}_2$  to determine N, P and K concentrations. The N concentration was determined with a continuous flow analyser (AA3, Seal Analytical Inc., Southampton, UK), P and K concentration was determined with an ICP-OES (optical emission spectroscopy with inductively coupled plasma, Longjumeau, HORIBA Jobin Ibon S.A.S., Paris, France). The uptake of N, P and K were calculated based on the dry mass and element concentration. At the same time, some fresh fruit samples were used to determine the quality indexes. For the determination of nitrate content, 2 g fresh fruits were taken, added 10 mL deionised water, and placed in a boiling water bath for 30 min. 0.1 mL of the extraction solution was taken and mixed with 0.4 mL of 5% salicylic acid-sulfuric acid solution at 25 °C for 20 min. The solution was added 9.5 mL of 8% NaOH and measured the absorbance using a visible spectrophotometer (UV-3600i Plus, Shimadzu, Tokyo, Japan). The vitamin C (VC) content was determined using an HPLC (high-performance liquid chromatography, LC-2040, Nexera-i, Shimadzu, Japan) after grinding, centrifuging and filtering with 10 mL 0.2% metaphosphoric acid. For the content of soluble sugar, 0.2 g fresh leaves were taken and added 10 mL distilled water to extract in a boiling water bath for 30 min. 0.5 mL of the extract was taken in a test tube, and 0.5 mL ethyl anthrone and 5 mL sulfuric acid were added to the extract. The test tube was placed in a boiling water bath and incubated for 1 min. After taking it out, it was naturally cooled to room tem-

perature and measured the absorbance at 630 nm. The content of soluble solids was detected by Abbe refractometer (NAR-1T, ATAGO, Tokyo, Japan). The final yield was an accumulation of the whole plot based on three batches of harvest.

#### Calculations and statistical analysis.

$$\text{Nutrient accumulation (kg/ha)} = \text{nutrient concentration (\%)} \times \text{dry mass (kg/ha)/100} \quad (1)$$

$$\text{Agronomic efficiency (AE, kg/kg)} = (\text{yield of the fertilised plot} - \text{yield of the no fertiliser plot}) / \text{applied nutrient rate} \quad (2)$$

$$\text{Recovery efficiency (RE, \%)} = (\text{total nutrient uptake of the fertilised plot} - \text{total nutrient uptake of no fertiliser plot}) / \text{applied nutrient rate} \times 100\% \quad (3)$$

The yield and nutrient uptake in the upper formulas were calculated according to dry mass.

In order to explore the best application amount of biochar based fertiliser on tomatoes, the Linear plus platform model was used to describe the response of tomato yield to BF application rate in the experiment. For the linear plus platform model, Y increased linearly with increasing X ranging from 0 to a critical value and reached the maximum at the critical value; after exceeding the critical value, Y remained the maximum no matter how X increased.

The linear plus platform model was:

$$Y = AX + B \quad (X \leq C); \quad Y = P \quad (X > C) \quad (4)$$

Y – yield (kg/ha); X – fertilisation application rate (kg/ha); A – regression coefficient; B – intercept; C – critical value; P – maximum yield (kg/ha).

Statistical analysis was performed using SPSS 18.0 (SPSS Inc., Chicago, USA). The data means were compared using the LSD (least significant difference) test ( $P < 0.05$ ). The figures were prepared with Origin 8.0 (Origin Lab Corporation, Northampton, USA).

## RESULTS AND DISCUSSION

**Effects of biochar-based fertiliser on the yield of tomato.** Yield is the first important factor to determine the price of tomatoes. Compared with CK treatment, the application of fertiliser (TFP and BF) significantly increased tomato production, with an average increase of 24 528 kg/ha in 2019 and an average increase of 38 620 kg/ha in 2020 (Figure 1). Compared with TFP treatment, the yield of tomatoes in BF-applied plots was increased by 3 471–6 918 kg/ha (5–9% increase) in 2019 and 8 199–15 995 kg/ha



<https://doi.org/10.17221/471/2021-PSE>




Figure 1. Effects of different fertilisation treatments on the yield of fresh tomato. CK – no fertiliser; TFP – traditional fertilisation practice; BF1 – 1 875 kg/ha of biochar-based fertiliser (BF); BF2 – 2 250 kg/ha of BF; BF3 – 2 625 kg/ha of BF; BF4 – 3 000 kg/ha of BF. Different letters indicate significant differences between different fertilisation treatments ( $P < 0.05$ ). Error bars represent the standard error of the mean

(12–23%) in 2020. The BF3 treatment showed the highest tomato yield in two years, which were 80 086 and 86 019 kg/ha, respectively. This indicates that BF could improve the yield of open field tomatoes and contribute to the improvement of tomato economic value. However, it is not that the more BF application, the higher the crop yield. Xu et al. (2021) reported that excessive application of BF might not continually improve plant growth, as the high C/N ratios might immobilise N and decrease plant N uptake. The optimal application amount of the BF was calculated using the linear plus platform model (Figure 2), and it was found that the maximum yield can be obtained when the application amount of the BF was 2 326 kg/ha. Therefore, the appropriate BF application rate is an important premise to ensure a high yield of tomatoes.

Studies demonstrated that the vast majority of biochar application approaches will never be economical if only productivity gains are realised (Bach et al. 2016). Thus, an alternative that has attracted growing interest in recent years is the development of BF, which could make the biochar technology more cost-effective due to the enhanced efficiency and associated lower application rates (Joseph et al. 2013). For instance, Puga et al. (2019a) observed a 26% average increase in maize yield and 12% higher N use efficiency for biochar with urea over conventional urea. Qian et al. (2014) observed rice yield increases of up to 24% using BF over conventional NPK in the field. In this study, BF increased the yield of tomatoes (5–9% in 2019 and 12–23% in 2020) in karst areas (Figure 1), which shows the positive responses of BF on crop productivity over conventional fertilisers. Sim et al. (2021) have pointed out that BF might

have superior characteristics for sustained release of nutrients as compared with conventional fertilisers, especially for controlled release of N fertilisers and reducing N losses under variable soil-plant-environment systems. Meanwhile, BF was more effective in eliminating those crop productivity constraints that conventional fertilisers could not sufficiently address, and have the potential to improve nutrient use efficiency and increase crop productivity by applying biochar in minor quantities (Melo et al. 2022).

**Effects of biochar-based fertiliser on the fruit quality of tomato.** Fruit quality is an important index of nutritional value and economic value. BF showed significant advantages in improving the yield and quality of crops (Sim et al. 2021), and the results of this study are similar to those of previ-




Figure 2. Effects of biochar-based fertiliser (BF) application rate on the yield of fresh tomato

<https://doi.org/10.17221/471/2021-PSE>

Table 2. Effects of different fertilisation treatments on nitrate, vitamin C, soluble sugar and total soluble solid of fresh tomato

Year	Treatment	Nitrate	Vitamin C	Soluble sugar	Total soluble solids
		(mg/kg)			
2019	CK	106.8 ± 3.6 <sup>d</sup>	130.1 ± 2.3 <sup>c</sup>	54.5 ± 2.3 <sup>b</sup>	45.0 ± 3.0 <sup>b</sup>
	TFP	134.2 ± 4.8 <sup>a</sup>	164.8 ± 6.7 <sup>b</sup>	56.2 ± 0.6 <sup>ab</sup>	45.0 ± 2.7 <sup>b</sup>
	BF1	115.8 ± 4.4 <sup>b</sup>	169.9 ± 3.9 <sup>ab</sup>	56.3 ± 0.7 <sup>ab</sup>	45.9 ± 0.8 <sup>ab</sup>
	BF2	114.6 ± 4.5 <sup>bc</sup>	176.0 ± 7.2 <sup>a</sup>	56.6 ± 0.7 <sup>ab</sup>	47.0 ± 1.8 <sup>ab</sup>
	BF3	107.9 ± 3.4 <sup>cd</sup>	180.0 ± 5.2 <sup>a</sup>	58.3 ± 0.7 <sup>a</sup>	49.2 ± 0.8 <sup>a</sup>
	BF4	106.4 ± 4.8 <sup>d</sup>	175.7 ± 8.3 <sup>a</sup>	58.0 ± 0.9 <sup>a</sup>	47.8 ± 2.0 <sup>ab</sup>
2020	CK	105.8 ± 4.8 <sup>b</sup>	154.2 ± 9.7 <sup>d</sup>	52.8 ± 2.1 <sup>c</sup>	42.5 ± 1.8 <sup>b</sup>
	TFP	126.1 ± 6.5 <sup>a</sup>	167.3 ± 5.0 <sup>c</sup>	56.5 ± 1.7 <sup>b</sup>	44.5 ± 1.5 <sup>b</sup>
	BF1	102.4 ± 3.0 <sup>bc</sup>	168.5 ± 5.5 <sup>c</sup>	58.4 ± 0.8 <sup>ab</sup>	47.8 ± 1.0 <sup>a</sup>
	BF2	98.2 ± 6.0 <sup>bc</sup>	183.6 ± 3.1 <sup>b</sup>	59.2 ± 1.5 <sup>a</sup>	48.9 ± 1.2 <sup>a</sup>
	BF3	97.6 ± 2.0 <sup>c</sup>	195.2 ± 3.9 <sup>a</sup>	60.3 ± 1.4 <sup>a</sup>	49.8 ± 1.4 <sup>a</sup>
	BF4	95.8 ± 2.3 <sup>c</sup>	186.3 ± 4.9 <sup>ab</sup>	60.0 ± 0.6 <sup>a</sup>	49.6 ± 1.0 <sup>a</sup>

CK – no fertiliser; TFP – traditional fertilisation practice; BF1 – 1 875 kg/ha of biochar-based fertiliser (BF); BF2 – 2 250 kg/ha of BF; BF3 – 2 625 kg/ha of BF; BF4 – 3 000 kg/ha of BF. Different letters in the same column indicates significant difference between different fertilisation treatments ( $P < 0.05$ )

ous studies (Sim et al. 2021). In this study, BF can significantly reduce the nitrate content in tomato fruits, and increase the content of VC, soluble sugar and the total soluble solid at the same time (Table 2). Compared with the TFP treatment, the BF treatments reduced the nitrate content in tomato fruits by 14–21% in 2019 and 19–24% in 2020. Tomato fruits in the BF4 treatment showed the lowest nitrate concentrations in the two years. BF treatments increased the VC content in tomato fruits by 3–9% in 2019 and 1–17% in 2020, and the VC content of tomato fruits treated with BF3 treatment was the highest. In addition, BF treatments also increased the content of the soluble sugar and the total soluble solids in tomato fruits, especially in 2020. Biochar is rich in mineral nutrients (Ca, Mg and Zn, etc.), which facilitates a balanced supply of nutrients (Liu et al. 2020). In addition, biochar possesses a strong capability of absorbing  $\text{NH}_4^+$ , which can regulate the rate of converting nitrogen from  $\text{NH}_4^+$  to  $\text{NO}_3^-$  in BF, and avoid the accumulation of nitrate due to excessive nitrogen uptake by crops (Khajavi-Shojaei et al. 2020). The relatively low application rate of BF (lower application rate of nitrogen) and the application of a variety of microelements may be beneficial to the increase of VC content in the fruit (Zhang et al. 2021b). Therefore, BF could improve the quality of open field tomatoes.

**Effects of biochar-based fertiliser on the nutrient accumulation of tomato.** Biochar adsorbs nutrients

from mineral fertiliser and later releases them to provide nutrients for plant growth, which could explain the persistent nutrients supply by BF (Yang et al. 2021). Compared with CK treatment, fertilisation significantly increased the accumulation of N, P and K by 100, 78, and 90%, respectively, in 2019 and 123, 93 and 108% in 2020 (Figure 3). Furthermore, compared with TFP treatment, these accumulations increased by 14–41, 6–32, and 6–42% respectively in 2019, and 6–33, 8–33, 15–47% respectively in 2020. The nutrient accumulations of N, P and K in the BF3 treatment were the highest, with 259.3, 37.4, 344.7 kg/ha respectively in 2019, and 263.3, 36.6, 351.2 kg/ha respectively in 2020. As we know, BF applied in soil could effectively act as the slow release of N fertiliser after one-time application which provides adequate N for plant growth (Ibrahim et al. 2020). On the other hand, the increased N uptake was associated with improved soil CEC, as soil with a high CEC is more likely to adsorb  $\text{NH}_4^+$ , which can effectively improve the utilisation of N in the soil (Liang et al. 2006). Moreover, Walter and Rao (2015) attributed the increased potassium uptake with BF to the presence of potassium in biochar itself and mineral fertiliser and enhanced potassium fertiliser use efficiency by biochar. It is worth noting that the increased potassium availability is vital for improving growth, biological nitrogen fixation, and the competitive ability of plants, as potassium is essential for bio-

<https://doi.org/10.17221/471/2021-PSE>




Figure 3. Effects of different fertilisation on nitrogen, phosphorus and potassium accumulations. CK – no fertiliser; TFP – traditional fertilisation practice; BF1 – 1 875 kg/ha of biochar-based fertiliser (BF); BF2 – 2 250 kg/ha of BF; BF3 – 2 625 kg/ha of BF; BF4 – 3 000 kg/ha of BF. Different letters indicate significant differences between different fertilisation treatments ( $P < 0.05$ ). Error bars represent the standard error of the mean

logical nitrogen fixation (Oram et al. 2014). When applying biochar or BF, the co-cropping mode of tomato and leguminous green manure may reduce the amount of nitrogen application (Gu et al. 2021) and improve the biological availability of nitrogen (Gatsios et al. 2021).

**Effects of biochar-based fertiliser on the fertiliser utilisation of tomato.** The fertiliser utilisation rate is an important indicator for evaluating the absorption and utilisation of fertiliser nutrients by crops, which is closely related to factors that affect nutrient conversions such as soil aeration status, soil

water content, and microbial activity (Peregrina et al. 2020). Compared with TFP treatment, the AE of N, P and K treated with the BF treatments increased by 20–71, 202–329, and 27–42% respectively in 2019, and 31–87, 229–365, and 10–56% respectively in 2020 (Table 3). The AE of N, P and K in BF1 treatment were both the highest, with 7.7, 19.3, and 6.4 kg/kg in 2019, and 9.7, 24.2, and 8.1 kg/kg respectively in 2020. The RE of N, P and K treated with the BF treatments increased by 92–138, 336–448, and 54–107% respectively in 2019, and 68–96, 318–417, and 78–111% respectively in 2020. The RE of N, P and

Table 3. Effects of different fertilisation on agronomic efficiency (AE) and recovery efficiency (RE)

Year	Treatment	AE (kg/kg)			RE (%)		
		N	P	K	N	P	K
2019	CK	–	–	–	–	–	–
	TFP	4.5 ± 0.8 <sup>b</sup>	4.5 ± 0.8 <sup>c</sup>	4.5 ± 0.8 <sup>b</sup>	15.6 ± 3.3 <sup>c</sup>	5.0 ± 0.3 <sup>d</sup>	23.2 ± 4.6 <sup>d</sup>
	BF1	7.7 ± 0.5 <sup>a</sup>	19.3 ± 1.3 <sup>a</sup>	6.4 ± 0.4 <sup>a</sup>	34.1 ± 3.9 <sup>ab</sup>	23.2 ± 1.5 <sup>bc</sup>	35.7 ± 6.2 <sup>c</sup>
	BF2	7.0 ± 0.8 <sup>a</sup>	17.5 ± 2.1 <sup>a</sup>	5.8 ± 0.7 <sup>a</sup>	35.3 ± 4.3 <sup>ab</sup>	24.9 ± 1.0 <sup>b</sup>	44.7 ± 5.3 <sup>ab</sup>
	BF3	6.9 ± 0.9 <sup>a</sup>	17.2 ± 2.3 <sup>a</sup>	5.7 ± 0.8 <sup>ab</sup>	37.1 ± 3.1 <sup>a</sup>	27.4 ± 1.1 <sup>a</sup>	48.1 ± 3.8 <sup>a</sup>
	BF4	5.4 ± 0.8 <sup>b</sup>	13.6 ± 2.0 <sup>b</sup>	4.5 ± 0.7 <sup>b</sup>	30.0 ± 2.5 <sup>b</sup>	21.8 ± 0.9 <sup>c</sup>	39.0 ± 2.9 <sup>bc</sup>
2020	CK	–	–	–	–	–	–
	TFP	5.2 ± 0.2 <sup>e</sup>	5.2 ± 0.2 <sup>e</sup>	5.2 ± 0.2 <sup>d</sup>	20.6 ± 0.9 <sup>d</sup>	5.6 ± 0.3 <sup>d</sup>	24.8 ± 1.3 <sup>c</sup>
	BF1	9.7 ± 0.6 <sup>a</sup>	24.2 ± 1.5 <sup>a</sup>	8.1 ± 0.5 <sup>a</sup>	37.3 ± 2.2 <sup>bc</sup>	26.5 ± 1.8 <sup>b</sup>	45.6 ± 2.9 <sup>b</sup>
	BF2	8.6 ± 0.3 <sup>b</sup>	21.6 ± 0.6 <sup>b</sup>	7.2 ± 0.2 <sup>b</sup>	38.8 ± 1.1 <sup>ab</sup>	27.0 ± 0.9 <sup>ab</sup>	49.6 ± 1.5 <sup>a</sup>
	BF3	7.9 ± 0.2 <sup>c</sup>	19.8 ± 0.6 <sup>c</sup>	6.6 ± 0.2 <sup>c</sup>	40.3 ± 1.4 <sup>a</sup>	28.9 ± 0.9 <sup>a</sup>	52.3 ± 2.1 <sup>a</sup>
	BF4	6.8 ± 0.5 <sup>d</sup>	17.1 ± 1.3 <sup>d</sup>	5.7 ± 0.4 <sup>d</sup>	34.5 ± 2.0 <sup>c</sup>	23.4 ± 1.6 <sup>c</sup>	44.2 ± 2.8 <sup>b</sup>

N – nitrogen; P – phosphorus; K – potassium; CK – no fertiliser; TFP – traditional fertilisation practice; BF1 – 1 875 kg/ha of biochar-based fertiliser (BF); BF2 – 2 250 kg/ha of BF; BF3 – 2 625 kg/ha of BF; BF4 – 3 000 kg/ha of BF. Different letters in the same column indicates significant difference between different fertilisation treatments ( $P < 0.05$ )

K in BF3 treatment were both the highest, with 37, 27, and 48% in 2019, and 40, 29 and 52% respectively in 2020. It can be seen that BF could improve the utilisation rate of fertiliser and reduce the loss of fertiliser nutrients.

Due to the presence of exchangeable ions and functional groups (Puga et al. 2019b), the biochar in biochar-based fertilisers can adsorb nitrogen ( $\text{NH}_4^+$  or  $\text{NO}_3^-$ ), which can avoid nitrogen leaching losses and gaseous losses, and thus improve nitrogen utilisation (Dong et al. 2019). Moreover, BF can not only change the physicochemical properties of soil, but also increase the enzyme activity related to soil nitrogen conversion, the abundance and activity of ammonia oxidising bacteria, and thereby increasing the biological availability of nitrogen (Liao et al. 2020). Meanwhile, the potassium content of biochar was relatively higher (Zornoza et al. 2016). In this experiment, the total potassium content of biochar was 21.4 g/kg, and the potassium in the biochar was mostly soluble (Tan et al. 2017). Therefore, the content of available potassium in the soil can increase quickly as well as be absorbed and used by crops in time after biochar is applied (Gwenzi et al. 2017). On the other hand, the temperature and pH of the soil as well as the content of organic carbon can be increased using biochar. This may indirectly increase the number and activity of solubilising silicate bacteria, which is conducive to the improve-

ment of soil potassium fixation and utilisation (Gao et al. 2021). Therefore, using potassium in biochar to replace potassium fertiliser may be a practical method, which is of great significance to alleviate the lack of potassium resources in China and improve the utilisation of straw potash resources.

It is worth noting that the prerequisite for the large-scale application of biochar was the economic feasibility of biochar application (Aguirre et al. 2021), however, most of the current research still focuses on the environmental effects of biochar rather than the economic benefits (Owsianiak et al. 2021). The main reason was the excessively high input costs of biochar and relatively slow benefits due to a large amount of biochar applied and the high price. Therefore, it is necessary to reduce the cost of raw materials as much as possible in the future, and thus reduce the price of BF, so that it can be widely used to maximise farmers' economic benefits.

Consequently, replacing the traditional mineral fertiliser with BF has been proved to be an alternative solution for managing sustainable fertilisation on open field tomato in karst mountainous areas, and the application of 2 326 kg/ha BF is worthy of popularising. Moreover, many reports have shown the positive effects of BF on crop production, the research on the mechanism of BF is still lacking. Further research should focus on studying the effects of BF on soil quality and microecology, to clarify the ecological and environmental effects.



<https://doi.org/10.17221/471/2021-PSE>

## REFERENCES

- Aguirre J.L., Martín M.T., González S., Peinado M. (2021): Effects and economic sustainability of biochar application on corn production in a mediterranean climate. *Molecules*, 26: 3313.
- Bach M., Wilske B., Breuer L. (2016): Current economic obstacles to biochar use in agriculture and climate change mitigation. *Carbon Management*, 7: 183–190.
- Bao S.D. (2000): *Soil Agricultural – Chemical Analysis*. Beijing, China Agricultural Press. (In Chinese)
- Chew J.K., Zhu L.L., Nielsen S., Graber E., Mitchell D.R.G., Horvat J., Mohammed M., Liu M.L., van Zwieten L., Donne S., Munroe P., Taherymoosavi S., Pace B., Rawal A., Hook J., Marjo C., Thomas D.S., Pan G.X., Li L.Q., Bian R.J., McBeath A., Bird M., Thomas T., Husson O., Solaiman Z., Joseph S., Fan X.R. (2020): Biochar-based fertilizer: supercharging root membrane potential and biomass yield of rice. *Science of The Total Environment*, 713: 136431.
- Das S.K., Ghosh G.K. (2021): Development and evaluation of biochar-based secondary and micronutrient enriched slow release nano-fertilizer for reduced nutrient losses. *Biomass Conversion and Biorefinery*, 2021.
- Dong D., Wang C., van Zwieten L., Wang H.L., Jiang P.K., Zhou M.M., Wu W.X. (2019): An effective biochar-based slow-release fertilizer for reducing nitrogen loss in paddy fields. *Journal of Soils and Sediments*, 20: 3027–3040.
- Dong D., Li J., Ying S.S., Wu J.S., Han X.G., Teng Y.X., Zhou M.R., Ren Y., Jiang P.K. (2021): Mitigation of methane emission in a rice paddy field amended with biochar-based slow-release fertilizer. *Science of The Total Environment*, 792: 148460.
- Dumroese R.K., Heiskanen J., Englund K., Tervahauta A. (2011): Pelleted biochar: chemical and physical properties show potential use as a substrate in container nurseries. *Biomass and Bioenergy*, 35: 2018–2027.
- Gao M.Y., Yang J.F., Liu C.M., Gu B.W., Han M., Li J.W., Li N., Liu N., An N., Dai J., Liu X.H., Han X.R. (2021): Effects of long-term biochar and biochar-based fertilizer application on brown earth soil bacterial communities. *Agriculture, Ecosystems and Environment*, 309: 107285.
- Gatsios A., Ntatsi G., Celi L., Said-Pullicino D., Tampakaki A., Savaas D. (2021): Legume-based mobile green manure can increase soil nitrogen availability and yield of organic greenhouse tomatoes. *Plants (Basel)*, 10: 2419.
- Gu C.M., Huang W., Li Y., Li Y.S., Yu C.B., Dai J., Hu W.S., Li X.Y., Brooks M., Xie L.H., Liao X., Qin L. (2021): Green manure amendment can reduce nitrogen fertilizer application rates for oilseed rape in maize-oilseed rape rotation. *Plants (Basel)*, 10: 2640.
- Gwenzi W., Nyambishi T.J., Chaukura N., Mapope N. (2017): Synthesis and nutrient release patterns of a biochar-based N-P-K slow-release fertilizer. *International Journal of Environmental Science and Technology*, 15: 405–414.
- Hong J.L., Ren L.J., Hong J.M., Xu C.Q. (2016): Environmental impact assessment of corn straw utilization in China. *Journal of Cleaner Production*, 112: 1700–1708.
- Ibrahim M.M., Tong C.X., Hu K., Zhou B.Q., Xing S.H., Mao Y.L. (2020): Biochar-fertilizer interaction modifies N-sorption, enzyme activities and microbial functional abundance regulating nitrogen retention in rhizosphere soil. *Science of The Total Environment*, 739: 140065.
- Joseph S., Graber E.R., Chia C., Munroe P., Donne S., Thomas T., Nielsen S., Marjo C., Rutledge H., Pan G.X., Li L., Taylor P., Rawal A., Hook J. (2013): Shifting paradigms: development of high-efficiency biochar fertilizers based on nano-structures and soluble components. *Carbon Management*, 4: 323–343.
- Joseph S., Anawar H.M., Storer P., Chia C., Lin Y., Munroe P., Donne S., Horvat J., Wang J.L., Solaiman Z.M. (2015): Effects of enriched biochars containing magnetic iron nanoparticles on mycorrhizal colonisation, plant growth, nutrient uptake and soil quality improvement. *Pedosphere*, 25: 749–760.
- Khajavi-Shojaei S., Moezzi A., Masir M.N., Taghavi M. (2020): Synthesis modified biochar-based slow-release nitrogen fertilizer increases nitrogen use efficiency and corn (*Zea mays* L.) growth. *Biomass Conversion and Biorefinery*, 2020.
- Kong W.W., Zhang M., Liu Y.L., Gou J.L., Wei Q.Q., Shen B.X. (2021): Physico-chemical characteristics and the adsorption of ammonium of biochar pyrolyzed from distilled spirit lees, tobacco fine and Chinese medicine residues. *Journal of Analytical and Applied Pyrolysis*, 156: 105148.
- Kumar A., Singh E., Mishra R., Kumar S. (2022): Biochar as environmental armour and its diverse role towards protecting soil, water and air. *Science of The Total Environment*, 806: 150444.
- Liang B., Lehmann J., Solomon D., Kinyangi J., Grossman J., O'Neill B., Skjemstad J.O., Thies J., Luizao F.J., Petersen J., Neves E.G. (2006): Black carbon increases cation exchange capacity in soils. *Soil Science Society of America Journal*, 70: 1719–1730.
- Liao J.Y., Liu X.R., Hu A., Song H.X., Chen X.Z., Zhang Z.H. (2020): Effects of biochar-based controlled release nitrogen fertilizer on nitrogen-use efficiency of oilseed rape (*Brassica napus* L.). *Scientific Reports*, 10: 11063.
- Liu Z.Y., Demisie W., Zhang M.K. (2013): Simulated degradation of biochar and its potential environmental implications. *Environmental Pollution*, 179: 146–152.
- Liu C.C., Liu Y.G., Guo K., Wang S.J., Liu H.M., Zhao H.W., Qiao X.G., Hou D.J., Li S.B. (2016): Aboveground carbon stock, allocation and sequestration potential during vegetation recovery in the karst region of southwestern China: a case study at a watershed scale. *Agriculture, Ecosystems and Environment*, 235: 91–100.
- Liu X.Y., Wang H.D., Liu C., Sun B.B., Zheng J.F., Bian R.J., Drosos M., Zhang X.H., Li L.Q., Pan G.X. (2020): Biochar increases maize yield by promoting root growth in the rainfed region. *Archives of Agronomy and Soil Science*, 67: 1411–1424.
- Luo X.X., Chen L., Zheng H., Chang J.J., Wang H.F., Wang Z.Y., Xing B.S. (2016): Biochar addition reduced net N mineralization of a coastal wetland soil in the Yellow River Delta, China. *Geoderma*, 282: 120–128.
- Maroušek J., Trakal L. (2021): Techno-economic analysis reveals the untapped potential of wood biochar. *Chemosphere*, 291: 133000.

<https://doi.org/10.17221/471/2021-PSE>

- Melo L.C.A., Lehmann J., Carneiro J.S. da S., Camps-Arbestain M. (2022): Biochar-based fertilizer effects on crop productivity: a meta-analysis. *Plant and Soil*, 2022: 486.
- Olsen S.R., Cole C.V., Watanabe F.S., Dean L.A. (1954): Estimation of available phosphorus in soil by extraction with sodium bicarbonate. Washington, United States Department of Agriculture.
- Oram N.J., van de Voorde T.F.J., Ouwehand G.-J., Bezemer T.M., Mommer L., Jeffery S., Groenigen J.W.V. (2014): Soil amendment with biochar increases the competitive ability of legumes *via* increased potassium availability. *Agriculture, Ecosystems and Environment*, 191: 92–98.
- Owsianiak M., Lindhjem H., Cornelissen G., Hale S.E., Sørmo E., Sparrevik M. (2021): Environmental and economic impacts of biochar production and agricultural use in six developing and middle-income countries. *Science of The Total Environment*, 755: 142455.
- Pan Q.L., Chen Q., Song T., Xu X.N., Zhan X.M., Peng J., Su H.Q., Wang Y., Han X.R. (2017): Influences of biochar and biochar-based compound fertilizer on soil water retention in brown soil. *Research of Soil and Water Conservation*, 24: 115–121. (In Chinese)
- Peregrina P.A., Grutzmacher P., Eduardo P.C.C., Sanches R.V., Alberto D.A.C. (2020): Biochar-based nitrogen fertilizers: greenhouse gas emissions, use efficiency, and maize yield in tropical soils. *Science of The Total Environment*, 704: 135375.
- Pituello C., Dal Ferro N., Francioso O., Simonetti G., Berti A., Piccoli I., Pisi A., Morari F. (2018): Effects of biochar on the dynamics of aggregate stability in clay and sandy loam soils. *European Journal of Soil Science*, 69: 827–842.
- Puga A.P., Grutzmacher P., Cerri C.E.P., Ribeirinho V.S., Andrade C.A. (2019a): Biochar-based nitrogen fertilizers: greenhouse gas emissions, use efficiency, and maize yield in tropical soils. *Science of The Total Environment*, 704: 135375.
- Puga A.P., Queiroz M.C.de.A., Ligo M.A.V., Carvalho C.S., Pires A.M.M., Marcatto J.de.O.S., Andrade C.A. (2019b): Nitrogen availability and ammonia volatilization in biochar-based fertilizers. *Archives of Agronomy and Soil Science*, 66: 992–1004.
- Qian L., Chen L., Joseph S., Pan G.X., Li L.Q., Zheng J.W., Zhang X.H., Zheng J.F., Yu X.Y., Wang J.F. (2014): Biochar compound fertilizer as an option to reach high productivity but low carbon intensity in rice agriculture of China. *Carbon Management*, 5: 145–154.
- Sim D.H.H., Tan I.A.W., Lim L.L.P., Hameed B.H. (2021): Encapsulated biochar-based sustained release fertilizer for precision agriculture: a review. *Journal of Cleaner Production*, 303: 127018.
- Stávková J., Maroušek J. (2021): Novel sorbent shows promising financial results on P recovery from sludge water. *Chemosphere*, 276: 130097.
- Sun P., Zhao Z.T., Fan P.S., Chen W., Ruan Y.Z., Wang Q. (2021): Ammonia- and nitrite-oxidizing bacteria are dominant in nitrification of maize rhizosphere soil following combined application of biochar and chemical fertilizer. *Frontiers in Microbiology*, 12: 715070.
- Tan Z.X., Liu L.Y., Zhang L.M., Huang Q.Y. (2017): Mechanistic study of the influence of pyrolysis conditions on potassium speciation in biochar "preparation-application" process. *Science of The Total Environment*, 599–600: 207–216.
- Wang Y.F., Xiao X., Zhang K., Chen B.L. (2019): Effects of biochar amendment on the soil silicon cycle in a soil-rice ecosystem. *Environmental Pollution*, 248: 823–833.
- Walter R., Rao B.K.R. (2015): Biochars influence sweet-potato yield and nutrient uptake in tropical Papua New Guinea. *Journal of Plant Nutrition and Soil Science*, 178: 393–400.
- Xu H., Cai A.D., Wu D., Liang G.P., Xiao J., Xu M.G., Colinet G., Zhang W.J. (2021): Effects of biochar application on crop productivity, soil carbon sequestration, and global warming potential controlled by biochar C:N ratio and soil pH: a global meta-analysis. *Soil and Tillage Research*, 213: 105125.
- Yan T.T., Xue J.H., Zhou Z.D., Wu Y.B. (2021a): Effects of biochar-based fertilizer on soil bacterial network structure in a karst mountainous area. *Catena*, 206: 105535.
- Yan T.T., Xue J.H., Zhou Z.D., Wu Y.B. (2021b): Biochar-based fertilizer amendments improve the soil microbial community structure in a karst mountainous area. *Science of The Total Environment*, 794: 148757.
- Yang W.H., Li C.J., Wang S.S., Zhou B.Q., Mao Y.L., Rensing C., Xing S.H. (2021): Influence of biochar and biochar-based fertilizer on yield, quality of tea and microbial community in an acid tea orchard soil. *Applied Soil Ecology*, 166: 104005.
- Yin H.J., Zhao W.Q., Li T., Cheng X.Y., Liu Q. (2018): Balancing straw returning and chemical fertilizers in China: role of straw nutrient resources. *Renewable and Sustainable Energy Reviews*, 81: 2695–2702.
- Zhang Q.Q., Song Y.F., Wu Z., Yan X.Y., Gunina A., Kuzyakov Y., Xiong Z.Q. (2020): Effects of six-year biochar amendment on soil aggregation, crop growth, and nitrogen and phosphorus use efficiencies in a rice-wheat rotation. *Journal of Cleaner Production*, 242: 118435.
- Zhang M., Liu Y.L., Wei Q.Q., Gou J.L. (2021a): Biochar enhances the retention capacity of nitrogen fertilizer and affects the diversity of nitrifying functional microbial communities in karst soil of southwest China. *Ecotoxicology and Environmental Safety*, 226: 112819.
- Zhang M.Y., Zhang L., Riaz M., Xia H., Jiang C.C. (2021b): Biochar amendment improved fruit quality and soil properties and microbial communities at different depths in citrus production. *Journal of Cleaner Production*, 292: 126062.
- Zheng J.L., Wang S.J., Wang R.M., Chen Y.L., Siddique K.H.M., Xia G.M., Chi D.C. (2021): Ameliorative roles of biochar-based fertilizer on morpho-physiological traits, nutrient uptake and yield in peanut (*Arachis hypogaea* L.) under water stress. *Agricultural Water Management*, 257: 107129.
- Zornoza R., Moreno-Barriga F., Acosta J.A., Muñoz M.A., Faz A. (2016): Stability, nutrient availability and hydrophobicity of biochars derived from manure, crop residues, and municipal solid waste for their use as soil amendments. *Chemosphere*, 144: 122–130.

Received: November 6, 2021

Accepted: March 11, 2022

Published online: March 18, 2022