

# Effect of manganese on cadmium toxicity in maize seedlings

P. Paľove-Balang, A. Kisová, J. Pavlovkin, I. Mistrík

*Institute of Botany, Slovak Academy of Sciences, Bratislava, Slovak Republic*

## ABSTRACT

The interaction of manganese with cadmium (Cd) toxicity was studied on maize plants grown in hydroponics. Manganese supplied as  $MnSO_4$  clearly alleviated the toxic effect of cadmium on the root growth of maize seedlings. The magnitude of alleviation was dose dependant and total abolition of  $10\mu M$  Cd toxicity on root growth was observed at Mn/Cd ratio of 20:1. The 12 h pre-treatment with  $10\mu M$  Cd was generally toxic for nitrate uptake and reduction (both determined in Cd-free media). The beneficial effect of  $100\mu M$  Mn on this toxicity was confirmed for the low-affinity nitrate uptake system, but on the other hand, Mn alone seems to be slightly toxic for high affinity nitrate uptake system and on the nitrate reductase activity.

**Keywords:** cadmium; manganese; membrane potential; nitrate uptake; nitrate reduction; maize roots

Cadmium (Cd) is a non-essential and toxic element without any known metabolic significance. The co-existence of Cd with other essential and non-essential heavy metals in the ecosystem can lead to various synergic and antagonistic interactions in their uptake and tissue content. Cadmium addition significantly reduced the Zn and Mn level in roots and shoots of barley (Wu et al. 2003) and durum wheat (Jalil et al. 1994). Similarly, the uptake and toxicity of Cd can be moderated significantly in the presence of excess amounts of certain essential metal nutrients like Zn, Ca, Fe, Cu, Mn (Das et al. 1997, Aravind and Prasad 2003), or non-essential Ti, as was reported recently (Leskó et al. 2002).

Manganese (Mn) is one of the essential heavy metals. The increased level of Mn reduced the uptake of Cd from nutrient medium, which resulted in lower accumulation of Cd in plant tissues (Jarvis et al. 1976, Baszynski et al. 1980). The decrease of plant Cd level after addition of Mn, and vice versa, might be explained by competition for the same membrane-transporters. Recently, due to an effort in sequencing of the whole plant genomes, many multispecific metal transporters have been identified and described. The NRAMP and IRT1 genes encode integral membrane proteins that can transport Mn as well as Cd among other metals (Rogers et al. 2000, Thomine et al. 2003, Cohen et al. 2004). The tobacco plants overexpressing the *Arabidopsis* CAX2 gene for Ca transporter,

accumulate significantly higher levels of Mn and Cd in tonoplast (Hirschi et al. 2000). The high Mn level in white lupine (*Lupinus albus* L.), a Mn hyperaccumulator, could contribute to the alleviation of negative effects of Cd, especially on the photosynthesis (Zornoza et al. 2002). A higher Mn accumulation in chloroplast when Cd is present in the growth medium was also found in lettuce (*Lactuca sativa* L., Ramos et al. 2002). Moreover, the partial restoration of chloroplast structure damaged by Cd treatment was found after addition of Mn (Baszynski et al. 1980).

The nitrogen nutrition is one of the most important metabolic processes in plants. Several authors found that nitrate uptake and assimilation is negatively affected by the presence of Cd in the growing medium. In particular, nitrate uptake decreased after Cd treatment in pea (Hernández et al. 1997), wheat (Stolt and Oscarson 2002) or barley (Boussana et al. 1999). Similarly, Cd inhibited the activity of nitrate reductase and nitrite reductase, the enzymes involved in nitrate reduction (Chugh et al. 1992, Ouariti et al. 1997) as well as glutamine synthetase, the enzyme involved in nitrogen assimilation (Chiraz et al. 2003).

The aim of the present study was to investigate the alternations in the growth, nitrate uptake and reduction in maize plants grown in the presence of different combinations of cadmium and manganese in culture medium. The possible positive effect of Mn on Cd toxicity is discussed.

## MATERIAL AND METHODS

**Plant material and growth conditions.** Maize (*Zea mays* L. cv. Tina) seeds were placed in filter papers rolls soaked with distilled water and germinated in the dark at 25°C and 98% RH. Three-day-old seedlings with 5–6 cm long primary seminal roots were used for growth experiments (see below), or transferred into N-free, half strength Hoagland's solution and kept in a growth chamber E8 (Convion, Canada) under controlled conditions (24°C, 70% RH, 14/10 h light/dark cycle and photosynthetic quantum flux density 180  $\mu\text{mol}/\text{m}^2/\text{s}$ ). Nutrient solution was continuously aerated and changed 2 times a week for a fresh one. 14-day-old seedlings were used for uptake, nitrogen content and nitrate reductase activity measurements after 12 h pre-treatment with induction solution containing 0.2mM (for uptake) or 2mM (for nitrate reductase activity)  $\text{KNO}_3$ , 0.5mM  $\text{CaSO}_4$  (control); and 10 $\mu\text{M}$   $\text{CdCl}_2$  with different concentrations of  $\text{MnSO}_4$  (0–1mM), (Cd and Mn treated plants). For root growth determination, three-day-old seedlings were selected for uniformity and transferred to glass pots containing 0.5mM  $\text{CaSO}_4$  and different concentrations of Cd (0–1mM), supplied as  $\text{CdCl}_2$  (pH 5.0). To screen the effect of Mn on root growth under Cd stress, the seedlings were treated by 10 $\mu\text{M}$  Cd, in combination with 1 $\mu\text{M}$  to 1mM  $\text{MnSO}_4$ . All seedlings were left in dark at 24°C. The growth rate of roots was calculated as an increase of root length between final and initial root lengths and expressed in cm/day.

**Nitrate uptake and nitrate content.** Nitrate uptake was calculated by following depletion from the uptake solution containing 0.5mM  $\text{CaSO}_4$  and 0.2 or 2mM  $\text{KNO}_3$  (for high and low affinity uptake systems, respectively). Concentrations of  $\text{NO}_3^-$  were determined spectrophotometrically (Braun-Systematic, Methodenblatt N60) and related to the root fresh weight. The minimal concentration of the solution was 0.05mM for high affinity uptake conditions and 1mM for low affinity uptake conditions at the end of experiment. Determinations of nitrate content in root crude extracts from 4 cm long apical segments were performed as described by Pajuelo et al. (2002). 200  $\mu\text{l}$  of 5% (w/v) salicylic acid dissolved in 96% (w/v) sulphuric acid was added to aliquots of 50  $\mu\text{l}$  from the root crude extracts (5 ml/g FW) and left to react for 20 min. 4.75 ml of 2M NaOH was added to the reaction mixtures and then the absorbance was read at 405 nm after cooling. A calibration curve of known amounts of nitrate

dissolved in the standard extraction buffer was used for analytical determinations. Blanks were set up without salicylic acid.

**Cadmium and manganese content.** The cadmium and manganese content was determined in 5 cm long root segment of the maize seedling treated the same way as for nitrate uptake measurements. The dry roots were digested in teflon autoclave in nitric acid for 4 h at 160°C, and dried in fluoric acid. The dry matter was digested in nitric acid. The Cd and Mn content were determined by atomic absorption spectrometry at 228.8 nm for Cd and 279.5 nm for Mn.

**Measurement of membrane potential.** Three-day-old maize seedlings after 24 h pre-incubation in solution containing 1mM  $\text{CaSO}_4$ , 0.2mM  $\text{KNO}_3$  and combinations of 10 $\mu\text{M}$   $\text{CdCl}_2$  and 10–200 $\mu\text{M}$   $\text{MnSO}_4$ , respectively, were attached to a Plexiglas holder and mounted in a vertical 5  $\text{cm}^3$  cuvette perfused with solution containing 1mM  $\text{CaSO}_4$  (pH 5.5). Nitrate was added by replacing the 1mM  $\text{CaSO}_4$  by 1mM  $\text{Ca}(\text{NO}_3)_2$ . Measurements of the membrane potential ( $E_m$ ) were carried out at 22°C, using standard microelectrode techniques (Pavlovkin et al. 1993) in the outermost layer of the root cortex cells 3–5 mm behind the root cap.

**Nitrate reductase assay.** For nitrate reductase activity, 4 cm tips or basis of plant roots were quickly frozen in liquid nitrogen, homogenised and then kept at  $-80^\circ\text{C}$  until use. Root material (0.1 g) was ground in a mortar at 4°C and resuspended in extraction buffer (0.6 ml). Extraction buffer contained: 100mM HEPES-KOH (pH 7.6), 10mM EDTA, 1mM dithiothreitol (DTT), 1mM phenylmethylsulphonyl fluoride (PMSF), 3% (w/v) bovine serum albumin (BSA), and 0.1% (v/v) Triton X-100. The homogenate was centrifuged for 15 min at 15 000  $\times g$  and insoluble material was removed. The clear supernatants were immediately assayed for NR activity (Pajuelo et al. 2002). The  $\text{NRA}_{\text{tot}}$  assay was carried out as follows: 500  $\mu\text{l}$  final volume reaction mixtures were placed in 1.5 ml Eppendorf tubes containing: 250  $\mu\text{l}$  of assay buffer (100mM HEPES-KOH pH 7.6, 10mM EDTA and 20 $\mu\text{M}$  FAD), 50  $\mu\text{l}$  100mM  $\text{KNO}_3$ ; 50  $\mu\text{l}$  of extraction buffer and 100  $\mu\text{l}$  root crude extract. The reaction was started by addition of 50  $\mu\text{l}$  0.7mM NADH and carried out at 30°C for 30 min. The  $\text{NRA}_{\text{act}}$  determination was performed by the same way, except that 10mM EDTA was replaced by 10mM  $\text{MgCl}_2$  both in extraction and assay buffer. Nitrite formation was measured as described elsewhere (De la Rosa et al. 1982), using sulphanilamide and N-(1-naphtyl)-ethylenediamine dihydrochloride reagents. Blank

tubes were processed as samples without NADH addition.

## RESULTS

Exposure of maize seedling to 10  $\mu\text{M}$  Cd for 48 h resulted in almost 50% reduction of root growth. In contrast, Cd-treated plants supplemented with Mn showed lower Cd-induced root growth inhibition, magnitude of which was Mn-dose dependent (Figure 1). Application of 200  $\mu\text{M}$  Mn to root solution completely alleviated the toxicity of 10  $\mu\text{M}$  Cd on the root growth. The addition of 10, 100 or 200  $\mu\text{M}$  Mn alone, without Cd had no significant effect on root growth (data not shown).

The net  $\text{NO}_3^-$  uptake, both at low and high affinity uptake conditions was also reduced by 12 h pre-treatment with 10  $\mu\text{M}$  Cd. Magnitude of inhibition was similar for both systems and represented 26.3% for HATS and 28.9% for LATS. Pre-treatment with 100  $\mu\text{M}$  Mn was found to be slightly inhibitory for high affinity  $\text{NO}_3^-$  uptake (18.4% inhibition), but not for low affinity one (Figure 2). Furthermore, Mn slightly, but not significantly ( $P = 0.14$ ) alle-

viated Cd-induced inhibition of  $\text{NO}_3^-$  uptake by LATS, but enhanced inhibition of  $\text{NO}_3^-$  uptake by HATS. Accumulation of  $\text{NO}_3^-$  in root tip tissues declined not only in the presence of Cd, but slightly also after Mn treatment, whereas Mn in combination with Cd did not show any improvement on nitrate accumulation (Figure 3).

Pre-treatment of maize roots with Cd showed a slight increase of the rest membrane potential value ( $E_m$ ) in contrary to Mn, pre-treatment with which caused a slight decrease of membrane potential (measured in the 1 mM  $\text{CaSO}_4$  solution, without Cd or Mn). After pre-treatment with both ions, the antagonistic effect of Mn to Cd on the membrane potential was observed. This effect was concentration dependent, and more-less equilibrated at 10:1 (Mn:Cd) stoichiometry (Table 1). Replacement of 1 mM  $\text{CaSO}_4$  by 1 mM  $\text{Ca}(\text{NO}_3)_2$  resulted in gradual hyperpolarization of membrane potential ( $\Delta = 33.2$  mV) (Figure 4). The initial electrical response to 2 mM nitrate was significantly lower in plants pre-treated with Cd, pre-treatment with 100 and 200  $\mu\text{M}$  Mn alleviated the inhibitory effect of Cd.

The cadmium content in roots treated with Mn + Cd mixture was nearly the same as in the




Figure 1. Effect of Cd and Mn ions on root growth of 3-day-old maize seedlings; root solution contained 0.5  $\mu\text{M}$   $\text{CaSO}_4$  (control – open column) supplemented with 10  $\mu\text{M}$   $\text{CdCl}_2$  and different concentrations of  $\text{MnCl}_2$  (0–200  $\mu\text{M}$  – filled columns); the root growth increments are expressed in cm per day; results are the means ( $\pm$  SE) of 4 experiments ( $n = 20$ )




Figure 2. Nitrate uptake by roots of 14-day-old maize plants; twelve hours before the uptake measurement plants were transferred to solution containing 0.5 mM  $\text{CaSO}_4$ , 0.2 mM  $\text{KNO}_3$ , 10  $\mu\text{M}$  Cd and/or 100  $\mu\text{M}$  Mn; the uptake solution was Cd and Mn-free and contained 0.5 mM  $\text{CaSO}_4$  and 0.2 mM  $\text{KNO}_3$  for HATS or 2 mM  $\text{KNO}_3$  for LATS; results are the means ( $\pm$  SE) of 4 experiments ( $n = 3$ )




Figure 3. Nitrate content in 4 cm long apical segments of maize roots after 12 h pre-treatment with 10 $\mu$ M Cd (Cd), with 100 $\mu$ M Mn (Mn) or with a mixture of 10 $\mu$ M Cd + 100 $\mu$ M Mn (Mn + Cd); results are the means ( $\pm$  SE) of 4 experiments ( $n = 3$ )

plants treated with Cd alone, despite the 10 times higher concentration of Mn. On the other hand, the relatively low cadmium concentration in Mn + Cd media, led to the much stronger reduction (12–36%) of the manganese level in maize root tissue in comparison with the plants treated with Mn alone (Table 2).

Nitrate reductase activity (NRA) significantly decreased in roots treated by 10 $\mu$ M Cd. The extent of inhibition reached 28% in case of the total nitrate reductase activity (NRA<sub>tot</sub>) and 39% in case of the

actual nitrate reductase activity (NRA<sub>act</sub>) (Figure 5). The same effect was observed in roots pre-treated with 100 $\mu$ M Mn. The extent of inhibition was very similar and reached 20 and 30% of both NRA<sub>tot</sub> and NRA<sub>act</sub>, respectively. The common effect of Mn + Cd was slightly synergic, resulting in 54% inhibition of NRA<sub>act</sub> and only 41% activity state of NRA (NRA<sub>act</sub>/NRA<sub>tot</sub>) in comparison to 63% activity state in control. The total protein content in root tissue remained unchanged during the NRA experiments.

## DISCUSSION

Our experiments significantly confirmed that sufficient amount of Mn may fully ameliorate the Cd-induced root growth inhibition in maize seedlings. At the 1:1 stoichiometry, which was in our conditions 10 $\mu$ M Mn and 10 $\mu$ M Cd, only weak alleviation effect of Mn was observed. The magnitude of Cd-toxicity alleviation increased with elevated Mn concentration and in the presence of 200 $\mu$ M Mn, when the Mn/Cd ratio reached 20:1, the root growth was approximately the same as in control plants. Such high Mn/Cd ratio could be usual in natural conditions, because natural Mn content in soils can reach mg/g range, even if the Mn availability for plants is partially reduced by Cd enrichment of the soil (Ramachandran and Souza 2002).

In spite of the fact that the total amelioration of Cd-induced root growth inhibition was observed after application of 200 $\mu$ M Mn concentration, further experiments (nitrate uptake and assimilation) were performed only with 100 $\mu$ M Mn. The main reason was that 200 $\mu$ M Mn alone negatively affected nitrate uptake and nitrate assimilation. Even a reduced concentration of Mn to 100 $\mu$ M

Table 1. Resting potential (Em) and hyperpolarization of membrane potential ( $\Delta$ ) during NO<sub>3</sub><sup>-</sup> uptake in intact roots of maize seedlings pre-treated for 12 h with different concentrations of Cd and Mn

Treatment	Em ( $\pm$ SE)	$\Delta$ ( $\pm$ SE)
Control	-131.8 ( $\pm$ 1.0) c	-33.2 ( $\pm$ 1.2) c
100 $\mu$ M Mn	-126.9 ( $\pm$ 0.8) b	-29.1 ( $\pm$ 0.8) b
10 $\mu$ M Cd	-141.8 ( $\pm$ 1.0) e	-20.0 ( $\pm$ 0.5) a
10 $\mu$ M Cd + 10 $\mu$ M Mn	-142.6 ( $\pm$ 1.2) e	-18.9 ( $\pm$ 1.2) a
10 $\mu$ M Cd + 100 $\mu$ M Mn	-136.9 ( $\pm$ 1.2) d	-26.8 ( $\pm$ 1.2) b
10 $\mu$ M Cd + 200 $\mu$ M Mn	-122.9 ( $\pm$ 0.8) a	-26.1 ( $\pm$ 1.2) b

Significancy ( $P_{0.05}$ ) is marked by letters

Table 2. Content of Mn and Cd in 4 cm root tips of maize seedlings in two separate experiments

Treatment	Cd content ( $\mu\text{g/g DW}$ )		Mn content ( $\mu\text{g/g DW}$ )	
	exp. 1	exp. 2	exp. 1	exp. 2
Control	4.2	7.6	4.4	4.2
Mn	n.d.	n.d.	411	419
Cd	370	323	n.d.	n.d.
Mn + Cd	312	307	263	326
% Mn + Cd/Cd	84.3%	95.0%	–	–
% Mn + Cd/Mn	–	–	64.0%	77.8%

n.d. = non detected; % Mn + Cd/Cd = the percentage of Cd content in roots treated with Mn + Cd to the Cd content in roots treated with Cd alone; % Mn + Cd/Mn = the percentage of Mn content in roots treated with Mn + Cd to the Mn content in roots treated with Mn alone

caused a significant inhibition of nitrate reductase activity.

Nitrate uptake in higher plants is mediated by two kinetically distinct systems. High-affinity transport system (HATS) works at low external  $\text{NO}_3^-$  concentrations, and low-affinity transport system (LATS), which is involved in nitrate uptake when external concentration of nitrate is higher than 1mM. Therefore, the uptake measurements at 0.2mM external  $\text{NO}_3^-$  represent the high affinity uptake conditions, and 2mM  $\text{NO}_3^-$  the low affinity

uptake conditions. Both systems, HATS as well as LATS were significantly affected by Cd-treatment. In our experiments with maize, 12 h pre-treatment with 10 $\mu\text{M}$  Cd resulted in 28.5% reduction of nitrate uptake by LATS and 26.3% reduction by HATS. On the other hand, pre-treatment with both ions (Cd + Mn in ratio 1:10) almost doubled the inhibitory effect on HATS (39.4%) while partially but not significantly alleviated Cd-induced inhibition of LATS (21.7%). A similar reduction of nitrate uptake was demonstrated for wheat (Stolt and




Figure 4. Representative changes of membrane potential of maize outer cortical root cells treated with 1mM  $\text{CaSO}_4$  (a); 10 $\mu\text{M}$  Cd + 100 $\mu\text{M}$  Mn (b); 100 $\mu\text{M}$  Mn (c) or 10 $\mu\text{M}$  Cd (d); arrows indicate addition of 2mM  $\text{Ca}(\text{NO}_3)_2$




Figure 5. Total ( $\text{NRA}_{\text{tot}}$ ) and actual activity ( $\text{NRA}_{\text{act}}$ ) of nitrate reductase measured *in vitro* in 4 cm apical segments of maize roots pre-treated with 10 $\mu\text{M}$  Cd, 100 $\mu\text{M}$  Mn, or 10 $\mu\text{M}$  Cd + 100 $\mu\text{M}$  Mn; results are the means of 4 experiments ( $\pm SE$ ,  $n = 3$ )

Oscarson 2002) and pea (Hernández et al. 1997). Boussama et al. (1999) firstly observed depression of the  $\text{NO}_3^-$  uptake after 12 h Cd pre-treatment during the  $\text{NO}_3^-$  induction period in barley, however, in this case 1mM  $\text{NO}_3^-$  was used for uptake measurements, which makes impossible to assign it to particular transport systems. This seems to be important for the effect of 100 $\mu\text{M}$  Mn, which was found to be slightly toxic for the HATS (18.2% reduction), but not for the LATS. Furthermore, a very slight alleviating effect of Mn was observed for LATS. The observed uptake at 2mM  $\text{NO}_3^-$  could be in fact a superposition of uptake provided by the LATS and HATS, and therefore alleviation effect of Mn for LATS could be partially masked by its slight toxicity for HATS.

Analysis of Mn and Cd content only partially confirmed the “competition theory” between Cd and Mn uptake (Rogers et al. 2000, Thomine et al. 2003, Cohen et al. 2004). In our experiments, plants treated for 12 h with solution contained 100 $\mu\text{M}$  Mn + 10 $\mu\text{M}$  Cd had lower Cd and Mn content than plants treated with Mn, or Cd alone as was reported by other authors (Jarvis et al. 1976, Baszyński et al. 1980, Wu et al. 2003). On the other hand, Mn-induced decrease of Cd content very low, and cannot explain the observed effects of Cd and Mn on nitrate uptake and assimilation.

To confirm the partially positive effect of Mn on Cd induced inhibition of LATS simultaneous electrophysiological measurements were performed on intact maize roots. Our electrophysiological measurements agree well with data presented by McClure et al. (1990) and Thibaud and Grignon (1981) working with intact maize roots. According to these authors, 2mM nitrate hyperpolarizes membrane potential to a value more negative than the original resting potential. Our results describe similar hyperpolarization and support a model for nitrate-dependent membrane potential hyperpolarization and nitrate uptake in co-transport with  $\text{H}^+$ . The magnitude of hyperpolarization was smaller in roots grown in presence of cadmium, and this effect was significantly alleviated in the presence of Mn, but only when Mn concentration was at least 10 times higher than Cd concentration. This observation is in good agreement with the results obtained by measurement of  $\text{NO}_3^-$  uptake upon depletion from the solution in low-affinity uptake conditions and supports the idea, that presence of Mn could decrease the toxic effect of Cd on LATS in root tissue.

As we mentioned earlier both Cd (10 $\mu\text{M}$ ) and Mn (100 $\mu\text{M}$ ) negatively affected NR activity. Reduction

of NR activity was reported also in plants grown under Cd-stress (Hernández et al. 1997, Ouariti et al. 1997, Boussama et al. 1999). Our experiments showed that Cd significantly inhibits total ( $\text{NRA}_{\text{tot}}$ ) as well as actual ( $\text{NRA}_{\text{act}}$ ) NRA and this inhibition represents 28.3 and 38.0%, respectively. NR assayed in the presence of EDTA ( $\text{NRA}_{\text{tot}}$ ) is proportional to the steady-state protein concentration. The assay in the presence of  $\text{Mg}^{2+}$  ( $\text{NRA}_{\text{act}}$ ) gives an estimation of the in situ NR activity (Kaiser and Huber 1997). The presence of 100 $\mu\text{M}$  Mn alone decreased both  $\text{NRA}_{\text{tot}}$  and  $\text{NRA}_{\text{act}}$  in similar pattern, but magnitude of reduction was lower than in plants treated with Cd. The effect of both ions was synergic, when added together, to the  $\text{NRA}_{\text{act}}$  but not to the  $\text{NRA}_{\text{tot}}$ . It can be supposed, that both ions partially downregulated the NR on the protein synthesis level, but more strongly interfered with the posttranslational regulation level. The down regulation of  $\text{NRA}_{\text{tot}}$  could be at least partially explained by lower induction caused by lower root  $\text{NO}_3^-$  concentration after Cd treatment. On the other hand Mn, that had no significant effect on tissue  $\text{NO}_3^-$ , also decreased the  $\text{NRA}_{\text{tot}}$ .

Based on our results, we can conclude that an addition of excess amount of Mn to the solution containing phytotoxic concentrations of Cd can improve growth parameters of maize plants. The mechanisms involved in this improvement as well as in the alleviation of toxic effect of Cd on plants are still unclear, and are probably not connected to nitrate metabolism. Slight alleviation of Cd-induced inhibition of LATS by Mn can not overbalance severe toxicity of Cd on total nitrate uptake in short term experiments, but could be beneficial for long term plant growth and development. Therefore future experiments focused on long term effects of Cd and Mn ions on plant metabolisms may bring positive explanation of this problem.

## REFERENCES

- Aravind P., Prasad M.N.V. (2003): Zinc alleviates cadmium-induced oxidative stress in *Ceratophyllum demersum* L.: a free floating freshwater macrophyte. *Plant Physiol. Biochem.*, 41: 391–397.
- Baszyński T., Wajda L., Król M., Wolińska D., Krupa Z., Tukendorf A. (1980): Photosynthetic activities of cadmium-treated tomato plants. *Physiol. Plant.*, 48: 365–370.
- Boussama N., Ouariti O., Ghorbal M.H. (1999): Changes in growth and nitrogen assimilation in barley seedlings under cadmium stress. *J. Plant Nutr.*, 22: 731–752.

- Chiraz Ch., Houda G., Habib G.M. (2003): Nitrogen metabolism in tomato plants under cadmium stress. *J. Plant. Nutr.*, 26: 1617–1634.
- Chungh L.K., Gupta V.K., Sawhney S.K. (1992): Effect of cadmium on enzymes of nitrogen metabolism in pea seedlings. *Phytochemistry*, 31: 395–400.
- Cohen C.K., Garvin D.F., Kochian L.V. (2004): Kinetic properties of a micronutrient transporter from *Pisum sativum* indicate a primary function in Fe uptake from the soil. *Planta*, 218: 784–792.
- Das P., Samantaray S., Rout G.R. (1997): Studies on cadmium toxicity in plants: A review. *Environ. Pollut.*, 98: 29–36.
- De la Rosa M.A., Márquez A.J., Vega J.M. (1982): Dissociation of FAD from NADH-nitrate reductase complex from *Ankistrodesmus braunii* and role of flavin in catalysis. *Z. Naturforsch., C*, 37: 24–30.
- Hernández L.E., Gárate A., Carpena-Ruiz R. (1997): Effects of cadmium on the uptake, distribution and assimilation of nitrate in *Pisum sativum*. *Plant Soil*, 189: 97–106.
- Hirschi K.D., Korenkov V.D., Wilganowski N.L., Wagner G.J. (2000): Expression of Arabidopsis CAX2 in tobacco. Altered metal accumulation and increased manganese tolerance. *Plant Physiol.*, 124: 125–133.
- Jalil A., Selles F., Clarke J.M. (1994): Effect of cadmium on growth and uptake of cadmium and other elements by durum-wheat. *J. Plat Nutr.*, 17: 1839–1858.
- Jarvis S.C., Jones L.H.P., Hopper M.J. (1976): Cadmium uptake from solution by plants and its transport from roots to shoots. *Plant Soil*, 44: 179–191.
- Kaiser W.M., Huber S.C. (1997): Correlation between apparent activation state of nitrate reductase (NR), NR hysteresis and degradation of NR protein. *J. Exp. Bot.*, 48: 1367–1374.
- Leskó K., Stefanovits-Bányai E., Pais I., Simon-Sarkadi L. (2002): Effect of cadmium and titanium-ascorbate stress on biological active compounds in wheat seedlings. *J. Plant Nutr.*, 25: 2571–2581.
- McClure P.R., Kochian L.V., Spanswick R.M., Shaff J.E. (1990): Evidence for cotransport of nitrate and protons in maize roots. I. Effects of nitrate on membrane potential. *Plant Physiol.*, 93: 281–289.
- Ouariti O., Gouia H., Ghorbal M.H. (1997): Responses of bean and tomato plants to cadmium: Growth, mineral nutrition, and nitrate reduction. *Plant Physiol. Biochem.*, 35: 347–354.
- Pajuelo P., Pajuelo E., Orea A., Romero J.M., Márquez A.J. (2002): Influence of plant age and growth conditions on nitrate assimilation in roots of *Lotus japonicus* plants. *Funct. Plant Biol.*, 29: 485–494.
- Pavlovkin J., Mistrík I., Zajchenko A.M., Dugovič L. (1993): Some aspects of phytotoxic action of trichothecene mycotoxin roridin H on corn roots. *Biologia, Bratislava*, 48: 435–439.
- Ramachandran V., D'Souza T.J. (2002): Plant uptake of cadmium, zinc and manganese from four contrasting soils amended with Cd-enriched sewage sludge. *J. Environ. Sci. Health. Toxic. Hazard Subst. Environ. Eng.*, 37: 1337–1346.
- Ramos I., Esteban E., Lucena J.J., Gárate A. (2002): Cadmium uptake and subcellular distribution in plants of *Lactuca* sp. Cd-Mn interaction. *Plant Sci.*, 162: 761–767.
- Rogers E.E., Eide D.J., Guerinot M.L. (2000): Altered selectivity in an Arabidopsis metal transporter. In: *Proc. Nat. Acad. Sci. USA*, 97: 12356–12360.
- Stolt J.P., Oscarson P. (2002): Influence of cadmium on net nitrate uptake kinetics in wheat. *J. Plant Nutr.*, 25: 2763–2774.
- Thibaud J.B., Grignon C. (1981): Mechanism of nitrate uptake in corn roots. *Plant Sci. Lett.*, 22: 279–289.
- Thomine S., Lelièvre F., Debarbieux E., Schroeder J.I., Brabier-Brygoo H. (2003): AtNRAMP3, a multispecific vacuolar metal transporter involved in plant responses to iron deficiency. *Plant J.*, 34: 685–695.
- Wu F., Zhang G., Yu J. (2003): Interaction of cadmium and four microelements for uptake and translocation in different barley genotypes. *Commun. Soil Sci. Plant Anal.*, 34: 2003–2020.
- Zornoza P., Vázquez S., Esteban E., Fernández-Pascual M., Carpena R. (2002): Cadmium-stress in nodulated white lupin: strategies to avoid toxicity. *Plant Physiol. Biochem.*, 40: 1003–1009.

Received on April 27, 2005

---

*Corresponding author:*

RNDr. Peter Paľove-Balang, Ph.D., Botanický ústav, Slovenská akadémia vied, Dúbravská cesta 14, 845 23 Bratislava, Slovenská republika  
phone: + 421 259 426 127, fax: + 421 254 771 948, e-mail: peter.palove-balang@savba.sk

---