

# *Silybum marianum* in vitro-flavonolignan production

L. Tůmová<sup>1</sup>, J. Řimáková<sup>1</sup>, J. Tůma<sup>2</sup>, J. Dušek<sup>1</sup>

<sup>1</sup>School of Pharmacy, Charles University, Hradec Kralove, Czech Republic

<sup>2</sup>Faculty of Education, University of Hradec Kralove, Czech Republic

## ABSTRACT

The effect of coniferyl alcohol as a precursor of flavonolignan biosynthesis on silymarin components production in *Silybum marianum* suspension culture was studied. Coniferyl alcohol showed the changes in silymarin complex production. Silydianin was detected mainly in the control samples of cultivated cells. A significant increase of silydianin was observed only after 72 h of the application of 46 μM coniferyl alcohol. No other components of the silymarin complex (silychristin and silybin) were detected; neither in control samples nor after the precursor feeding. But the increased accumulation of taxifolin (flavanole) was very interesting. The highest taxifolin production was reached after 48 hours of treatment (about 554% compared to control).

**Keywords:** milk thistle; *Silybum marianum*; in vitro culture; flavonolignans; coniferyl alcohol

Milk thistle, *Silybum marianum* (L.) Gaertn., Asteraceae is widely used for hepatic and biliary disorders in Europe; it has been used medicinally since the first century. It was also mentioned in the writings of Dioscorides, Jacobus Theodorus and Culpepper. Its leaves, flowers and roots were historically considered a vegetable in European diets, and fruits (achenes), which resemble seeds, were roasted as a coffee bean substitute. The leaves of the plant are eaten in fresh salads and as a spinach substitute, the stalks like asparagus, and the flower heads serve as artichokes (Luper 1998).

The milk thistle is an annual herb or biannual herbaceous plant that is widespread in temperate climate of the American countries, Australia, Southern and Western Europe. This plant is also cultivated in the Czech Republic.

The dried seeds contain 1–4% of silymarin flavonoids. Silymarin is a mixture of three flavonolignans, including silybin (silibinin), silidianin, and silichrystin (Figure 1). Other flavonolignans identified in *S. marianum* include dehydrosilybin, deoxysilycistin, deoxysilydianin, silandrin, silybinome, silyhermin, and neosilyhermin. In addition, milk thistle contains apigenin, taxifolin, silybonol, myristic, oleic, palmitin, and stearin acids (Wichtl 1994).

The parts of silymarin complex are of a considerable pharmacological interest owing to their strong anti-hepatotoxic and hepatoprotective activity (Valenzuela et al. 1986, Morazoni and Bombardelli 1995), and a recent work has shown that silymarin also acts as an anticholesterolaemic agent (Krecman et al. 1998). Milk thistle fruit is used for the prophylaxis and treatment of liver damage caused by metabolite toxins, e.g. alcohol, tissue poison, in liver dysfunction and after hepatitis (post-hepatic syndrome), and in chronic degenerative liver conditions (Wichtl 1994).

*In vitro* culture of cells and tissues may offer an alternative for the production of silymarin but until now, only few studies addressing this possibility have been carried out (Becker and Schroll 1977), and in all cases flavonolignan production in *in vitro* cultures is very low and even disappears in prolonged cultures. This is a common occurrence in the production of secondary metabolites in many species. However, in some cases (Corchete et al. 1991, Hagendoorn et al. 1994), the manipulation of the components of the culture medium substantially increased the production of these compounds.

Feeding experiments with the precursors of biosynthesis of secondary metabolites resulted

---

Supported by the Ministry of Education, Youth and Sports of the Czech Republic, Project No. MSM 0021620822.




Figure 1. Main components of silymarin complex and taxifolin structure

in a 3–5 fold increase in levels of secondary metabolites. For example precursor L-phenylalanine showed a 3–5 fold increase in 5-methoxypodophyllotoxin in suspension culture of *Linum flavum* (van Uden et al. 1990).

Coniferyl alcohol is a key precursor of biosynthetic pathway of flavonolignans. Flavonolignans are adducts of flavanole taxifolin with coniferyl alcohol (Figure 2). Coniferyl alcohol was chosen as a model substance in this study. The aim of our experiment was to increase flavonolignan production by feeding nutrient medium with precursor of coniferyl alcohol. The same precursor – coniferyl alcohol in the form of complex with  $\beta$ -cyclodextrin was used as precursor for podophyllotoxin accumulation in *Podophyllum hexandrum* cell suspension cultures. The  $\beta$ -cyclodextrin itself was proven to be non-toxic for cells. It did not influence podophyllotoxin content and it was not metabolized or used as a carbon source by cells. For comparison, coniferin, the water-soluble  $\beta$ -D-glucoside of coniferyl alcohol, was also fed in the same concentrations. The effect of coniferin on the podophyllotoxin accumulation was stronger than that of coniferyl alcohol complexed with  $\beta$ -cyclodextrin, but coniferin is not commercially available (Woerdenbag et al. 1990).

## MATERIAL AND METHODS

**Plant material.** Cell suspensions were established from 3-month-old undifferentiated cotyledon calluses and grown in MS liquid medium (Murashige and Skoog 1962) supplemented with 10 mg/dm<sup>3</sup> NAA. These suspensions were cultivated in growth chambers at 25°C in the shaker at 120 rpm and 16 hours daylight. Ethanollic coniferyl alcohol solution in concentrations  $c_1$  33.11 mg/dm<sup>3</sup> (184 $\mu$ M);  $c_2$  16.56 mg/dm<sup>3</sup> (92 $\mu$ M) and  $c_3$  8.28 mg/dm<sup>3</sup> (46 $\mu$ M) was added into MS nutrient medium as biosynthesis precursor of flavonolignans. After 12, 24, 48, 72, and 168 hours of precursor application, the suspension cells were taken out. The cells were separated from nutrient medium by a reduced press filtration and were dried at the laboratory temperature. The flavonolignan content was detected not only in suspension cells but also in the nutrient medium.

**Analytical method** (Řimáková 2005). Dried suspension cells were extracted in methanol for 30 min and methanolic solution was concentrated to 5 ml and analyzed by HPLC. Solid fraction after evaporation of nutrient medium was dissolved in 5 ml of methanol. The content of flavonolignans was determined by HPLC on a UNICAM CRYSTAL






Figure 3. Taxifolin content in the medium after the treatment of *Silybum marianum* suspension culture with coniferyl alcohol

podophyllotoxin production, but only to a maximum of 0.006%. Feeding with 3mM coniferin caused a giant increase, with the maximum of 0.056% on day 10 (Woerdenbag et al. 1990).

Increased silydianin production after coniferyl alcohol feeding (as a biosynthesis precursor) was also achieved but other components of silymarin complex were not influenced. It could be caused by the use of free coniferyl alcohol only, not in the form with complex of  $\beta$ -cyclodextrin for better water solubility. Some precursors are very poorly soluble in aqueous media and reaction rates are therefore too low to be detected. So far, poorly water-soluble precursors have often been applied in two-phase systems. However, many of the plant cells used hardly converted precursors in the presence of organic phases, often due to dramatic decrease of cell vitality (Beiderbeck and Knoop 1988).

From the results presented, it is concluded that feeding a medium with precursor of coniferyl alcohol occurring in the lignan biosynthesis offers the possibility to enhance the content of some components of silymarin complex and flavonoid-taxifolin in *Silybum marianum* cultures *in vitro*.

## REFERENCES

Becker H., Schroll R. (1977): Callus und suspension-kulturen von *Silybum marianum*. *Planta Med.*, 31: 185–192.

Beiderbeck R., Knoop B. (1988): Enhanced production of secondary substance: addition of artificial accumulation sites to cultures. In: Bajaj Y.P.S. (ed.): *Biotechnology in Agriculture and Forestry* 4. Medical and

Aromatic Plants I. Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo: 123–135.

Corchete M.P., Jiménez M.A., Morán M., Cacho M., Fernández-Tárrago J. (1991): Effect of cell suspension cultures of *Digitalis thapsi* L. *Plant Cell Rep.*, 10: 394–396.

Fett-Neto A.G., Melanson S.J., Nicholson S.A., Pennington J.J., DoCosmo F. (2004): Improved taxol yield by aromatic carboxylic and aminoacid feeding to cell cultures of *Taxus cuspidata*. *Biotechnol. Bioengin.*, 8: 967–971.

Hagendoorn M.J.M., Wagner A.M., Segers G., van der Plas L.H.W. et al. (1994): Cytoplasmic acidification and secondary metabolite production in different plant cell suspensions. *Plant Physiol.*, 10: 723–730.

Krecman V., Skotová N., Waterová D., Ulrichová J., Simánek V. (1998): Silymarin inhibits the development of diet-induced hypercholesterolemia in rats. *Planta Med.*, 64: 138–142.

Luper S. (1998): A review of plants used in the treatment of liver disease: part 1. *Alter. Med. Rev.*, 3: 410–421.

Morazoni P., Bombardelli E. (1995): *Silybum marianum* (*Carduus marianum*). *Fitoterapia*, 66: 3–42.

Murashige T., Skoog F. (1962): A revised medium for rapid growth and bioassays with tobacco tissue cultures. *J. Plant Physiol.*, 15: 473–497.

Řimáková J. (2005): Possibility of influencing the secondary metabolite production in medicinal plants culture *in vitro*. [Doctor Thesis.] Faculty of Pharmacy, Charles University, Hradec Králové.

Valenzuela R., Guerra R., Videla L.A. (1986): Antioxidant properties of the flavonoids silybin and (+)-cyanidanol-3: comparison with butylated hydroxyanisole and butylated hydroxytoluene. *Planta Med.*, 52: 438–440.

Van Uden W., Pras N., Vossebeld E.M., Jos N.M. (1990): Production of 5-methoxypodophyllotoxin in cell suspension cultures of *Linum flavum* L. *Plant Cell Tiss. Organ Cult.*, 20: 81–87.

Wichtl M. (1994): *Herbal Drugs and Phytopharmaceuticals*. Medpharm Scientific Publishers, Stuttgart.

Woerdenbag H.J., van Uden W., Frijlink H.W., Lerk C.F., Pras N., Malingré T.M. (1990): Increased podophyllotoxin production in *Podophyllum hexandrum* cell suspension cultures after feeding coniferyl alcohol as a  $\beta$ -cyclodextrin complex. *Plant Cell Rep.*, 9: 97–100.

Received on March 24, 2006

---

*Corresponding author:*

Doc. PharmDr. Lenka Tůmová, CSc., Univerzita Karlova, Farmaceutická fakulta v Hradci Králové, katedra farmako-gnosie, Heyrovského 1203, 500 05 Hradec Králové, Česká republika  
phone: + 420 495 067 364, e-mail: tumova@faf.cuni.cz

---