

The effect of carbon source on plant regeneration in tomato

J. GUBIŠ, Z. LAJCHOVÁ, L. KLČOVÁ

Division of Applied Genetics and Breeding, Research Institute of Plant Production Piešťany, Piešťany, Slovak Republic

ABSTRACT: The effect of different tomato cultivars and different sugar types (sucrose, glucose and maltose) and concentrations (1.0, 2.0 and 3.0%) on shoot regeneration from aseptically grown hypocotyl and cotyledon explants were studied. Among sugar types, sucrose at a concentration of 3.0% induced the highest number of shoots from both types of explants. In hypocotyl explants, cv. Premium showed the best regeneration capacity (0.23 shoots per explant), and in cotyledon explants, cv. Hana produced the maximal number of shoots (0.43 or 0.37 for media with 2.0% or 3.0% sucrose, respectively).

Keywords: *in vitro* culture; MS medium; sugars; organogenesis

In vitro regeneration of cultivated tomato (*Lycopersicon esculentum* Mill.) has been a subject of research because of the commercial value of the crop and its amenability for further improvement via genetic manipulation (EVANS 1989). The most successful procedure is regeneration through adventitious organogenesis (VAN ROEKEL et al. 1993; FRARY, EARLE 1996; PERES et al. 2001). The *in vitro* morphogenetic responses of cultured plants are influenced by several different components of the culture media and it is important to evaluate their effect on plant regeneration. The purpose of our study was to evaluate the influence of carbon source on plant regeneration in tomato.

Three cultivars of tomato (*Lycopersicon esculentum* Mill.), Premium, Hana and UC 82, were used. The seeds were surface-sterilised by immersion into a 4% solution of sodium hypochlorite for 15 min and rinsed four times with sterile distilled water. The seeds were then germinated in glass containers with 25 cm³ of a half-strength medium of MURASHIGE and SKOOG (1962) (abbreviated hereinafter as "MS"), 100 mg/m³ myo-inositol, 2 mg/dm³ thiamine.HCl, 0.5 mg/dm³ pyridoxine.HCl, 0.5 mg/dm³ nicotinic acid, 1% sucrose and 0.7% agar. The cultures were initially kept in the dark at 27 ± 1°C for two days and then maintained under a 16h photoperiod at

50 µmol/m² s, with day/night temperature of 25°C/20°C. Hypocotyl and cotyledon segments were cut from the seedlings grown *in vitro*. The hypocotyls were cut into three segments. Each cotyledon was transversally cut into two segments. Hypocotyls were transversally cut into 4–7mm segments and leaf-blades into pieces of 30–40 mm². The hypocotyl explants were placed horizontally on the medium surface and cotyledon explants with the adaxial surface in contact with the medium. The effect of different concentrations of sucrose, glucose and maltose (1.0, 2.0 and 3.0%) that were added to the MS basal medium supplemented with 1 mg/dm³ zeatin (ZEA) and 0.1 mg/dm³ indole-3-acetic acid (IAA) (ICHIMURA, ODA 1995) on regeneration capacity of explants was studied. The media were adjusted to pH 5.8 prior to autoclaving. Glass containers with 25 cm³ of medium were used. The regeneration capacity of explants was assessed 6 weeks later. The following parameters were evaluated: frequency of regeneration (percents of regenerating explants) and the number of shoots per explant. Significance of differences between the results was estimated by analysis of variance (Statgraphics Version 5.0). Variation among means was analysed using LSD ($P \leq 0.05$) procedure according to the method described by SNEDECOR and COCHRAN (1956).

Supported by the Ministry of Agriculture of the Slovak Republic, Grant No. 2003 SP 27/028 0D 01/028 0D 01/03/02/06.

Table 1. Influence of different concentrations (1, 2 and 3%) of sugars on plant regeneration in tomato – the number of shoots/plated explant are means of 30 explants; data were taken after 6 weeks of culture. Different letters mean significant difference at $P \leq 0.05$

Sugar (%)	Hypocotyls				Cotyledons				
	Premium	Hana	UC 82	mean	Premium	Hana	UC 82	mean	
Sucrose	1.0	0.03	0.00	0.00	0.01 ^a	0.07	0.10	0.03	0.07 ^{ab}
	2.0	0.00	0.07	0.00	0.02 ^a	0.03	0.43	0.07	0.18 ^b
	3.0	0.23	0.03	0.03	0.10 ^b	0.03	0.37	0.13	0.18 ^b
Glucose	1.0	0.00	0.00	0.00	0.00 ^a	0.07	0.33	0.03	0.14 ^{ab}
	2.0	0.00	0.03	0.03	0.02 ^a	0.07	0.30	0.13	0.17 ^{ab}
	3.0	0.10	0.03	0.00	0.04 ^a	0.10	0.17	0.03	0.10 ^{ab}
Maltose	1.0	0.00	0.00	0.00	0.00 ^a	0.10	0.20	0.00	0.10 ^{ab}
	2.0	0.13	0.00	0.00	0.04 ^a	0.27	0.23	0.00	0.17 ^{ab}
	3.0	0.07	0.07	0.00	0.05 ^a	0.00	0.13	0.03	0.05 ^a
Cultivar class means		0.06 ^b	0.03 ^a	0.01 ^a		0.08 ^a	0.25 ^b	0.05 ^a	

The use of different tomato cultivars and different sugar types and concentrations for shoot regeneration from aseptically grown hypocotyls and cotyledons resulted in significant effects of both the genotypes and the sugars. A large variability in the number of shoots was observed across cultivars and across the different sugar concentrations (Table 1). Cultivar mean comparisons (LSD, $\alpha = 0.05$) showed only two classes differing in the induction potential. The cultivar that produced the highest number of shoots per hypocotyl (0.06) was cv. Premium while cv. Hana produced the highest number of cotyledon explants (0.25). However, the lowest number of shoots per explant in both explant types was induced in the cv. UC 82. The highest induction class of cotyledon explants included two different concentrations (2.0 and 3.0%) of sucrose, which produced an average number of regenerated shoots per explant 0.18 (Table 1). Contrary to sucrose, the medium supplemented with 2.0% glucose or maltose also showed a good regeneration capacity (0.17) on cotyledon explants. For hypocotyl explants the highest number of shoots was regenerated on the medium with 3.0% sucrose.

Normally for the cells, tissues or organ cultures it is essential to add a carbon source into the growth medium. Sucrose is almost universally used for micropropagation purposes as it is readily utilisable by cells. The optimum concentration of sucrose required induction of morphogenesis or growth differs among genotypes. Sucrose seems to be essential for the healthy growth of tomato cultures, and most researchers have used it as the sole source of 'carbon' (CHEN et al. 1999; COSTA et al. 2000; VENKATACHALAM et al. 2000).

Sucrose concentration of 30 g/dm³ (compared to 5, 10, or 20 g/dm³) was found to be optimal for the microplant growth of tomato (SCHNAPP, PREECE 1986). The majority of researchers have used this sucrose concentration in their initiation and multiplication media (CHEN et al. 1999; COSTA et al. 2000; VENKATACHALAM et al. 2000).

EL-BAKRY (2002) compared the effect of carbon source on shoot induction and plant regeneration in tomato. He used glucose, fructose, maltose and sucrose at a 3% concentration. In his experiment maltose gave the highest number of shoots in the genotypes he used (Castlerock, UC97-3 and Peto 86).

Acknowledgements

We thank Doc. M. VALŠÍKOVÁ from the Research Institute of Vegetables in Nové Zámky for supplying seeds of the tomato cultivars.

References

- COSTA M.G.C., NOGUEIRA F.T.S., FIGUEIRA M.L., OTONI W.C., BROMMONSCHENKEL S.H., CECON P.R., 2000. Influence of the antibiotic timentin on plant regeneration of tomato (*Lycopersicon esculentum* Mill.) cultivars. *Plant Cell Reports*, 19: 327–332.
- CHEN H.Y., ZHANG J.H., ZHUANG T.M., ZHOU G.H., 1999. Studies on optimum hormone levels for tomato plant regeneration from hypocotyl explants cultured *in vitro*. *Acta Agriculture Shanghai*, 15: 26–29.
- EL-BAKRY A.A., 2002. Effects of genotype, growth regulators, carbon source, and pH on shoot induction and plant regeneration in tomato. *In vitro Cellular and Developmental Biology – Plant*, 38: 501–507.

- EVANS D.A., 1989. Somaclonal variation – genetic basic and breeding applications. *Trends in Genetics*, 5: 46–50.
- FRARY A., EARLE E.D., 1996. An examination of factors affecting the efficiency of *Agrobacterium*-mediated transformation of tomato. *Plant Cell Reports*, 16: 235–240.
- ICHIMURA K., ODA M., 1995. Stimulation of shoot regeneration from cotyledon segments of tomato (*Lycopersicon esculentum* Mill.) by agar and its extract. *Journal of the Japanese Society of Horticultural Science*, 64: 135–141.
- MURASHIGE T., SKOOG F., 1962. A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiologia Plantarum*, 15: 473–497.
- PERES L.E.P., MORGANTE P.G., VECCHI C., KRAUS J.E., VAN SLUYS M.A., 2001. Shoot regeneration capacity from roots and transgenic hairy roots of tomato cultivars and wild related species. *Plant Cell Tissue Organ Culture*, 65: 37–44.
- SCHNAPP S.R., PREECE J.E., 1986. *In vitro* growth reduction of tomato and carnation microplant. *Plant Cell Tissue Organ Culture*, 6: 3–8.
- SNEDECOR G.W., COCHRAN W.G., 1956. *Statistical Methods* 5th ed. Ames, Iowa, The Iowa State University Press: 367.
- VAN ROEKEL J.S.C., DAMM B., MELCHERS L.S., HOEKEMA A., 1993. Factors influencing transformation frequency of tomato (*Lycopersicon esculentum*). *Plant Cell Reports*, 12: 644–647.
- VENKATACHALAM P., GEETHA N., PRIYA P., RAJASEGER G., JAYABALAN N., 2000. High frequency plantlet regeneration from hypocotyl explants of tomato (*Lycopersicon esculentum* Mill.) via organogenesis. *Plant Cell Biotechnology and Molecular Biology*, 1: 95–100.

Received for publication September 29, 2004
Accepted after corrections November 11, 2004

Vplyv zdroja uhlíka na regeneráciu rastlín rajčiaka

ABSTRAKT: Bol študovaný vplyv rozdielnych odrôd rajčiaka jedlého, typov cukru (sacharóza, glukóza a maltóza) a ich koncentrácie (1, 2 a 3%) na regeneráciu výhonkov z asepticky naklíčených hypokotylových a klíčolistových explantátov. Spomedzi rôznych typov cukrov 3% sacharóza indukovala najvyšší počet výhonkov pri oboch typoch explantátov. Najlepšiu regeneračnú schopnosť (0,23 výhonku na explantát) pre hypokotylové explantáty vykazovala odroda Premium a maximálny počet výhonkov (0,43 alebo 0,37 pre médium s 2 % alebo 3 % sacharózy) pri klíčolistových explantátoch produkovala odroda Hana.

Kľúčové slová: *in vitro* kultúra; MS médium; cukry; organogenéza

Corresponding author:

Ing. JOZEF GUBIŠ, Výskumný ústav rastlinnej výroby, Divízia aplikovanej genetiky a šľachtenia, Bratislavská cesta 122, 921 68 Piešťany, Slovenská republika
tel.: + 421 337 722 311, fax: + 421 337 726 306, e-mail: gubis@vurv.sk
