

Porcine neonates failing vitality score: physio-metabolic profile and latency to the first teat contact

M.E. TRUJILLO-ORTEGA¹, D. MOTA-ROJAS², O. JUÁREZ², D. VILLANUEVA-GARCÍA³, P. ROLDAN-SANTIAGO², M. BECERRIL-HERRERA⁴, R. HERNÁNDEZ-GONZÁLEZ⁵, P. MORA-MEDINA¹, M. ALONSO-SPILSBURY², A.M. ROSALES², R. MARTÍNEZ-RODRÍGUEZ¹, R. RAMÍREZ-NECOECHEA²

¹National Autonomous University of Mexico (UNAM), FMVZ, CEIEPP and FESC, Mexico City and State of Mexico, Mexico

²Department of Animal Production and Agriculture, Stress Physiology and Farm Animal Welfare, Metropolitan Autonomous University, Campus Xochimilco (UAM-X), Mexico D.F., Mexico

³Division of Neonatology, Mexican Children's Hospital "Federico Gomez", Mexico D.F., Mexico

⁴EIAH- Benemerita Autonomous University of Puebla, Teziutlán, Puebla, Mexico

⁵Department of Experimental Research and Animal Resources, National Institute of Medical Sciences and Nutrition "Salvador Zubirán", Mexico D.F., Mexico

ABSTRACT: The objective of this study was to compare the metabolic and electrolytic profile as well as the morphological appearance of the umbilical cord and newly born piglets' weight that failed the vitality test scale compared to those who passed. Newborn piglets were divided into three groups according to the vitality with a modified Apgar score at birth: Group 1, failing with a score < 5 (G_1 ; $n = 218$), Group 2 had a score of 6 to 7 (G_2 ; $n = 439$) and Group 3 had scores > 8 (G_3 ; $n = 464$). Results showed significant differences among groups ($P < 0.05$) in the physio-metabolic pH, PCO_2 , PO_2 , Na^+ , Ca^{2+} , glucose, lactate and bicarbonate values. Regarding weight, temperature and latency to connect the maternal teat, there were also significant differences ($P < 0.05$) among groups; it took 23.38 min for G_3 while neonatal piglets from G_1 took 30 min longer ($P < 0.05$) to make the first teat contact. The neonates from the latter group had a higher percentage (75.68%) of broken umbilical cords, with higher birth weight (+200 g, $P < 0.05$), showed higher than normal blood glucose concentrations, and had lower body temperature at birth ($-0.7^\circ C$, $P < 0.05$) and PO_2 in comparison with the other groups of neonates that passed the vitality score. A novel point of this study is the profile characterization of piglets that failed and passed the vitality score; we expect that the data provided may be applicable as reference values of metabolic and electrolyte blood profiles in newborn piglets according to their vitality. In conclusion, this study demonstrates that low vitality newborn piglets had clearly undergone through perinatal asphyxia. Potential indicators increasing this condition are: high birth weight, low body temperature, vitality score ≤ 5 , and the presence of the broken umbilical cord at birth.

Keywords: piglet; welfare; haemodynamics; acidosis; asphyxia; neonate; Apgar score

Clinical studies by Mota-Rojas et al. (2007) carried out on commercial pig farms showed that out of 1000 newly born piglets between 150 and 200 animals experienced apnoea after birth and it was related to perinatal asphyxia longer than 30 s, affecting the vitality in a negative way and delaying

the latency of connecting to the dam's teat, which represents an economic impact for the producer. In another study where 11 324 farrowings were analysed from 5000 breeding Mexican pig farms, Mota-Rojas and Ramírez (1996) concluded that the cost of each newborn was approximately 16 USD, which

multiplied by the number of stillbirths per year (28 430) represented an annual approximated loss of 445 000 USD. Therefore it is important to have the resources that allow quick and efficient evaluations of the physiological state of newborn piglets. The number of piglets born alive is an indicator of the economic benefits of a farm (Houška et al., 2010).

The transitional period during expulsion from the uterus to the birth of a newborn is of utmost importance since many adaptation problems may be detected, such as thermoregulation alterations and breathing difficulties (Odehnalová et al., 2008). The understanding of normal transitional physiological changes of the newborn piglets is important for evaluating their vitality (Sanchez et al., 2009; Maniau et al., 2010). The vitality of the neonatal piglets is described as the capacity to develop and survive, recovering from any imbalance caused by stress in the birth process. Vitality of the neonatal piglet depends on various factors (see review by Alonso-Spilsbury et al., 2007) influenced by the mother, the piglet and the environment.

The Apgar scoring system is an easy and reliable method for routine assessment of both human and animal neonates, in terms of viability and short-term survival prognosis (Alonso-Spilsbury et al., 2005; Castagnetti et al., 2007, 2010; Veronesi et al., 2009; González-Lozano et al., 2009; Orozco-Gregorio et al., 2010). However, its use is not widespread in veterinary medicine. In addition, behavioural traits such as time until first upright standing and latency to first udder contact are indicators of pig vitality (Leenhouders et al., 2001; González-Lozano et al., 2010). Randall (1971) demonstrated effects of viability scoring at birth on the time to reach the udder and colostrum intake; low viability scores were directly correlated with low birth weights and hypercapnia. Similarly, Zaleski and Hacker (1993) showed that viability score was highly correlated with pH and PCO_2 and De Roth and Downie (1976) studies found a strong relation between birth weight and viability scores.

Another indicator of piglet vitality is the umbilical cord (UC) pathology. Morphologic aspects of the UC such as adherence and rupture have usually been studied and retrospectively correlated with the perinatal outcome in piglets (Mota-Rojas et al., 2002, 2005a,b,c; Castro-Nájera et al., 2006).

Although breathlessness and hypothermia usually represent less severe neonatal welfare insults than do hunger, sickness and pain (Mellor and Stafford, 2004), measurements of blood gases and estimates of non-invasive physiological variables of the new-

born piglets still provide crucial information on oxygen delivery, important for subsequent survival. Reference values of serum electrolytes in healthy piglets during the first 48 h of life (Wehrend et al., 2003) and acid-base balance in piglets at birth (van Dijk et al., 2006) were described elsewhere. Also, there are references for blood biochemistry for survivors and piglets that died during the first 10 days of life (Tuchscherer et al., 2000). However, information on the metabolic and electrolyte state of the neonates that failed the vitality scale test at birth is scarce (Herpin et al., 1996; Casellas et al., 2004). Thus, the objective of this study was to thoroughly compare metabolic and electrolytic blood profiles as well as the morphological appearance of the umbilical cord and ability to contact the dam teat in piglets with different vitality scores at birth.

MATERIAL AND METHODS

Animals, facilities and management

This is a retrospective study, evaluating data from all neonates that were expelled (1121 newborn piglets) from 100 hybrid Yorkshire × Landrace sows from 1st to 5th parities, as part of a more comprehensive larger project dealing with asphyxia of newly born piglets (Trujillo-Ortega et al., 2007). The experiment was carried out on a commercial pig farm with intensive production, located in central Mexico. During the experiment, all sows remained in farrowing crates, following the guidelines established by the Mexican Official Standard (NOM-062-ZOO, 1999). Sows were fed twice daily with a commercial lactation diet. Dams and their litters had ad libitum access to water.

Sows were moved from the gestation area (housed in stalls) to the farrowing unit with individual crates one week before the expected farrowing date. As a standard procedure on the farm, prostaglandins were used to synchronize farrowings. The farrowing unit had electronic ventilation systems and was lit with both natural and electric light set at 39.8 candelas, room temperature averaged 26°C with a relative humidity of 60%.

Variables

To avoid interference with the vitality criteria, no aid was given to newborn piglets at the time of

farrowing and during the vitality evaluation. The neonates were evaluated at birth using Randall's (Randall, 1971) adaptation of the Apgar score for human neonates, described by Zaleski and Hacker (1993) and modified by Mota-Rojas et al. (2005c), measuring the following variables within one minute after birth: heart rate (beats per min): < 120 (bradycardia), between 121 and 160 (normal), and > 161 (tachycardia); latency to breathing (min), the interval between birth and the first breath: > 1 min, between 16 s and 1 min, and < 15 s; colour of the skin on the snout: pale, pink or cyanotic; latency to standing, measured as the interval between birth and the first time the neonate stands on all four legs. This was classified as > 5 min, between 1 and 5 min, and < 1 min and the skin stain with meconium as severe, mild, or absent (Mota-Rojas et al., 2006). Each variable was scored from 0 (the least favourable) to 2 (the most favourable) and a global score ranging from 1 to 10 was obtained for each neonatal piglet. A scoring of less than 6 indicated failure to pass the vitality scale test.

Heart rate was measured with a stethoscope. The first breath was considered when a movement was observed in the thoracic area accompanied by exhalation of air. The time it took each piglet to stand up was measured with a chronometer until the newborn was able to stand on all four feet. In addition to the scale, the latency to the first mammary contact was recorded through individual piglet observation from continuous video recording. This latency was the interval between birth and the moment when the newborn first made a contact with the dam's teat, independently if there was milk consumption or not. The time registered for this variable began from the moment the newborn was relocated close to the mother's vulva, immediately after taking the first blood sample. All the piglets were monitored for blood values and vitality score, and were individually identified with numbers on their backs.

The inclusion criteria to form the groups were taken exclusively from the vitality score, independently of the physio-metabolic profile, birth order and gender. Animals were classified into three groups: Group 1 (G_1 ; $n = 218$) made up of neonates that failed the vitality scale test with a rating of ≤ 5 ; Group 2 (G_2 ; $n = 439$) included those with a score between 6 and 7 points, and Group 3 (G_3 ; $n = 464$) those newborn piglets with a vitality score ≥ 8 points; all vitality measurements were done within the first minute of life.

The blood samples of the neonates were taken immediately after birth, prior to the onset of regular respiratory movements. Sampling was performed by retro-orbital bleeding with a 100 μ l microcapillary tube in the medial canthus of the eye at a 30°–45° angle towards the back of the eye and inserting the needle into the venous suborbital sinus, according to the technique used by Zaleski and Hacker (1993) and Orozco-Gregorio et al. (2008). The pH, partial pressure of carbon dioxide (PaCO₂/mmHg) and oxygen (PaO₂/mmHg), electrolytes [Na⁺, K⁺ and Ca²⁺ (mEq/l)], glucose and lactate (mg/dl), HCO₃⁻ (mmol/l) levels and haematocrit (%) were analysed simultaneously using a third generation gas analyser (GEM Premier 3000, Instrumentation Laboratory Co., Lexington, USA and Instrumentation Laboratory SpA, Milano, Italy).

Body temperature was recorded within the first and on the fifth minute after expulsion and was registered instantly (1 s) with a tympanic membrane thermometer (ThermoScan Braun GmbH, Kronberg, Germany). The tympanic temperature was used as it correlates with the cerebral temperature significantly better than the colonic temperature, and has been useful as non-invasive measurement in experiments involving survival after hypoxia/ischemia (Haaland et al., 1996a).

The umbilical cords were evaluated at birth and classified as attached (normal) or broken according to the criteria from Mota-Rojas et al. (2002). Having assessed the vitality and gross morphology of the umbilical cord, piglets were individually weighed using a digital scale (Salter Weight-Tronix Ltd., West Bromwich, UK).

The experiment received ethical approval by the Research Institutional Animal Care and Use Committee (IACUC), at the Faculty of Veterinary Medicine and Zootechnics at UNAM in Mexico City. To shorten the handling duration, the sampling occurred simultaneously whenever possible, e.g. by measuring heart rate and ear temperature at the same time. At the end, all procedures were carried out within 5 min.

Statistical analysis

A completely randomized design was used. Continuous data were summarized as mean \pm standard deviation and were compared among the three groups by means of the ANOVA test under the GLM (General Linear Models) procedure of

Table 1. Mean \pm standard deviation (SD) of blood critical values from newborn piglets classified according to the vitality score

Blood trait	Vitality score		
	G ₁ (0–5) (n = 218)	G ₂ (6–7) (n = 439)	G ₃ (8–10) (n = 464)
pH*	7.1 \pm 0.6 ^c	7.2 \pm 0.6 ^b	7.3 \pm 0.5 [§]
PCO ₂ (mmHg)	86.6 \pm 23.3 ^a	53.7 \pm 12.8 ^b	46.4 \pm 10.6 ^c
PO ₂ (mmHg)	21.7 \pm 10.5 ^c	26.2 \pm 6.8 ^b	27.9 \pm 7.3 ^a
Na ⁺ (mmol/l)	134.0 \pm 3.5 ^b	134.8 \pm 3.5 ^{ab}	135.7 \pm 3.8 ^a
K ⁺ (mmol/l)	6.8 \pm 0.6 ^a	6.6 \pm 0.6 ^a	6.7 \pm 4.3 ^a
Ca ²⁺ (mmol/l)	1.8 \pm 0.2 ^a	1.6 \pm 0.1 ^b	1.5 \pm 0.1 ^c
Glucose (mg/dl)	83.5 \pm 38.9 ^a	63.3 \pm 15.3 ^b	67.1 \pm 15.8 ^b
Lactate (mg/dl)	90.0 \pm 21.2 ^a	44.3 \pm 17.7 ^b	36.1 \pm 15.1 ^c
HCO ₃ ⁻ (mmol/l)	18.9 \pm 1.9 ^c	21.3 \pm 2.3 ^b	23.0 \pm 2.9 ^a
Hematocrit (%)	32.6 \pm 4.3 ^a	32.5 \pm 4.4 ^a	31.1 \pm 4.6 ^a

Within a row, means with different letters (a, b, c) differ (ANOVA, $P < 0.05$)

*Kruskal-Wallis test, expressed as a mean and standard deviation

the SAS statistical programme (2004). The multiple comparisons of means were performed using Tukey's test. In addition, the Kruskal-Wallis test was run for variables without normal distribution like the pH of the umbilical cord and a Chi-square test was performed to compare UC morphology among groups.

RESULTS

Metabolic profile

There were significant statistical differences ($P < 0.05$) among groups (G₁, G₂ and G₃) catalogued according to their vitality score in the

blood variables pH, PCO₂, PO₂, HCO₃⁻, glucose and lactate. There were also significant differences among groups in plasma electrolytes Na⁺ and Ca²⁺, ($P < 0.05$). We found no differences among groups for K⁺ concentrations ($P > 0.05$) and for the haematocrit percentages ($P > 0.05$) (Table 1).

Umbilical cord gross morphology

G₃ showed a large significant difference ($P < 0.0001$), with a higher percentage of normal intact umbilical cords compared to that of G₁. There was also a significant difference regarding broken UC ($P < 0.0001$) between groups, G₁ (failing) showed a two-fold higher incidence compared to G₂ (Table 2).

Table 2. Condition of the umbilical cord (number and percentage) of newborn piglets classified according to the vitality score

Umbilical cord condition	Vitality score			P-value
	G ₁ (0–5) (n = 218)	G ₂ (6–7) (n = 439)	G ₃ (8–10) (n = 464)	
	<i>n</i> (%)			
Adhered (normal)	53 (24.31)	358 (81.54)	420 (90.5)	0.0001
Broken (abnormal)	165 (75.68)	81 (18.45)	44 (9.48)	0.0001

Kruskal-Wallis test

Table 3. Mean \pm standard deviation (SD) of weight, sex, body temperature and latency to first teat contact from newborn piglets classified according to the vitality score

Trait	Vitality score		
	G ₁ (0–5) (<i>n</i> = 218)	G ₂ (6–7) (<i>n</i> = 439)	G ₃ (8–10) (<i>n</i> = 464)
Weight (g)*	1590.9 \pm 229.9 ^a	1472.4 \pm 207.3 ^b	1393.9 \pm 218.2 ^c
Body temperature at birth (°C)*	37.0 \pm 0.4 ^c	37.5 \pm 0.6 ^b	37.7 \pm 0.7 ^a
Body temperature at 5 min (°C)*	36.6 \pm 1.1 ^a	35.7 \pm 1.3 ^a	36.4 \pm 1.2 ^a
Latency to teat contact (min)*	54.4 \pm 8.9 ^a	29.2 \pm 10.9 ^b	23.3 \pm 8.1 ^c
Males†	131 (60.09%)	193 (43.96%)	197 (42.45%)
Females†	87 (39.90%)	246 (56.03%)	267 (57.54%)

Within a row, means with different letters (a, b, c) differ (*ANOVA, $P < 0.05$)

†Kruskal-Wallis test, $P = 0.05$

Weight, temperature, latency to connect the teat and gender

The weight of neonates that failed the vitality score in G₁ was higher by 197 g ($P < 0.05$) compared to the weight of newborn piglets in G₃ with high vitality scores (Table 3). Body temperature immediately after birth between the neonates in G₁ versus G₃ was by 0.7°C lower ($P < 0.05$). However, there were no differences among groups ($P > 0.05$) in body temperature at 5 min.

As expected and in agreement with Trujillo-Ortega et al. (2007), the high vitality score group (G₃) had the shortest time to contact the udder, whereas the group of newborn piglets that failed the vitality score (G₁) took 30 min longer ($P < 0.05$) to make the teat contact. For the vitality performance regarding gender in the failed viability neonates, 60% of the piglets were males, whereas the larger corresponding percentage of newborn piglets that reached passing qualifications was for the females.

DISCUSSION

Vitality: umbilical cord and asphyxia

A physio-metabolic profile suggested for the neonates from the study is depicted in Figure 1. The integrity of the umbilical cord and assessment of acid-base balance at birth correlated with neonatal survival and birth asphyxia (Zaleski and Hacker, 1993; Herpin et al., 1996; Mota-Rojas et al., 2005c; Van Dijk et al., 2006, 2008a,b). We found significant differences among groups regarding the umbilical

cord condition and the vitality score. Pigs with a score of ≤ 5 had a greater incidence of broken umbilical cords, in concordance with earlier studies from Zaleski and Hacker (1993).

Increased levels of plasma lactate in the failed scored vitality group confirm that this neonates suffered early tissue hypoxia (Ježková and Smrčková, 1990; Haaland et al., 1996b; Herpin et al., 1996). Recently, Harris et al. (2008) showed that piglet capacity to tolerate asphyxia depends on each individual, since there is a large variation in the cardiovascular response and cortisol concentrations allowing them to survive birth asphyxia. The incidence and the degree of birth asphyxia experienced in newborn piglets during birth are highly variable and rather unpredictable. Furthermore, Herpin et al. (1996) found a broad range of acid-base values at birth in different members of the same litter. Newly born piglets that failed the vitality scale in our study showed a blood rise in PCO₂ and lactic acid, with a low PO₂, in accordance with previous studies (Herpin et al., 1996; Orozco-Gregorio et al., 2007; Trujillo-Ortega et al., 2007). A reduction of oxygen supply through the umbilical cord leads to increased PCO₂ causing respiratory acidosis resulting in a higher risk of intrapartum anoxia and prenatal mortality (Mota-Rojas et al., 2005a,b,c). The neonatal piglets with a score of ≤ 5 on the vitality scale clearly suffered a degree of asphyxia during birth (Mota-Rojas et al., 2011).

Vitality: body temperature and blood gases

One of the consequences of apnoea is hypoxia. The neonatal piglets with a vitality of ≤ 5 had two




Figure 1. Physio-metabolic performance of porcine neonates with < 5 in the vitality score

notable characteristics – low body temperature and PO_2 within one min after birth. Both traits are related since hypoxia elicits an array of compensatory responses, among them is anapyrexia, the regulated drop of body temperature, which is a very efficient way to prevent cellular damage caused by hypoxia (Steiner et al., 2002). The general importance of anapyrexia resides in the fact that it reduces the consumption of oxygen and increases the affinity of haemoglobin to oxygen (Bicego et al., 2002; Steiner et al., 2002). Variations in temperature tend to affect the entire organism (Odehnalová et al., 2008); the neonate piglets that obtained a low vitality score showed hypercapnia, results in agreement with previous work by Randall (1971) and Herpin et al. (1996); whereas more recently, Groenendaal et al. (2009) indicated that hypothermia affects PCO_2 diminishing this gas concentration at low temperatures.

Vitality: acid-base imbalance and electrolyte profile

A novel point of this study is the blood profile characterization of piglets that either failed or passed the vitality score; the data provided may be applicable as

a reference for electrolyte performance in newborn piglets depending on the score, since we know the limited capacity of the neonatal kidney to retain and excrete sodium (Aggarwal et al., 2001).

As expected, pH values were lower for the low viability group compared with the medium and high viability groups, results are in accordance with studies of Herpin et al. (1996). Differences in the Ca^{2+} levels of neonatal piglets are probably due to changes in pH, affecting Ca^{2+} transport proteins. In other words, the more acid the plasma, the fewer binding sites with negative charge are available for setting the calcium to plasma proteins, hence the concentrations of Ca^{2+} ions were elevated in neonatal piglets with failed scores on the vitality scale.

Vitality: energy performance

Herpin et al. (1996) observed an increasing glucose concentration ($P < 0.01$) with blood PCO_2 in piglets with a low vitality score compared to the higher scored ones; similarly, the results from this study indicate that piglets that failed the vitality scale test showed higher than normal glucose concentrations compared to those piglets that passed. Higher blood glucose concentrations were reported in still-born and weak born piglets (Svendsen et al., 1986) and highly asphyxiated piglets (Herpin et al., 1996), whereas in newborn pigs with low birth weight the utilization of glucose as an energy source is more intensive (Kabalin et al., 2008). The larger difference in blood glucose concentration in the failed score group is most probably due to the effects of catecholamines and stimulation of liver glycogenolysis (Herpin et al., 1996), caused by the stress of neonates during birth, indicating adaptation problems to extra-uterine life (Tuchscherer et al., 2000).

Vitality: weight and asphyxia

Another factor that is predisposed to the process of asphyxia is the neonatal weight, in human babies low birth weight is related to poor vitality scores. Similarly, Okere et al. (1997) found that the piglet viability score was highly correlated with piglet weight ($r = 0.66$). However, in this study, piglets with a score of ≤ 5 were the heaviest among groups and showed asphyxia signs. A possible explanation is that heavier piglets had a greater difficulty in travelling down the birth canal, not to

forget that these failing neonatal pigs had a greater incidence of broken umbilical cords. These three characteristics – weight, a score of < 5 and broken umbilical cords, constitute a higher level of foetal hypoxia. According to Hoy et al. (1995) neonatal asphyxia delays the first contact with the udder and the first intake of colostrum and is associated with a reduction of rectal temperature at 24 h of life being of the prognostic value for early postnatal vitality.

Vitality: temperature and latency to teat contact

The survival of newborn piglets with low birth weight is reduced because they show poor thermoregulatory abilities and are slow to acquire colostrum (Baxter et al., 2008). Trujillo-Ortega et al. (2007) found a positive correlation between latency to the first teat contact and vitality score. Indeed, Leenhouders et al. (2001) results showed that the survival chances of small pigs decreased considerably with prolonged first colostrum intake. Surprisingly, as indicated in this study, neonates with a vitality score ≤ 5 weighed more among groups; this indicates that vitality scoring is independent of weight. Contrary to Hoy et al. (1994) and Baxter et al. (2008) results, it took the heavier piglets (G_1) longer to make the first udder contact.

Casellas et al. (2004) observed a positive correlation between viability score and rectal temperature at birth and 60 min after birth, whereas Leenhouders et al. (2001) reported no correlation between rectal temperature and early postnatal survival. In the present study, neonate piglets with a score ≤ 5 showed lower temperatures immediately after birth, results similar to those reported by Casellas et al. (2004) and Orozco-Gregorio et al. (2008), who found a correlation between low body temperature and lower vitality scores and prolonged time taken to contact the teat, which indicates deterioration of the newborn piglet self-regulation caused by not ingesting colostrum.

Vitality: gender

Gender differences were observed in the different vitality classes, males represented the highest percentage in the failed vitality score group (G_1). The physiological explanation has not been clear up

to now, however, regarding rabbit foetuses, Nielsen and Torday (1981) found that while the biochemical indices of lung maturation were the same for full-term males and females, the male foetuses reached the maturity values later than females. Moreover, in human medicine, low Apgar scores are predominantly for boys.

CONCLUSIONS

Traditionally, low birth weight piglets are more prone to have higher rates of pre-weaning mortality. Although we did not measure the mortality rates in this study, our results show that the piglets with a failing score on the vitality scale were the heaviest among groups, it took them 30 min longer to contact the maternal teat, they had a higher percentage of broken umbilical cords, low body temperature (first minute of life), low blood PO_2 and higher than normal blood glucose and lactate levels showing perinatal asphyxia, compared to the other groups of neonates with a passing score on the vitality test; thus this study demonstrates that low vitality piglets had clearly undergone through perinatal asphyxia. Our results confirm that hypoxia and anapyrexia lead to longer latencies to the first teat contact. Critical metabolic and electrolyte blood profiles and latency to the first teat contact according to the vitality scores of piglets at birth are provided as reference values. Gender differences in the vitality score are interesting findings and require further research, as females have significantly higher survival rates than males, despite the higher birth weight in the males.

Acknowledgment

Daniel Mota-Rojas, María Elena Trujillo-Ortega, Rafael Hernández-González, Marcelino Becerril-Herrera, María Alonso-Spilsbury and Ramiro Ramírez-Necoechea, were supported, as members, by the Sistema Nacional de Investigadores (SNI) in Mexico.

REFERENCES

- Aggarwal R., Deorari A.K., Paul K., (2001): Fluid and electrolyte management in term and preterm neonates. *Indian Journal of Pediatrics*, 68, 1139–1142.

- Alonso-Spilsbury M., Mota-Rojas D., Villanueva-García D., Martínez-Burnes J., Orozco G.H., Ramírez-Necoechea R., López, A., Trujillo-Ortega M.E. (2005): Perinatal asphyxia pathophysiology in fetal and human neonate: a review. *Animal Reproduction Science*, 90, 1–30.
- Alonso-Spilsbury M., Ramírez-Necoechea R., González-Lozano M., Mota-Rojas D., Trujillo-Ortega M.E. (2007): Piglet survival in early lactation: a review. *Journal of Animal and Veterinary Advances*, 6, 76–86.
- Baxter E.M., Jarvis S., D'Eath R.B., Ross D.W., Robson S.K., Farish M., Nevison I.M., Lawrence A.B., Edwards S.A. (2008): Investigating the behavioural and physiological indicators of neonatal survival in pigs. *Theriogenology*, 69, 773–783.
- Bicego K.C., Steiner A.A., Gargaglioni L.H., Branco L.G. (2002): Is lactate a mediator of hypoxia-induced ana-pyrexia? *Pflugers Archiv*, 444, 810–815.
- Casellas J., Rau W.M., Piedrafita J., Sánchez A., Arqué M., Noguera J.L. (2004): Viability of Iberian × Meishan F2 newborn pigs. I. Analysis of physiological and vitality variables. *Journal of Animal Science*, 82, 1919–1924.
- Castagnetti C., Mariella J., Serrazanetti G.P., Grandis A., Merlo B., Fabbri M., Mari G. (2007): Evaluation of lung maturity by amniotic fluid analysis in equine neonate. *Theriogenology*, 67, 1455–1462.
- Castagnetti C., Pirrone A., Mariella J., Mari G. (2010): Venous blood lactate evaluation in equine neonatal intensive care. *Theriogenology*, 73, 343–357.
- Castro-Nájera J.A., Martínez-Burnes J., Mota-Rojas D., Cuevas-Reyes H., López A., Ramírez-Necoechea R., Gallegos-Sagredo R., Alonso-Spilsbury M. (2006): Morphological changes in the lungs of meconium-stained piglets. *Journal of Veterinary Diagnostic Investigation*, 18, 622–627.
- Chou Y.H., Tsou-Yau K.I., Wang P.J. (1998): Clinical application of the measurement of cord plasma lactate and pyruvate in the assessment of high-risk neonates. *Acta Paediatrica*, 87, 764–768.
- De Roth L., Downie H.G. (1976): Evaluation of viability of neonatal swine. *Canadian Veterinary Journal*, 17, 275–279.
- Fahey J.O., King T.L. (2005): Intrauterine asphyxia: cellular respiration and acid-base balance. *Midwifery Womens Health*, 50, 498–506.
- Fraser D. (1990): Behavioural perspectives on piglet survival. *Journal of Reproduction Fertility*, 40 (Suppl.), 355–370.
- González-Lozano M., Trujillo-Ortega M.E., Becerril-Herrera M., Alonso-Spilsbury M., Ramírez-Necoechea R., Hernández-González R., Mota-Rojas D. (2009): Effect of oxytocin on critical blood variables from dystocic sows. *Veterinaria Mexico*, 40, 231–245.
- González-Lozano M., Trujillo-Ortega M.E., Becerril-Herrera M., Alonso-Spilsbury M., Rosales-Torres A.M., Mota-Rojas D. (2010): Uterine activity and fetal electronic monitoring in parturient sows treated with vetrabutin chlorhydrate. *Journal of Veterinary Pharmacology and Therapeutics*, 33, 28–34.
- Groenendaal F., De Vooght K.M., van Bel F. (2009): Blood gas values during hypothermia in asphyxiated term neonates. *Pediatrics*, 123, 170–172.
- Haaland K., Steen P.A., Thoresen M. (1996a): Cerebral, tympanic and colonic thermometry in the piglet. *Reproduction Fertility and Development*, 8, 125–128.
- Haaland K., Kofstad J., Apricena F., Thoresen M. (1996b): Haemoglobin is inversely related to plasma lactate and heart rate in the newborn piglet. *Biology of the Neonate*, 69, 350–356.
- Harris T.A., Healy G.N., Colditz P.B., Lingwood B.E. (2008): Associations between serum cortisol, cardiovascular function and neurological outcome following acute global hypoxia in the newborn piglet. *Stress*, 24, 1–11.
- Herpin P., Le Dividich J., Hulin J.C., Fillaut M., de Marco F., Bertin R. (1996): Effects of the level of asphyxia during delivery on viability at birth and early postnatal vitality of newborn pigs. *Journal of Animal Science*, 74, 2067–2075.
- Herpin P., Hulin J.C., Le Dividich J., Fillaut M. (2001): Effect of oxygen inhalation at birth on the reduction in early postnatal mortality in pigs. *Journal of Animal Science*, 79, 5–10.
- Houška L., Wolfová M., Nagy I., Csörnyei Z., Komlósi I. (2010): Economic values for traits of pigs in Hungary. *Czech Journal of Animal Science*, 55, 139–148.
- Hoy S., Lutter C., Wähner M., Puppe B. (1994): The effect of birth weight on the early postnatal vitality of piglets. *Deutsche Tierärztliche Wochenschrift*, 101, 393–396.
- Hoy S., Lutter C., Puppe B., Wahner, M. (1995): Correlations between vitality of newborn piglets, teat order, mortality, and live weight development up to weaning. *Berliner und Munchener Tierärztliche Wochenschrift*, 108, 224–228.
- Ježková D., Smrčková M. (1990): Variations of glucosae-mia and lactacidemia in pregnant sows, fetuses and in sows and piglets till the 10th day after delivery. *Veterinarni Medicina*, 35, 613–620. (in Czech)
- Kabalin A.E., Balenovic T., Speranda M., Gradinski-Vrbanc B., Speranda T., Valpotic I., Pavicic Z., Balenovic M., Valpotic H. (2008): Preliminary investigation of birth weight influence on growth, metabolite and biocatalysator concentrations in sera of suckling piglets. *Tieraerztliche Umschau*, 63, 142–145.
- Leenhouwers J.I., de Almeida C.A., Knol E.F., van der Lende T. (2001): Progress of farrowing and early post-

- natal pig behavior in relation to genetic merit for pig survival. *Journal of Animal Science*, 79, 1416–1422.
- Mainau E., Dalmau A., Ruiz-de-la-Torre J.L., Manteca X. (2010): A behavioural scale to measure ease of farrowing in sows. *Theriogenology*, 74, 1279–1287.
- Mellor D.J., Stafford K.J. (2004): Animal welfare implications of neonatal mortality and morbidity in farm animals. *Veterinary Journal*, 168, 118–133.
- Mota-Rojas D., Ramírez, N.R. (1996): Stillbirths. *Agro México*, 18, 227–234.
- Mota-Rojas D., Martínez-Burnes J., Trujillo-Ortega M. E., Alonso-Spilsbury M., Ramírez-Necoechea R., López-Mayagoitia A. (2002): Effect of oxytocin treatment in sows on umbilical cord morphology and meconium staining and neonatal mortality of piglets. *American Journal of Veterinary Research*, 63, 1571–1574.
- Mota-Rojas D., Martínez-Burnes J., Trujillo O.M.E., López M.A., Rosales T.A.M., Ramírez N.R., Orozco G.H., Merino P.A., Alonso-Spilsbury M. (2005a): Uterine and fetal asphyxia monitoring in parturient sows treated with oxytocin. *Animal Reproduction Science*, 86, 131–141.
- Mota-Rojas D., Nava-Ocampo A., Trujillo-Ortega M. E., Velázquez-Armenta Y., Ramírez-Necoechea R., Martínez-Burnes J., Alonso-Spilsbury M. (2005b): Dose minimization study of oxytocin in early labor in sows: Uterine activity and fetal outcome. *Reproductive Toxicology*, 20, 255–259.
- Mota-Rojas D., Martínez-Burnes J., Trujillo-Ortega M. E., López A., Rosales A.M., Ramírez-Necoechea R., Orozco H., Merino A., Alonso-Spilsbury M. (2005c): The effects of vetrabutín chlorhydrate and oxytocin on stillbirth rate and asphyxia in swine. *Theriogenology*, 64, 1889–1897.
- Mota-Rojas D., Martínez-Burnes J., Alonso-Spilsbury M., López A., Ramírez-Necoechea R., Trujillo-Ortega M.E., Medina-Hernández F.J., de la Cruz N.I., Albores-Torres V., Gallegos-Sagredo R. (2006): Meconium staining of the skin and meconium aspiration in porcine intrapartum stillbirths. *Livestock Science*, 102, 155–162.
- Mota-Rojas D., Acosta M.B., Olmos H.A., Aceves R.D. (2007): The newborn: factors related to low vitality. *Porcicultores y su Entorno*, 58, 80–85.
- Mota-Rojas D., Orozco-Gregorio H., Villanueva-García D., Bonilla-Jaime H., Suárez-Bonilla X., Hernández-González R., Roldan-Santiago P., Trujillo-Ortega M.E. (2011): Fetal and neonatal energy metabolism in pigs and humans: A review. *Veterinarni Medicina*, 56, 215–225.
- Nielsen H.C., Torday J.S. (1981): Sex differences in fetal rabbit pulmonary surfactant production. *Pediatrics Research*, 15, 1245–1247.
- NOM-062-ZOO (1999): Norma Oficial Mexicana: Especificaciones técnicas para la producción, cuidado y uso de los animales de laboratorio. Diario Oficial de la Federación. Miércoles 22 de agosto de 2001. México.
- Odehmalová S., Vinkler A., Novák P., Drábek J. (2008): The dynamics of changes in selected parameters in relation to different air temperature in the farrowing house for sows. *Czech Journal of Animal Science*, 53, 195–203.
- Okere C., Hacker R.R., Werchola G. (1997): Relationships between serum IGF-I concentrations and piglet development or neonatal viability following porcine somatotropin (pST) and insulin administration to gestating gilts. *Theriogenology*, 47, 1403–1412.
- Orozco-Gregorio H., Mota-Rojas D., Alonso-Spilsbury M., González-Lozano M., Trujillo-Ortega M.E., Olmos-Hernández S.A., Sánchez-Aparicio P. Ramírez-Necoechea R., Hernández-González R. Uribe-Escamilla R., Villanueva-García D. (2007): Importance of blood gas measurements in perinatal asphyxia and alternatives to restore the acid base balance status to improve the newborn performance. *American Journal of Biochemistry and Biotechnology*, 3, 131–140.
- Orozco-Gregorio H., Mota-Rojas D., Alonso-Spilsbury M., Olmos-Hernández A., Ramírez-Necoechea R.R., Velázquez-Armenta E.Y., Nava-Ocampo A.A., Hernández-González R., Trujillo-Ortega M.E., Villanueva-García D. (2008): Short-term neurophysiologic consequences of intrapartum asphyxia in piglets born by spontaneous parturition. *International Journal of Neuroscience*, 118, 1299–1315.
- Orozco-Gregorio H., Mota-Rojas D., Bonilla-Jaime H., Trujillo-Ortega M.E., Becerril-Herrera M., Hernández-González R., Villanueva-García D. (2010): Effects of administration of caffeine on metabolic variables in neonatal pigs with peripartum asphyxia. *American Journal of Veterinary Science*, 71, 1214–1219.
- Randall G.C.B. (1971): The relationship of arterial blood pH and pCO₂ to the viability of the newborn piglet. *Canadian Journal of Comparative Medicine*, 35, 141–146.
- Sanchez A.P., Mota-Rojas D., Trujillo-Ortega M.E., Zarco-Quintero L., Becerril-Herrera M., Alonso-Spilsbury M., Alfaro-Rodríguez A. (2009): Effect of prostaglandins for inducing birth, on weight, vitality and physiological response in newborn pigs. *Journal of Applied Animal Research*, 36, 113–118.
- Steiner A., Rocha M.J.A., Branco L.G. (2002): A neurochemical mechanism for hypoxia-induced anapnoea. *American Journal of Physiology*, 283, R1412–R1422.
- Svendsen J., Svendsen L.S., Bengtsson A.C. (1986): Reducing perinatal mortality in pigs. In: Leman, A.D., Straw

- B.E., Mengeling W.L., D'Allaire S., Taylor D.J. (eds): Diseases of Swine. Ames, USA, Iowa State University Press, 813–825.
- Trujillo-Ortega M.E., Mota-Rojas D., Olmos-Hernández A., Alonso-Spilsbury M., González M., Orozco H., Ramírez-Necochea R., Nava-Ocampo A. (2007): A study of piglets born by spontaneous parturition under uncontrolled conditions: could this be a naturalistic model for the study of intrapartum asphyxia? *Acta Biomedica*, 78, 29–35.
- Tuchscherer M., Puppe B., Tuchscherer A., Tiemann U. (2000): Identification of neonate at risk: traits of newborn piglets with respect to survival. *Theriogenology*, 54, 371–388.
- van Dijk A.J., van der Lende T., Taverne M.A.M. (2006): Acid-base balance of umbilical artery blood of liveborn piglets at birth and its relation with factors affecting delivery of individual piglets. *Theriogenology*, 66, 1824–1833.
- van Dijk A.J., van Loon J.P.A.M., Taverne M.A.M., Jonker F.H. (2008a): Umbilical cord clamping in term piglets: A useful model to anterior study perinatal asphyxia? *Theriogenology*, 70, 662–674.
- van Dijk A.J., Parvizi N., Taverne M.A.M., Fink-Gremmels J. (2008b): Placental transfer and pharmacokinetics of allopurinol in late pregnant sows and their fetuses. *Journal of Veterinary Pharmacology and Therapeutics*, 31, 489–495.
- Veronesi M.C., Panzani S., Faustini A., Rota A. (2009): An Apgar scoring system for routine assessment of newborn puppy viability and short-term survival prognosis. *Theriogenology*, 72, 401–407.
- Wehrend A., Failing K., Tschachtschal J., Bostedt H. (2003): Reference values of electrolyte plasma concentrations in healthy piglets during the first 48 hours of life. *Journal of the American Veterinary Medical Association*, 50, 67–71.
- Zaleski H.M., Hacker R.R. (1993): Comparison of viability scoring and blood gas analysis as measures of piglet viability. *Canadian Journal of Animal Science*, 73, 649.

Received: 2011–04–25

Accepted after corrections: 2011–05–05

Corresponding Author

Prof. Ph D. Daniel Mota-Rojas, Metropolitan Autonomous University (UAM-X), Department of Agriculture and Animal Production, Calzada del Hueso 1100, Colonia Villa Quietud, México D.F., 04960 México
Tel. +52 55 548 375 35, e-mail: dmota100@yahoo.com.mx
