

<https://doi.org/10.17221/51/2019-PPS>

The significance of anthropochory in *Hercinothrips femoralis* (Thysanoptera: Thripidae) – Short Communication

MARTIN ŠTEFÁNIK*, MARTINA ZVARÍKOVÁ, RUDOLF MASAROVIČ, PETER FEDOR

Department of Environmental Ecology, Faculty of Natural Sciences, Comenius University,
Bratislava, Slovakia

*Corresponding author: stefanik16@uniba.sk

Citation: Štefánik M., Zvaríková M., Masarovič R., Fedor P. (2019): The significance of anthropochory with *Hercinothrips femoralis* (Thysanoptera: Thripidae). *Plant Protect. Sci.*, 55: 262–265.

Abstract: After the first record of banded greenhouse thrips, *Hercinothrips femoralis* (Reuter, 1891) in Slovakia in 2008, a peculiar occurrence in the extreme subalpine environment of Podbanské (High Tatra Mts.), Slovakia, during the summer of the same year was observed. Since this species of exotic thrips is known to be almost sedentary, the mode of dispersal was hypothesised to be related to passive transport via weather currents. According to our observations of unintentional dispersal, a test was conducted to research a previously unidentified introduction pathway of this species. Our preliminary results show that passive transport by humans plays a crucial role in the dispersal of *H. femoralis*.

Keywords: anthropochorous; banded greenhouse thrips; basil; pest; spread; thrips

The importance of dynamic dispersal pathways in insects (STINNER *et al.* 1983; ROBINET *et al.* 2009; CARRASCO *et al.* 2010; BULLOCK *et al.* 2018), including thrips (KIRK & TERRY 2003; FEDOR 2004; FEDOR *et al.* 2010, MASAROVIČ *et al.* 2017) is widely recognised. The increases in global trade and the movement of people and goods inadvertently created and strengthened pathways for insects to invade previously inaccessible environments (KIRITANI 2001; KIRK & TERRY 2003; MAKRA *et al.* 2017). The minute size and often cryptic nature of thrips predisposes them as potentially effective pests, when in reality, only about 1%, out of more than 6 200 species of thrips (ThripsWiki 2019), are currently considered serious pests (FEDOR *et al.* 2004; MORSE & HODDLE 2005).

Hercinothrips femoralis is a known pest in greenhouses with records reaching back to the end of the

19th century, with the first description credited to Prof. O. M. Reuter (WHITE 1916). Few occurrences were recorded through the 20th century (HARTZELL 1926; BUCHANAN 1932; DENMARK 1976) until the second half of the century, when it resurfaced in Hungary (JENSER & CZENCZ 1988) and a first record in Australia was made (HOUSTON *et al.* 1991). Routine occurrences of *H. femoralis* in greenhouses around Europe were noted in recent decades with the first records from Slovenia (TRDAN 2002), Greece (RODITAKIS *et al.* 2006), Croatia (SIMALA & MILEK 2008) and Slovakia (VARGA 2008). The most recent reoccurrence was published in 2016 from South Korea (LEE & LEE 2016) where it was recorded severely damaging ornamental plants in a private household. The northernmost outdoor record came from Podbanské (the High Tatra Mountains), Slovakia, during June and July of 2008 (MASAROVIČ *et al.* 2014).

Supported by the Scientific Grant Agency of the Ministry of Education, Science, Research and Sport of the Slovak Republic and Slovak Academy of Science – VEGA, Grant No. 1/0104/16.

MATERIAL AND METHODS

During the winter of 2016, we commenced the rearing of *H. femoralis* for the purposes of invasive ecology research at the Faculty of Natural Sciences, Comenius University in Bratislava. The rearing was conducted by one researcher only with access to the grow room. The duties of the researcher consisted of watering the plants, the removal of the withered leaves and the exchange of the withered plants for fresh ones. After a month, we noticed the presence of *H. femoralis* in the researcher's office, where it already started to damage the ornamental plant *Tradescantia pallida* (Rose) DR Hunt. This pattern repeated with the most visited office spaces containing any ornamental plants. Therefore, we conducted a simple test of the dispersal rate and the conditions under which this rate may vary:

4 different indoor sites in a thrips free environment were chosen, each fitted with 2 basil plants (*Ocimum basilicum*), the host plant of *H. femoralis*. Sites 1–3 (university offices) were situated approximately 350 m from the laboratory with the grow room, as the centre of the *H. femoralis* distribution. Site 4 was located 3 km from the University building (a room in a private flat). The visiting frequency and duration for each site varied (Table 1). To maintain the same probability of introduction, only one site (1–3) was visited per day. Site 4 was visited daily, but the visit was conducted later in the day, after one of the sites 1–3 was visited. The light conditions were maintained by natural light and the temperature was maintained by a central heating system with a temperature range of 20–25°C. The laboratory grow room was fitted with 10 basil plants, which were changed for new ones as soon as any withering was observed. The basil plants in the grow room were introduced to approximately 300 *H. femoralis* females. The maintenance and thrips presence at sites 2 and 3 were checked by a second researcher. For each site, the date of the first appearance of the thrips and the

number of visits was recorded, if no observation was made after 150 days, the test was ended. After the successful detection of *H. femoralis*, the basil plants were removed from the study site.

The same test was repeated a second time with the grow room full of basil plants with a suboptimal water supply. The plants were changed for new ones, only if they were almost completely withered.

The collected specimens were mounted on slides using standard laboratory methods (MOUND & KIBBY 1998) and determined according to the nomenclature (ZUR STRASSEN 2003).

RESULTS AND DISCUSSION

The visiting frequency, the number of visits and days to the first thrips occurrence at each site is summarised in Table 1.

No thrips were observed at site 3 under optimal conditions, therefore, we do not refer to this observation further in the results. All remaining study sites were infested by thrips under optimal and suboptimal conditions, showing a pattern similar to the pre-test observations. The fastest introduction of *H. femoralis* to the new environment (8 days under the optimal conditions/6 days under the suboptimal conditions) was recorded at site 1. Although it took 31 days for *H. femoralis* to invade site 3 from suboptimal conditions, the observation was made right after the 2nd visit. Since site 4 was not visited right after the contact of the researcher with the host plants and the site was located further from the grow room, the period to the first detection of *H. femoralis* was longer compared to the office space, visited right after the interaction.

According to these preliminary results we believe that the record of *H. femoralis* in the subalpine environment of Podbanské (the High Tatra Mts.) made by MASAROVÍČ *et al.* (2014) may have been the result of an unintentional human-assisted dispersal.

Table 1. Monitoring of the *H. femoralis* dispersal under optimal and suboptimal conditions

Site	Visit length (h)	No. of visits	Days to first collected individual		No. of times the site was visited	
			optimal	suboptimal	optimal	suboptimal
S1	5 – 8	5 × week	8	6	8	6
S2	1	1 × week	35	28	5	4
S3	1	1 × 30 days	–	31	5	2
S4	6 – 12	home – daily	26	13	26	13

<https://doi.org/10.17221/51/2019-PPS>

Since the presence of *H. femoralis* was recorded at a greenhouse located at the same faculty (VARGA 2008) as Masarovič's research team, members of which repeatedly visited the sites at Podbanské (High Tatra Mts.), Slovakia and, thus, possibly introduced a population that was able to reside outdoors during the collection in June and July of 2008 (MASAROVÍČ *et al.* 2014). No further records suggest that the population was able to survive the latter months, as is often the case for exotic thrips species (MCDONALD *et al.* 1997; MCDONALD *et al.* 1999; MCDONALD *et al.* 2000; LARENTZAKI *et al.* 2007; RAMANAND *et al.* 2017). Although it seems that *H. femoralis* is not suited to survive the outdoors conditions in Slovakia throughout the whole year, novel indoor conditions such as office spaces that are kept at relatively constant temperatures, containing ornamental plants are providing a potential niche that could be exploited year-round (TRDAN *et al.* 2007; LEE & LEE 2016).

The effects on the speed of introduction was tested, by creating suboptimal rearing conditions, as previously demonstrated in *Frankliniella occidentalis* (Pergande, 1895) (RHAINDS & SHIPP 2009). Our simple test reveals a 31.67% average increase in the speed of introduction under the suboptimal conditions. It was previously suggested that *H. femoralis* is almost sedentary if the host plant is healthy and, therefore, the food source and possible oviposition sites are available. Although, if the food sources are scarce, *H. femoralis* starts to be much more mobile, searching for novel host plants (LAUGHLIN 1971).

The goal of this study was to add a possible pathway of introduction of *H. femoralis* not yet described by previous authors. The presented work demonstrates how an artificial environment with specific conditions creates new possibilities for the polyphagous opportunist *H. femoralis* and how the unintentional human-assisted dispersal pathway may be an important factor in the thrips introduction to novel environments. We present our preliminary results in hope that they may serve as a background for further and more complex studies of this dispersal mechanism.

References

- Buchanan D. (1932): A Bacterial disease of beans transmitted by *Hellothrips femoralis* (Reuter). *Journal of Economic Entomology*, 25: 49–53.
- Bullock J.M., Bonte D., Pufal G., da Silva Carvalho C., Chapman D.S., García C., García D., Matthysen E., Delgado M.M. (2018): Human-mediated dispersal and the rewiring of spatial networks. *Trends in Ecology & Evolution*, 33: 958–970.
- Carrasco L.R., Mumford J.D., MacLeod A., Harwood T., Grabenweger G., Leach A.W., Knight J.D., Baker R.H. (2010): Unveiling human-assisted dispersal mechanisms in invasive alien insects: Integration of spatial stochastic simulation and phenology models. *Ecological Modelling*, 221: 2068–2075.
- Denmark H.A. (1976): The banded greenhouse Thrips, *Hercinothrips Femoralis* (O. M. Reuter) damage to ornamental plants. *Proceedings of the Florida State Horticultural Society*, 89: 330–331.
- Fedor P. (2004): First records of *Dendrothrips degeeri* Uzel, 1895 (Thysanoptera, Thripidae) in Slovakia. *Biologia*, 59: 211–212.
- Fedor P., Sierka W., Majzlan O. (2004): The thrips (Thysanoptera) of Slovakia. *Acta Phytopathologica et Entomologica Hungarica*, 39: 301–309.
- Hartzell A. (1926): Naphthalene fumigation of greenhouses. *Journal of Economic Entomology*, 19: 780–786.
- Houston K.J., Mound L.A., Palmer J.M. (1991): Two pest thrips (Thysanoptera) new to Australia, with notes on the distribution and structural variation of other species. *Journal of the Australian Entomological Society*, 30: 231–232.
- Jenser G., Czencz K. (1988): Thysanoptera species occurring frequently on cultivated plants in Hungary. *Acta Phytopathologica et Entomologica Hungarica*, 23: 285–289.
- Kiritani K. (2001): Invasive insect pests and plant quarantine in Japan. *Extension Bulletin of the Food and Fertilizer Center*, 498: 1–12.
- Kirk W.D.J., Terry L.I. (2003): The spread of the western flower thrips *Frankliniella occidentalis* (Pergande). *Agricultural and Forest Entomology*, 5: 301–310.
- Larentzaki E., Powell G., Copland M.J.W. (2007): Effect of temperature on development, overwintering and establishment potential of *Frankliniella vespiformis* in the UK. *Entomologia Experimentalis et Applicata*, 124: 143–151.
- Laughlin R. (1971): A culture method for *Hercinothrips femoralis* (Reuter) (Thysanoptera). *Australian Journal of Entomology*, 10: 301–303.
- Lee G.S., Lee W. (2016): Rediscovery of an exotic thrips *Hercinothrips femoralis* (Reuter), in Korea, severely damaging to ornamental plants. *Korean Society of Applied Entomology*, 1: 102–102.
- Makra L., Bodnár K., Fülöp A., Orosz S., Szénási Á., Csépe Z., Jenser G., Tusnády G., Magyar D. (2017): The first record of subtropical insects (Thysanoptera) in central Europe: long-distance transport of airborne thrips, ap-

- plying three-dimensional backward trajectories. *Agricultural and Forest Entomology*, 20: 301–326.
- Masarovič R., Doričová M., Prokop P., Fedor P. (2014): “Testing the limits” - an interesting record of the exotic banded greenhouse thrips *Hercinothrips femoralis* (Thysanoptera: Thripidae: Panchaethripinae) at high Carpathian mountain altitudes. *Biologia*, 69: 1631–1634.
- Masarovič R., Zvaríková M., Fedorová J., Fedor P. (2017): Thigmotactic behavior of *Limothrips cerealium* (Thysanoptera: Thripidae) leads to laboratory equipment damage in the Czech Republic. *Journal of Entomological Science*, 52: 308–310.
- McDonald J.R., Bale J.S., Walters K.F.A. (1997): Low temperature mortality and overwintering of the western flower thrips *Frankliniella occidentalis* (Thysanoptera: Thripidae). *Bulletin of Entomological Research*, 87: 497–505.
- McDonald J.R., Bale J.S., Walters K.F.A. (1999): Temperature, development and establishment potential of Thrips palmi (Thysanoptera: Thripidae) in the United Kingdom. *European Journal of Entomology* 96: 169–174.
- McDonald J.R., Head J., Bale J.S., Walters K.F.A. (2000): Cold tolerance, overwintering and establishment potential of *Thrips palmi*. *Physiological Entomology*, 25: 159–166.
- Morse J.G., Hoddle M.S. (2005): Invasion biology of thrips. *Annual Review of Entomology*, 51: 67–89.
- Mound L.A., Kibby G. (1998): Thysanoptera: An Identification Guide. 2nd Ed. Oxford-New York, CAB International: 1–70.
- Ramanand H., McConnachie A.J., Olckers T. (2017): Thermal tolerance of *Liothrips tractabilis*, a biological control agent of *Campuloclinium macrocephalum* recently established in South Africa. *Entomologia Experimentalis et Applicata*, 162: 234–242.
- Rhainds M., Shipp L. (2009): Dispersal of adult western flower thrips (Thysanoptera: Thripidae) on chrysanthemum plants: impact of feeding-induced senescence of inflorescences. *Environmental Entomology*, 32: 1056–1065.
- Robinet C., Roques A., Pan H., Fang G., Ye J., Zhang Y., Sun J. (2009): Role of human-mediated dispersal in the spread of the pinewood nematode in China. *PLOS One* (serial online). 4: e4646. Available from www.journals.plos.org/plosone/ (accessed Mar 10, 2019).
- Roditakis E., Mound L.A., Roditakis N.E. (2006): NOTE: First record in Crete of *Hercinothrips femoralis* in greenhouse banana plantations. *Phytoparasitica*, 34: 488–490.
- Simala M., Milek T.M. (2008): Thysanoptera species recorded in greenhouses in Croatia from 2003–2006. *Acta Phytopathologica et Entomologica Hungarica*, 43: 373–383.
- Stinner R.E., Barfield C.S., Stimac J.L., Dohse L. (1983): Dispersal and movement of insect pests. *Annual Review of Entomology*, 28: 319–335.
- zur Strassen R. (2003): Die terebranten Thysanopteren Europas und des Mittelmeer-Gebietes. *Tierwelt Deutschlands*.
- ThripsWiki- providing information on the World's thrips - Available at http://thrips.info/wiki/Main_Page (accessed Mar 3, 2019).
- Trdan S. (2002): *Hercinothrips femoralis* (Reuter) also recorded in Slovenia. *Sodobno kmetijstvo*, 35: 242–244.
- Trdan S., Kužnik L., Vidrih M. (2007): First results concerning the efficacy of entomopathogenic nematodes against *Hercinothrips femoralis* (Reuter). *Acta agriculturae Slovenica*, 89: 5–13.
- Varga L. (2008): *Hercinothrips femoralis* (Reuter, 1891) – a new pest thrips (Thysanoptera: Panchaethripinae) in Slovakia. *Plant Protection Science*, 44: 114–118.
- White W.H. (1916): The sugar-beet thrips: United States Department of Agriculture Bulletin No. 421. In: Howard L.O. (ed): Washington D.C., Government Printing Office: 1–12.

Received: April 4, 2019

Accepted: May 29, 2019

Published online: August 26, 2019