

Dynamics of oak mycorrhizas

V. PEŠKOVÁ

Forestry and Game Management Research Institute, Jíloviště-Strnady, Czech Republic

ABSTRACT: A decline of macromycetes, especially of ectomycorrhizal species, has been described in the last decade in various parts of Europe. These changes are attributed to direct or indirect effects of air pollution. In Central Europe these changes correlate with visual damage of forest. Many ectomycorrhizal fungi seem to be very suitable bioindicators of the disturbance of forest ectotrophic stability. Stages of this disturbance can be linked directly to particular phases of impoverishment of ectomycorrhizal mycocoenoses. These conclusions have been obtained from long-term research on permanent plots (1,000–2,500 m²) in spruce and oak forests in the Czech Republic, but it seems that they are valid generally. For the quantification and study of mycorrhizal activity a special method has been developed and applied. The analyses of mycorrhizas from the same plots in the period 2000–2002 indicate large mycorrhizal dynamics. These data correlate with data obtained from a fruiting bodies survey. Both the percentage of ectomycorrhizal species and the ratio of active mycorrhizas are highly sensitive to outer impacts (air pollution, acidification, fertilization). Their decrease is in correlation with the strong defoliation of trees and can be used for the prediction of further development in comparable stands.

Keywords: ectomycorrhizas; ectomycorrhizal fungi; *Quercus* – oak; root; forest stability

In the last decades a large decline of the tree health status has been observed in Europe. Different studies did not indicate any simple reason for it. Direct impacts of pathogens and poisoning agents are combined with probably more important dysfunction of the nutritional transfer process. In general, it is a synergic effect of many factors (THOMAS et al. 2002) but their particular involvement cannot be clearly detected and their individual role is still discussed. Among others, great attention is focused on the study of root systems and the role of symbiotic fungi.

Mycorrhizal associations

Mycorrhizas are specific forms of symbiosis between plants and fungi. This phenomenon often described as mycorrhizal infection is in fact a complex process. It is known from more than 95% of plant species. Mycorrhizal associations are less often among water plants and they were not detected on many species from ruderal and boggy stands.

Experiments proved that plants with appropriate levels of mycorrhizas revealed increased uptake of nutrients (mainly phosphorus and nitrogen) especially in situations when these elements were present in low concentrations or in an insoluble form. Another important effect is connected with their ability to concentrate in their body mineral nutrients that are slowly released and delivered to the symbiotic plant in a period of insufficiency. Reciprocal transport of organic compounds such as monosaccharides ensures a bilateral advantageous effect.

There are three main types of mycorrhizas from the morphological point of view: ectomycorrhizas, endomycorrhizas and ectendomycorrhizas (ROSPAL et al. 2003; PETERSON et al. 2004).

Ectomycorrhizas

Roots of woody plants from the temperate zone form symbioses with species-specific mushrooms. The majority of ectomycorrhizal (ECM) roots reveals a characteristic anatomic structure. Short ECM

Supported by the Ministry of Agriculture of the Czech Republic, Project No. QD 0332.

roots lack root hairs that are characteristic of non-infected roots or roots with endomycorrhizas. They develop predominantly in upper soil layers with high percentage of raw humus.

The surface of roots infected by mycorrhizal fungi is changed being covered by the hyphal mantle. The growth of roots is retarded and the dichotomous branching of root tips is giving them a distinct shape. Hyphae from the mantle often extend to the soil forming here extraradical mycelium and rhizomorphs. In the body of the root, hyphae form Hartig net – a network between root cells. Hartig net is involved in nutrient exchange that was proved by autoradiographic analysis (e.g. BÜCKING, HEYSER 2001). In the period of maximum extent the contact area between host and mushroom hyphae are quite a large interface. The lifespan of ECM roots is dependent on many internal and external factors. The expected maximum lifespan is two years.

Until now, mycorrhizal associations have been identified in about 2,000 plant species. They play an important role in major natural ecosystems, especially in forests as all important woody species (spruce, pine, oak, beech, birch) are characterized by high mycorrhizal activity. It is supposed that about 2,000 species of fungi, mostly *Basidiomycetes*, can form these associations.

Endomycorrhizas

Endomycorrhizal infections do not produce an obvious change. They are not evident with the naked eye. In this case hyphae form the extraradical mycelium and they also grow not only in intercellular areas but also within root cells. Roots are not altered having the normal surface, structure and root hairs. Finely branched intraradical hyphae constitute “arbuscules” responsible for nutrient exchange and enlarged “vesicles” with storage function (MEJSTŘÍK 1988).

Endomycorrhizas are known from about 1,000 plant genera of 200 families but it is supposed that this association is important for much more, perhaps about 300,000 species. On the other hand, the number of involved fungi is very low. These fungi belong to *Zygomycetes* (ROSYPAL et al. 2003).

Ectendomycorrhizas

Ectendomycorrhizas resemble ectomycorrhizas. They have the mantle and Hartig net but after Hartig net formation, intracellular hyphae develop into epidermal and cortical cells. They are formed between a limited number of ascomycetous fungi and the conifer genera *Pinus* and *Larix*.

Reasons for decline of forests

According to a majority of authors, the decline of forests in Central Europe is caused by the synergic influence of abiotic and biotic factors. A list of abiotic stress factors comprises: repeated dry periods, limited precipitation in sensitive periods and in general non-optimal distribution of rains, deep winter frosts in some years and mild winters with lack of necessary extent of dormancy, winters without snow cover and fast dramatic changes of weather. Important factors are also long-term changes of site conditions connected with climatic changes (fast water drain and deficit of soil humidity, decrease of water level), anthropogenic changes (emissions followed by acidification, soil wash out processes, deposition of toxic emissions and general changes of soil chemistry) or direct anthropic influence (direct devastation, false husbandry strategy or ecologically incorrect decisions). Other factors are connected with high density of game animals followed by great winter damage by grazing and deer barking with subsequent increase in fungi infections or insect infestations. Trees under attack much more likely reduce the growth of roots and thus mycorrhizal associations are also failing (FELLNER, PEŠKOVÁ 1995).

MATERIAL AND METHODS

Dynamics of mycorrhizas on oak study plots

Sampling strategy

Fine roots are concentrated in the depth of 8–30 cm, mainly close to litter and humus horizons. MARKS et al. (1967) and ALEXANDER (1985) calculated the optimum volume for borehole samples. The diameter of about 4 cm is enough when only roots smaller than 2 mm are significant. The sample size used in different studies fluctuates from 1.2 to 10 cm of diameter, and sampling depth from 7.5 to 90 cm. Larger samples can contain a more representative composition, but the labour costs of root preparation and analyses grow significantly. We use the cylinder soil probe 6 cm in diameter and 15 cm in height. This seems to be the optimum between sample size and acceptable time for root evaluation. Five samples were taken from each study plot in order to prevent accidental deviations. They are scattered not randomly, but regularly at the distance of 1 m from selected trees. With standard sampling strategy (PEŠKOVÁ 2000) we obtained in total 210 samples.

In the early 90ties, a standard quantification method was developed for comparative studies of mycorrhizas (PEŠKOVÁ 2000). From soil samples

Table 1. Short characteristic of oak study plots

Working-plan area	Březka 1 114201	Březka 2 114201	Dřevíč 1 112000	Dřevíč 2 112000	Postoloprty 406000	Třebotov 114701
Organ. unit	Březka game preserve	Březka game preserve	LČR, Nižbor Forest Dist.	LČR, Nižbor Forest Dist.	LČR, Žatec Forest Dist.	Lesy Steinských
Stand	2C12	3A13	715D7	714G8	506B11	017J12
Forest region	10	10	8a	8a	2b	8
Forest type	3K6	3S1	2B4	2B4	1B5	2S2
Age	113	122	160	110	109	117
Stand density	7	7	7	8	10	10
Oak proportion (%)	94	100	80	95	59	97
Height a.s.l. (m)	425	430	430	390	260	350
Location	14°32'E, 49°54'N	14°32'E, 49°54'N	13°58'E, 50°01'N	13°58'E, 50°01'N	13°47'E, 50°22'N	14°18'E, 49°58'N

taken constantly by the soil probe (400 cm³) all root fragments were extracted and, besides others, their number, length and mycorrhizas were surveyed. Evaluation of ectomycorrhizal infection was repeatedly performed on selected study plots together with scoring of the health status of trees (defoliation) and quantification of all fungi fructifications (*Macromycetes*). Chosen plots represent different ecological environments. Sampling was performed in the period of mycorrhizal growth maximum (spring, autumn).

Preparation of root samples and their evaluation

From each probe all roots were separated manually, the total remaining soil was washed out. All roots were classified into four root classes according to root diameters (to 1 mm, 1–2 mm, 2–5 mm and +5 mm), and for detailed analyses only the thinnest roots (to 1 mm of diameter) were stored in the fixa-

tion solution (glutaraldehyde). The root dry matter and the total (cumulated) length were measured for roots from all four classes.

Evaluation of mycorrhizal infection

The main analyzed factors were absolute and relative numbers of active and non-active mycorrhizal tips. Mycorrhizas were evaluated within the root fraction under 1 mm. We applied the modified method of JAKUCS et al. (1986) using 20 root sections of 5 cm in length. The level of mycorrhizal infection was evaluated using two parameters: density and percentage of mycorrhizal tips. The density of active and non-active mycorrhizas was counted as an average value of the number of mycorrhizal tips connected with 1 cm of root. The percentage proportion of mycorrhizas was calculated as a ratio of active and non-active mycorrhizal tips (VOGT et al. 1983; CAISOVÁ 1994).




Fig. 1. Relative quantity of active mycorrhizas on oak roots between 2000 and 2002

Table 2. Sampling for root mycorrhizal analyses

Plot	Sample	Tree No	Date
Březka 1	1-5	49, 16, 19, 12, 31	02/05/2000
	1a-5a	49, 16, 19, 12, 31	23/04/2001
	1b-5b	49, 16, 19, 12, 31	12/10/2000
	1c-5c	49, 16, 19, 12, 31	18/10/2001
Březka 2	1-5	7, 17, 36, 43, 45	02/05/2000
	1-5	2, 38, 26, 15, 10	23/04/2001
	1a-5a	2, 38, 26, 15, 10	12/10/2000
	1b-5b	2, 38, 26, 15, 10	18/10/2001
Dřevíč 1	1-5	2, 38, 26, 15, 10	04/05/2000
	1a-5a	2, 38, 26, 15, 10	25/04/2001
	1b-5b	2, 38, 26, 15, 10	06/10/2000
	1c-5c	2, 38, 26, 15, 10	16/10/2001
Dřevíč 2	1-5	28, 30, 38, 35, 43	04/05/2000
	1-5	1, 7, 28, 33, 49	25/04/2001
	1-5	1, 12, 18, 21, 34	06/10/2000
	1-5	1, 12, 18, 21, 34	16/10/2001
Postoloprty	1-5	1, 7, 28, 33, 49	16/06/2000
	1-5	1, 12, 18, 21, 34	11/10/2000
	1-5	1, 12, 18, 21, 34	06/10/2000
	1-5	1, 12, 18, 21, 34	10/10/2001
Třebotov	1-5	1, 12, 18, 21, 34	07/06/2000
	1-5	1, 12, 18, 21, 34	15/05/2001
	1-5	1, 12, 18, 21, 34	14/05/2002
	1-5	1, 12, 18, 21, 34	14/10/2002

RESULTS

Comparison of root development and mycorrhizas in 2000–2002

Comparisons of the main studied parameters, i.e. densities and relative numbers of active (AM) and non-active (NM) mycorrhizas, show large differences from year to year. In 2000 the best situation was found in Dřevíč 1 with average level of AM density and very low NM density. The lowest density and percentage of AM were found in Postoloprty. In the next year in 2001 we found the highest AM quality in Dřevíč 2 and the worst quality in Postoloprty again. The highest increase in AM was observed on plots that had lower values in the previous year (Dřevíč 2, Postoloprty, Třebotov). In 2002 the highest levels of AM (both density and percentage) were detected in Dřevíč 2 again and lowest in Postoloprty and Březka.

It seems that convenient weather conditions, i.e. higher temperatures and more rains, were positively correlated with all mycorrhizal parameters. There was an increase in AM percentages on all oak study plots except for Postoloprty in 2002 (Figs. 1 and 2) and also an increase in AM densities at all sites (Figs. 3 and 4). Average values of AM calculated from all study plots were increasing gradually and the values of NM decreased correspondingly between 2000 and 2002. Average values of dry roots (below 1 mm) were constant (Fig. 5).


The following comparison and statistical tests of AM percentages and densities were performed in the statistic package Kwikstat.

Percentage of active mycorrhizas (2000–2002):

2000: mean = 17.5	s. d. = 9.14032	n = 12
2001: mean = 34.75	s. d. = 13.15035	n = 12
2002: mean = 43.91667	s. d. = 16.3788	n = 12

Analysis of variance:

Source	S.S.	DF	MS	F	Appx P
Total	10,089.89	35			
Treatment	4,317.72	2	2,158.86	12.34	0.000
Error	5,772.17	33	174.91		






Fig. 2. Relative quantity of non-active mycorrhizas on oak roots between 2000 and 2002


Densities of active mycorrhizas (2000–2002):

2000: mean = 0.405 s. d. = 0.1828313 n = 12
 2001: mean = 0.7091667 s. d. = 0.228173 n = 12
 2002: mean = 1.059167 s. d. = 0.4714091 n = 12

Analysis of variance:

Source	S.S.	DF	MS	F	Appx P
Total	5.96	35			
Treatment	2.57	2	1.29	12.54	0.000
Error	3.38	33	0.10		

	Minimum	25 th %tile	Median	75 th %tile	Maximum
0	0.18	0.27	0.32	0.61	0.71
1	0.42	0.53	0.66	0.89	1.11
2	0.39	0.58	1.13	1.48	1.67



We found statistically significant differences between years in percentage and also in densities of active mycorrhizas.

Comparison of macromycete diversity based on their fructifications observed between 2000 and 2002.

The macromycete maximum was found on plot Dřevíč 1 (128 species) and Dřevíč 2 (126 species). The plot of Třebotov was also very rich (107 spe-




Fig. 3. Density of active mycorrhizas on oak roots between 2000 and 2002




Fig. 4. Density of nonactive mycorrhizas on oak roots between 2000 and 2002




Fig. 5. Comparison of root dry matter (root section to 1mm in diameter)

cies). Other study plots like Březka 1, Březka 2 and Postoloprty reveal low diversity with 84, 54 and 56, respectively. We observed a gradual increase in the species number.

Evaluation of defoliation between 2000 and 2003

The health status of oak (*Q. petraea* and *Q. robur*) forests on study plots was evaluated using a standard scoring method. In this period we observed improvement of defoliation (Fig. 6). This amelioration is most probably a result of very convenient weather conditions with high level of precipitation in 2001 and 2002.

DISCUSSION

Our research on mycorrhizas in broad-leaved and coniferous forest on permanent plots produced

partly expected results and also indicated some interesting trends. For example, we confirmed the observation of JANSEN (1991) from Douglas fir stands in the Netherlands that the proportion of active mycorrhizal tips was positively, and mostly very closely, correlated with the proportion of mycorrhizal species, and negatively correlated with the proportion of trees with a severe crown decline. However, this observation for beech and oak stands is valid only for trees where crown defoliation exceeds 60% and mostly only in stands with a similar density of mycorrhizas. Similar data were obtained from spruce stands (FELLNER, SOUKUP 1994). The conditions, i.e. similar density of mycorrhizas and severe defoliation seem to be very important factors. Some contradictory results, e.g. CAUSIN et al. (1994), could perhaps be explained by differences between compared stands.




Fig. 6. Changes of oak defoliation on study plots between 2000 and 2003

Our data on the dry matter of roots indicates that the finest roots, i.e. those < 1 mm in diameter, are generally the most sensitive to air pollution, fertilization or other external impacts (FELLNER 1993). However, the results from our plots are not clear-cut, and only indicate trends. Analyses of data from individual trees may give a clearer picture than data from entire plots.

The fluctuation of the level of mycorrhizal infection reveals a clear correlation between the tree health status and absolute or relative numbers of active mycorrhizas. A conspicuous decline of macromycetes, especially in ectomycorrhizal species, that was described in our study (FELLNER, PEŠKOVÁ 1995) and also in the last decade in various parts of the temperate zone of Europe can be attributed to direct or indirect effect of air pollution. As a collateral effect it has an impact on nutrient budgets of oak forests. This can be documented by worse development of mycorrhizas, roots (lower biomass) or finally also by a defoliation of trees.

In view of the fact that in Central Europe these changes preceded the visual damage of forest or even its decline, many ectomycorrhizal fungi seems to be very suitable bioindicators of the disturbance of forest ectotrophic stability. Defined stages of this disturbance can be linked directly to particular phases of impoverishment of ectomycorrhizal mycocoenoses and of enrichment of lignicolous mycocoenoses. These conclusions have been obtained from long-term research on permanent plots 1,000–2,500 m² in spruce and recently also in oak forests in the Czech Republic, but it seems that they are valid generally, at least in temperate forests. The analyses of true mycorrhizas from the same plots confirmed data obtained from fruiting bodies. Both

the percentage of ectomycorrhizal species and the ratio of active mycorrhizas are highly sensitive to some outer impacts on forest ecosystems (air pollution, acidification, fertilization). Their decrease is in negative correlation with the strong defoliation of trees and can be used for the prediction of further development in the locality, at least in comparable stands, i.e. in the case of similar density of active mycorrhizas, and water stress.

CONCLUSIONS

Despite of the fact that data from individual samples can reveal a relatively large variability, the total values are repeatable and they reliably reflect the parameters tested. Changes of mycorrhizal activity observed on oak plots can be described as follows:

- Between the years 2000 and 2002 we observed an evident increase in active mycorrhizas and a corresponding decrease in inactive ones.
- Gradual increase in active mycorrhizas correlates with an improvement of tree health status measured as grade of their defoliation.
- Between the years 2000 and 2002 in the sum of all macromycete fructifications the percentage of mycorrhizal mushrooms increased correspondingly.
- Dry matter of the root fraction to 1 mm in diameter was roughly constant in the studied period.

The results show that there is a link between the quantity of mycorrhizal infection and the health status of trees. It also seems that many ectomycorrhizal fungi (evaluated in their phase of fructification) can be suitable bioindicators of forest ectotrophic situation. The decline of ectomycorrhizal species can

indicate disturbances and subsequently decline of oak forests.

The evaluations of various root and mycorrhizal parameters indicate that the density of mycorrhizal tips is affected by the long-term quality of local soil conditions while the proportion of active mycorrhizal tips is a more sensitive indicator of present factors such as drought stress, air pollution, use of fertilizers, etc. Despite of these general correlations it is not yet completely known what factors directly influence the mycorrhizas and how they affect their development.

References

ALEXANDER I., 1985. Measuring mycorrhizal biomass: principles and problems. In: Proceedings of the 6th NACOM, June 25–29, 1984, Bend, Oregon, U.S.A. Forest Research Laboratory: 136–138.

BÜCKING H., HEYSER W., 2001. Microradiographic localization of phosphate and carbohydrates in mycorrhizal roots of *Populus tremula* × *Populus alba* and the implication for transfer processes in ectomycorrhizal associations. *Tree Physiology*, 21: 101–107.

CAISOVÁ V., 1994. Hodnocení mykorrhizace sazenic smrku pichlavého (*Picea pungens*) po aplikaci tekutého a suchého inokula lakovky (*Laccaria proxima*). *Práce VÚLHM*, 79: 117–123.

CAUSIN R., MONTECCHIO L., MUTTO ACCORDI S., 1994. The mycorrhizal status of a declining wood of *Quercus robur* L. In: Environmental constraints and oaks: Ecological and physiological aspects. Symposium INRA – IUFRO – EUROSILVA, August 29–September 1, Velaine en Haye, Nancy, France. Centre National de Formation Forestiere, Velaine en Haye: 141.

FELLNER R., 1993. Air pollution and mycorrhizal fungi in Central Europe. In: PEGLER D.N., BODDY L.B., KIRK P.M. (eds.), *Fungi of Europe: Investigation, Recording and Conservation*. Kew, Royal Botanic Gardens: 239–250.

FELLNER R., PEŠKOVÁ V., 1995. Effects of industrial pollutants on ectomycorrhizal relationships in temperate forests. *Canadian Journal of Botany*, 73 (Suppl. 1): 1310–1315.

FELLNER R., SOUKUP E., 1994. Mycological monitoring in the air polluted regions of the Czech Republic. *Communications Instituti Forestalia Čechosloveniae*, 17: 125–137.

JAKUCS P., MESZAROS I., PAPP B.L., TOTH J.A., 1986. Acidification of soil and decay of sessile oak in the „Sikfokaj project“ area (N-Hungary). *Acta Botanica Hungarica*, 32: 303–322.

JANSEN A.E., 1991. The mycorrhizal status of Douglas fir in the Netherlands: its relation with stand age, regional factors, atmosphere pollutants and tree vitality. *Agriculture, Ecosystems and Environment*, 35: 191–208.

MARKS G.C., DITCHBURNE N., FOSTER R.C., 1967. A technique for making quantitative estimates of mycorrhiza populations in *Pinus radiata* forest and its application in Victoria. In: Proceedings of the XIVth IUFRO Congress, September 4–9, 1967. Germany, Munich: 66–83.

MEJSTŘÍK V., 1988. *Mykorrhizní symbiózy*. Praha, Academia: 150.

PEŠKOVÁ V., 2000. Odběry a rozborů kořenů pro studium mykorrhizních poměrů v lesních porostech. *Zpravodaj ochrany lesa*, VI (8): 9–10.

PETERSON R.L., MASSICOTTE H.B., MELVILLE L.H., 2004. *Mycorrhizas: Anatomy and Cell Biology*. Ottawa, NRC Research Press: 173.

ROSYPAL S. et al., 2003. *Nový přehled biologie*. Praha, Scientia: 797.

THOMAS F.M., BLANK R., HARTMANN G., 2002. Abiotic and biotic factors and their interactions as causes of oak decline in Central Europe. *Forest Pathology*, 32: 277–307.

VOGT K.A., MOORE E.E., VOGT D.J., REDLIN M.J., EDMONDS R.L., 1983. Conifer fine root and mycorrhizal root biomass within the forest floors of Douglas-fir stands of different ages and site productivities. *Canadian Journal of Forest Research*, 13: 429–437.

Received for publication December 28, 2004

Accepted after corrections January 20, 2005

Dynamika dubových mykorrhiz

V. PEŠKOVÁ

Výzkumný ústav lesního hospodářství a myslivosti, Jíloviště-Strnady, Česká republika

ABSTRAKT: V práci jsou uvedeny výsledky výzkumu ze šesti trvalých zkusných dubových ploch v ČR (Březka 1 + 2, Dřevíč 1 + 2, Postoloprty, Třebotov). Na vybraných plochách byly provedeny odběry kořenů a mykorrhiz, hodnocen zdravotní stav dubů, sbírání plodnice makromycet. Z výsledků rozborů kořenů a mykorrhiz je od r. 2000 do r. 2002 patrný zřetelný nárůst aktivních mykorrhiz

a naopak pokles neaktivních mykorhiz. Tento postupný nárůst aktivních mykorhiz koresponduje s pozvolným zlepšováním zdravotního stavu stromů (hodnoceným především stupněm defoliace). Srovnání údajů procentuálního zastoupení mykorhizních hub na sledovaných dubových plochách ukazuje zvyšující se podíl mykorhizních hub v celkovém druhovém spektru makromycet.

Klíčová slova: ektomykorhiza; ektomykorhizní houba; *Quercus* – dub; kořen; stabilita lesa

V posledních desetiletích došlo v Evropě ke zhoršení zdravotního stavu lesních porostů. Příčiny nelze jednoznačně určit. Vedle přímého působení patogenních organismů a znečištění prostředí zde má hlavní úlohu i narušení nutričního mechanismu. Velká pozornost je v současnosti věnována výzkumu kořenových systémů a funkci symbiotických hub.

Ve druhé polovině osmdesátých let publikovali v Maďarsku výsledky mnohaletých výzkumů o ekologických souvislostech odumírání dubů v Maďarsku. Při interpretaci svých závěrů zvažovali vlivy nejrozličnějších činitelů, které mohou vést k poškozování lesních porostů, i nejrozličnější projevy negativních změn, které lze v poškozovaných lesních ekosystémech zjistit. Scénář základních příčinných vztahů v tomto složitém procesu degradace lesních stanovišť vychází z klíčové role vzdušného znečištění, vedoucí k zániku mykorhizních hub, což ve svých důsledcích navozuje procesy odumírání kořínků a inhibici mykorhiz.

Mykorhizní houby mohou sloužit jako bioindikátory narušení ektotrofní stability lesa. Dosavadní výsledky výzkumu mykorhiz ukazují na diagnostický význam stanovení procentuálního podílu mykorhizních druhů makromycet vzhledem k nemýkorhizním druhům.

Studium biomasy jemných kořenů je důležitým parametrem pro poznání dynamiky lesních ekosystémů. Dynamika rozvoje jemných kořenových systémů a ektomykorhiz je řízena jednak vnitřními faktory dřeviny, podmínkami půdního prostředí (dostupností vody, aciditou, dostupností minerálních látek, obsahem organické hmoty v půdě atd.) a povětrnostními vlivy.

Během výzkumu mykorhiz na dubových plochách bylo kořenovou sondou v letech 2000–2002 odebí-

ráno pět vzorků, celkem bylo získáno 210 vzorků kořenů a mykorhiz. Odběry kořenů a mykorhiz byly realizovány v období růstového maxima mykorhiz (na jaře a na podzim), a to vždy na stejném místě.

Úroveň mykorhizních poměrů byla hodnocena s využitím dvou parametrů: hustota mykorhizních špiček a jejich procentuální podíl. Hustota aktivních a neaktivních mykorhiz je počítána jako průměrná hodnota zjištěného počtu mykorhiz vztažená na 1 cm délky kořene. Procentuální podíl mykorhiz je kalkulován jako poměr aktivních a neaktivních mykorhiz.

V průběhu let 2000–2002 byly na těchto dubových plochách sbírány makromycety jednou až dvakrát měsíčně v období od června do listopadu podle průběhu počasí a růstu hub. Nasbíraný materiál byl determinován, byla stanovena jeho trofická příslušnost (zjednodušeně: mykorhizní a saprotrofní – ostatní – druhy hub).

Hodnocení zdravotního stavu dubů bylo v letech 2000–2003 zjišťováno posouzením defoliace korun. Výzkum přinesl tyto výsledky:

- Celkově je od r. 2000 do r. 2002 patrný zřetelný nárůst aktivních mykorhiz a naopak pokles neaktivních mykorhiz.
- Postupný nárůst aktivních mykorhiz koresponduje s pozvolným zlepšováním zdravotního stavu stromů (hodnoceným stupněm defoliace).
- Srovnání údajů procentuálního zastoupení mykorhizních hub na sledovaných dubových plochách v r. 2000–2002 ukazuje zvyšující se podíl mykorhizních hub v celkovém druhovém spektru makromycet.
- Výsledky analýzy sušiny kořenů v r. 2000–2002 vykazují celkem vyrovnané hodnoty, bereme-li v úvahu nejcitlivější frakci do 1 mm.

Corresponding author:

Ing. VÍTĚZSLAVA PEŠKOVÁ, Výzkumný ústav lesního hospodářství a myslivosti, Jíloviště-Strnady,
156 04 Praha 5-Zbraslav, Česká republika
tel.: + 420 257 892 299, fax: + 420 257 920 648, e-mail: peskova@vulhm.cz
