

Immunohistochemical characterisation of cholinergic nerve fibres supplying accessory genital glands in the pig

M. KLIMCZUK, J. KALECZYC, A. FRANKE-RADOWIECKA, K. CZAJA, P. PODLASZ, M. LAKOMY

Department of Animal Anatomy, Faculty of Veterinary Medicine, University of Warmia and Mazury, Olsztyn, Poland

ABSTRACT: Our previous immunohistochemical investigations revealed three major populations of nerve fibres supplying the porcine accessory genital glands (AGG) including noradrenergic, non-noradrenergic putative cholinergic and sensory nerve terminals (Kalczyk et al., 1997). However, it is still unclear whether the non-noradrenergic nerve fibres are cholinergic in nature. The knowledge of the population of cholinergic nerve fibres in mammalian AGG based upon vesicular acetylcholine transporter (VACHT) immunohistochemistry is very limited. Therefore, the aim of the present investigation was to disclose the occurrence and colocalization patterns of VACHT, dopamine β -hydroxylase (D β H) and some neuropeptides including vasoactive intestinal polypeptide (VIP), neuropeptide Y (NPY) and somatostatin (SOM) within nerve fibres supplying the porcine AGG. Double-immunohistochemical labellings showed that VACHT-positive nerve terminals were non-adrenergic (D β H-negative), however, many of them contained immunoreactivities to VIP, NPY and/or SOM. The coexistence patterns of these biologically active substances in nerve fibres supplying particular glands are similar but the density of cholinergic innervation varies between the organs. The innervation of the seminal vesicle and prostatic body is better developed than that of the disseminated part of the prostate and bulbourethral glands. The majority of cholinergic nerve fibres associated with blood vessels supplying the glands contain VIP and NPY and, to a lesser degree, SOM. The possible function and origin of the cholinergic nerve fibre population are discussed.

Keywords: accessory genital glands; cholinergic innervation; neuropeptides; immunohistochemistry; pig

In the boar, accessory genital glands include the body and disseminated part of the prostate, paired seminal vesicles and bulbourethral glands. These glands receive an abundant innervation arising primarily from pelvic ganglia and, to a lesser extent, from the inferior mesenteric ganglion as well as from the sympathetic chain and dorsal root ganglia (DRG) (Sjostrand, 1965; Costa and Furness, 1973; Kalczyk et al., 1993, 1994, 2002; Kolbeck and Steers, 1993; Dhami and Mitchell, 1994; Danuser et al., 1997).

Two groups of autonomic nerves are thought to control the secretory function of AGG (Dail, 1993). The expulsion of glandular excretions dur-

ing seminal emission is thought to be largely under sympathetic-noradrenergic control. Noradrenergic nerves induce contractions of smooth muscle cells surrounding the secretory epithelium. Sympathetic non-adrenergic (putative cholinergic) nerves are believed to be secretomotor to the glandular epithelium.

It has been demonstrated in a variety of species including the rat, mouse, hamster, guinea-pig, cat, rabbit, dog and humans that nerves innervating AGG, in addition to classical neurotransmitters such as acetylcholine and noradrenalin, may also contain other biologically active substances including substance P (SP), met-enkephalin (Met-ENK),

leu-enkephalin (leu-ENK), neuropeptide Y (NPY), vasoactive intestinal peptide (VIP), somatostatin (SOM), calcitonin gene-related peptide (CGRP) and nitric oxide (Alm et al., 1980, 1981; Stjernquist et al., 1987; Aumuller et al., 1989; Higgins and Gossling, 1989; Crowe et al., 1991; Kepper and Keast, 1995; Jen et al., 1995; Tainio, 1995; Hedlund et al., 1996; Chow et al., 1997; Sjostrand et al., 1998).

The knowledge of neurochemical properties of adrenergic nerve fibres supplying mammalian AGG is comprehensive and detailed because methods used to investigate adrenergic nerve structures are well developed and considered to be reliable. On the other hand, until quite lately, the exact identification of peripheral cholinergic nerve structures was impossible because specific markers to investigate this part of the peripheral nervous system (PNS) were not available. The most commonly used histochemical method for localization of acetylcholinesterase (AChE) is thought to be controversial in assessment of cholinergic nerve fibres because this enzyme has been found to be related not exclusively to cholinergic axons but also to nerve terminals belonging to other subdivisions of the PNS (Koelle, 1955; Burn and Rand, 1965; Jacobovitz and Koelle, 1965). Recently, efficient antibodies against an isoform of choline acetyltransferase (ChAT; an enzyme of acetylcholine synthesis) found in the peripheral nerve structures were raised and characterised, and using these antisera the presence of ChAT-containing nerve fibres in the wall of the vas deferens was revealed (Kujat et al., 1993; Kaleczyc et al., 2000). Moreover, an alternative marker for cholinergic nerve structures has become available in the form of an antibody to vesicular acetylcholine transporter (VACHT – the protein associated with the storage of acetylcholine within vesicles in cholinergic nerve terminals). VACHT-positive nerve fibres were found in the human prostate and seminal vesicle (Dixon et al., 2000) and in the rat prostate (Schafer et al., 1998; Nadelhaft, 2003). However, the knowledge of neurochemical properties of cholinergic nerve fibres supplying AGG is still very limited. Therefore, the present study was designed to investigate the presence and coexistence of VACHT and other biologically active substances including NPY, VIP, SOM (neuropeptides found earlier to occur most frequently in non-adrenergic nerve fibres supplying the porcine AGG; Kaleczyc et al., 1999) and D β H in nerve terminals supplying the porcine AGG using single and double-labelling immunofluorescence.

MATERIAL AND METHODS

The experiment was performed on three juvenile (12–15 kg body weight, b.w.) and three adult (90–100 kg b.w.) boars of the Large White Polish race obtained from a commercial fattening farm in Purda (Poland). All the animals were housed and treated in accordance with the rules approved by the local Ethics Commission (conforming the principles of Laboratory Animal Care, NIH publication No. 86-23, revised in 1985).

Removal, fixation and sectioning of AGG. Prior to perfusion, the pigs were deeply anaesthetised. Thirty minutes before the main anaesthetic pentobarbital (Vetbutal, Biowet, Poland; mg/kg b.w.) was applied intravenously, all the animals were pretreated with propionylpromazine (Combelen; Biowet, Poland 0.7 mg/kg b.w., *i.m.*). Then they were transcatheterially perfused with 0.5 l (juvenile animals) or 1 l (adult boars) of preperfusion solution containing 0.9% sodium chloride (Chemia, Gliwice, Poland), 2.5% polyvinylpyrrolidone (Sigma, Deisenhofen, Germany), 0.5 procaine hydrochloride (Polfa, Warsaw, Poland) and 20 000 IU of heparin (Heparinum; Polfa; added *ex tempore*), followed by 4 l (juvenile animals) or 8 l (adult boars) of 4% ice-cold buffered paraformaldehyde (pH 7.4).

Samples of bulbourethral glands, seminal vesicles and prostates (including the body and disseminated part) were dissected out and postfixed by immersion in the same fixative for 2 hours, then rinsed with phosphate buffer (pH 7.4) and transferred to and stored in 18% buffered sucrose solution (pH 7.4) until further processing. Eight- μ m cryostat sections from the glands were cut longitudinally or transversally. All the sections were mounted on chrome alum-gelatine-coated glass slides.

The sections were processed for single- or double-labelling immunohistochemistry to study the presence and coexistence of VACHT with other biologically active substances including D β H, VIP, NPY and SOM in nerve fibres supplying the glands.

Immunohistochemical procedure. The sections were air-dried at room temperature (RT) for 30 min, rinsed (3 \times 5 min) with phosphate-buffered saline (PBS; pH 7.4), preincubated with a blocking mixture containing 0.25% Triton X-100, 1% bovine serum albumin and 10% normal goat serum in PBS (1 h, room temperature, RT), and incubated overnight (ON) in a humid chamber with one (single labellings) or a mixture of two (double-labellings) primary antisera raised in different

species. Afterwards, in the case of single-labellings, the sections were incubated with an appropriate biotinylated antiserum (1h, RT, for single- and double-labellings this step was sometimes omitted) followed by incubation with CY-3-conjugated streptavidin or an appropriate fluorescein isothiocyanate (FITC)-conjugated secondary antiserum (1h, RT); in the case of double-stainings, mixtures of the appropriate secondary reagents mentioned above were used. Primary antisera and secondary reagents are listed in Table 1. Each step of immunolabelling was followed by raising the sections with PBS (3 × 5 min). Finally, they were mounted with carbonate-buffered glycerol (pH 8.6). The stained sections were studied with Zeiss-Axiophot microscope equipped for epifluorescence and an appropriate filter set for CY3 and FITC.

Controls. Preabsorption of the diluted antiserum with 20 µg/ml of the appropriate antigen (besides DβH) completely abolished the specific immunoreaction. Additionally, the primary antisera were omitted or replaced by non-immune sera or by PBS in order to check the method specificity.

RESULTS

Because no distinct differences were found in the innervation of AGG between the juvenile and adult pigs, the results are described together. Moreover, double-labelling investigations revealed

no colocalization of VACHT and DβH in nerve fibres supplying the glands, thus the nerve terminals immunoreactive to these substances formed two separate populations of axons.

Distribution of VACHT-immunoreactive (VACHT-IR) nerve fibres supplying the seminal vesicles and their chemical coding

Single-labelling immunofluorescence disclosed that the seminal vesicle was intensely supplied with VACHT-positive, varicose nerve fibres. The varicose nerve terminals were often observed to occur beneath but never penetrating into the epithelium. They were also related to smooth muscle cell bundles located in the interstitial tissue of the glands (Figures 1, 2, 3a).

Both intrinsic as well as extrinsic arteries were moderately supplied with VACHT-IR nerve fibres while veins were apposed by solitary nerve terminals only.

Double-labelling immunofluorescence revealed that many VACHT-IR nerve fibres supplying the seminal vesicle contained also NPY, and some of them stained also for VIP or SOM. Most of VACHT-positive nerve terminals located beneath the tubular epithelium displayed immunoreactivities to NPY, VIP or SOM (Figures 1, 2, 3). The vast majority of VACHT-IR fibres found in the interstitial tissue was simultaneously NPY- or VIP-positive

Table 1. List of primary antisera and secondary reagents used in the study

Antigen	Code	Dilution	Species	Supplier
Primary antibodies				
VACHT	H-V006	1 : 8 000	rabbit	Phoenix Pharmaceuticals, USA
NPY	NZ1115	1 : 150	rat	Affiniti, UK
SOM	8330-0009	1 : 50	rat	Biogenesis, UK
VIP	MaVIP	1 : 1 500	mouse	East Acres, USA
Secondary reagents				
Biotinylated anti-rabbit IgG		1 : 400	goat	Dako, Denmark
Biotinylated anti-rat IgG		1 : 400	rabbit	Dako, Denmark
Biotinylated anti-mouse IgG		1 : 400	goat	Dako, Denmark
FITC-conjug. anti-mouse IgG		1 : 400	goat	Jackson Immun. Lab. USA
FITC-conjug. anti-rat IgG		1 : 50	goat	Jackson Immun. Lab. USA
Streptavidin-conjug. C _Y 3		1 : 4 000		Dianova, Hamburg, Germany

Table 2. Coexistence patterns of VAcHT and neuropeptides in nerve fibres supplying the porcine accessory genital glands and semiquantitative evaluation of their relative frequency

Coexistence pattern	Seminal vesicle		Prostatic body		Disseminated prostate		Bulbourethral gland		A	V
	BE	IT	BE	IT	BE	IT	BE	IT		
VAcHT/NPY	++	+++	+++	+	++	+/-	++	+/-	+	+/-
VAcHT/SOM	+	+/-	+/-	+	+	+/-	+	+/-	+/-	+/-
VAcHT/VIP	+	++	++	+	++	+/-	++	+	++	+/-

BE – beneath the epithelium, IT – interstitial tissue, A – arteries, V – veins, – not found, +/- single fibres, + few fibres, ++ moderate number of fibres, +++ many fibres

but only the solitary nerve endings exhibited immunoreactivity to SOM (Figure 3). Most of the perivascular VAcHT-IR nerve fibres stained also for NPY or VIP.

Distribution of VAcHT-IR nerve fibres supplying the prostate and their chemical coding

Single-labelling immunofluorescence revealed that immunoreactivity to VAcHT was expressed in numerous nerve fibres supplying the prostatic body while they were less frequently encountered in the disseminated part of the gland. VAcHT-IR nerve terminals closely apposing glandular tubules

were more numerous than those associated with the smooth muscle cell bundles in the interstitial tissue (Figures 4, 5, 6). Both intrinsic and extrinsic arteries were moderately supplied with VAcHT-positive nerve fibres. Veins were only occasionally apposed by the solitary axons.

Double-immunostaining revealed the presence of immunoreactivity to NPY, VIP and SOM within VAcHT-IR nerve fibres supplying the body and disseminated part of the prostate. Many VAcHT-positive nerve terminals, especially those located beneath the tubular epithelium, contained NPY- and VIP-immunoreactivity (Figures 4, 5), however, only solitary nerve terminals stained also for SOM (Figure 6). The majority of VAcHT-IR perivascular nerve fibres also displayed immunoreactivity to




Figure 1. The seminal vesicle in the juvenile boar. Vesicular acetylcholine transporter (VAcHT)- [a CY3 visualisation (CY3)] and neuropeptide Y-immunoreactive (NPY-IR) [b fluorescein isothiocyanate visualisation (FITC)], nerve fibres (arrows) associated with the glandular epithelium (E) or located within the interstitial tissue (IT) contained both substances studied. In contrast, NPY-positive but VAcHT-negative nerve fibre (arrowhead) was located within the interstitial tissue (IT). Magnification 450×




Figure 2. The seminal vesicle in the juvenile boar. VACHT- (a CY3) and vasoactive intestinal polypeptide (VIP)-IR (b FITC) nerve fibres (arrows) associated with the glandular epithelium (E) or located within the interstitial tissue (IT). Magnification 450×

Figure 3. The seminal vesicle in the juvenile boar. VACHT-IR (a CY3) and somatostatin (SOM)-positive (b FITC) nerve fibres (arrows) associated with the glandular epithelium (E) or interstitial tissue (IT). In contrast, VACHT-positive (a) but SOM-negative (b) nerve fibre (arrowhead) was located within the interstitial tissue (IT). Magnification 450×

NPY and VIP, and the solitary nerve endings were simultaneously SOM-positive.

Distribution of VACHT-IR nerve fibres supplying the bulbourethral gland and their chemical coding

As revealed by single-labelling immunofluorescence, VACHT-positive nerve fibres moderately supplied the bulbourethral gland. These nerve terminals were often found beneath the tubular epithelium while they were less numerous within

the interstitial tissue (Figures 7, 8, 9). Both intrinsic and extrinsic arteries were moderately supplied with VACHT-positive nerve fibres. Veins were only occasionally apposed by the solitary axons.

Double-labelling investigations revealed that many VACHT-IR nerve terminals supplying the bulbourethral gland, especially those located beneath the epithelium, contained also NPY-, VIP- or SOM-immunoreactivity (Figures 7, 8, 9). Most of the perivascular VACHT-containing nerve fibres also displayed immunoreactivity to NPY or VIP and the solitary axons were simultaneously SOM-positive.




Figure 4. The disseminated part of the prostate in the juvenile boar. VACHT-IR (**a** CY3) nerve fibres (*arrows*) associated with the glandular epithelium (E) also contained neuropeptide Y (NPY)-IR [**b** FITC]. Magnification 450×

Figure 5. The disseminated part of the prostate in the juvenile boar. The majority of VACHT-IR (**a** CY3) nerve terminals (*arrows*) supplying glandular acini stained also for VIP (**b** FITC). Magnification 450×

Figure 6. The disseminated part of the prostate of the juvenile boar. VACHT-IR (**a** CY3) and somatostatin (SOM)-positive (**b** FITC) nerve fibres (*arrow*) associated with the glandular epithelium (E). In contrast, VACHT-positive (**a**) but SOM-negative (**b**) nerve fibre (*arrowhead*) was located within the interstitial tissue (IT). Magnification 450×




Figure 7. The bulbourethral gland in the juvenile boar. VACHT- (a CY3) and NPY-IR (b FITC) nerve fibres (arrows) running in the interstitial tissue between the tubules of the gland. In contrast, VACHT-positive (a) but NPY-negative (b) varicose nerve fibre (arrowhead) was located beneath the epithelium of the tubule. Magnification 450×

Figure 8. The bulbourethral gland in the juvenile boar. VACHT- (a CY3) and VIP-IR (b FITC) varicose nerve fibre (arrow) associated with the glandular epithelium. Some VACHT-positive nerve fibres (arrowheads) located within the interstitial tissue (IT) were VIP-negative. Magnification 450×

Figure 9. The bulbourethral gland in the juvenile boar. VACHT- (a CY3) and SOM-IR (b FITC) nerve fibres (arrows) associated with the glandular epithelium (E). Magnification 450×

DISCUSSION

The present study has shown that the porcine AGG are supplied by moderate numbers of nerve fibres containing immunoreactivity to VAcHT, therefore they should be regarded as cholinergic nerve terminals. The VAcHT-positive nerve fibres closely apposing the tubules of the glands are more numerous than those associated with smooth muscle cell bundles in the interstitial tissue. The general distribution of cholinergic nerve terminals in the porcine AGG is similar to that of AChE-positive axons found in other mammals including the hamster, guinea pig, rat and humans (Al-Zuhair et al., 1975; Dunzerdorfen et al., 1976; Vaalasti and Hervonen, 1979; Moss et al., 1987; Higgins and Gosling, 1989; Vega Alvarez et al., 1989; Chow et al., 1997). The literature dealing with the population of cholinergic nerve fibres supplying mammalian AGG based upon VAcHT immunohistochemistry is very limited. The presence of cholinergic (VAcHT-positive) fibres was shown in the human prostate and seminal vesicle (Dixon et al., 2000) and in the rat prostate (Schafer et al., 1998; Nadelhaft, 2003). Dixon et al. (2000) found that the majority of VAcHT-positive nerve endings localized beneath the epithelium of the human prostate and seminal vesicle contained also NPY whereas the small proportion stained for nitric oxide synthase (NOS).

In our recent study dealing with the innervation of the porcine AGG we found that many non-adrenergic nerve fibres were localized beneath the tubular epithelium and in the interstitial tissue (Kaleczyc et al., 1999). These fibres coexpressed immunoreactivities to NPY and VIP, VIP and SOM or to NPY and SOM, thus it is likely that these three peptides coexist within some of the nerve terminals. The present investigations using single- and double-immunofluorescence confirmed these findings and revealed that these non-adrenergic nerve terminals were cholinergic in nature (VAcHT-positive).

The biologically active substances found to coexist within cholinergic nerve fibres supplying porcine AGG probably perform an inhibitory function with regard to their smooth muscle cells and blood vessels. VIP has been demonstrated in nerve fibres supplying the prostate and seminal vesicles in the guinea-pig, rat, cat, rabbit, mouse and humans. VIP-immunoreactive nerve fibres have been mainly associated with the basis of the glandular epithelium and, in smaller proportion, with the smooth muscle coat and blood vessels (Alm et al., 1980;

Vaalasti et al., 1980, 1986; Larsen et al., 1981; Gu et al., 1983; Stjernquist et al., 1983; Polak and Bloom, 1984; Lamano Carvalho et al., 1986; Higgins and Gosling, 1989; Lange and Unger, 1990; Properzi et al., 1992; Jen et al. 1995; Kepper and Keast, 1995; Tainio, 1995). Very little is known about the physiological role of VIP in AGG. Pinho et al. (1994) reported that in the seminal vesicle of the hamster VIP-IR nerve fibres also manifested AChE activity and suggested a modulatory function of this peptide on muscarine receptors. VIP is typically present in cholinergic nerves in other tissues (Ottesen and Fahrenkrug, 1995). On the other hand, there is a widespread agreement that cholinergic nerves are secretomotor to the glandular epithelium of the prostate and seminal vesicles (Dail, 1993). Moreover, normal rat prostate epithelial cells have been shown to contain VIP receptors and to produce cAMP in response to the action of this peptide (Carmena and Prieto, 1985a,b). It is therefore possible, that VIP plays a role in the regulation of the secretory function of epithelial cells in this and other AGG. Immunoreactivity to SOM was found in a few nerve fibres supplying the human seminal vesicle and prostate (Gu et al., 1983; Chapple et al., 1991; Crowe et al., 1991; Tainio, 1995). Mirabella et al. (2003) revealed SOM-IR nerve fibres innervating the prostate in the water buffalo. In this species non-vascular SOM-positive axons contain neither VIP nor TH, thus these fibres probably belong to a subpopulation of non-adrenergic nerves distinct from those containing NOS, VIP and NPY.

The presence of NPY-positive nerve endings in the reproductive organs which receive a dense sympathetic nerve supply such as the vas deferens and accessory sex glands was reported earlier. In the seminal vesicle of man (Adrian et al., 1984; Lange and Unger, 1990; Tainio, 1995) and rodents (Dhami and Mitchell, 1984; Lamano Carvalho et al., 1986; Stjernquist et al., 1987; Yuri, 1990; Properzi et al., 1992; Irvani and Zar, 1994) NPY-immunoreactive nerve fibres appear to be the major peptide-containing neuronal component. These fibres have been predominantly found in the smooth muscle layer of the glands, however, some of them have been detected in the subepithelial tissue and in association with blood vessels. The rich supply of NPY-IR nerve fibres in the non-vascular smooth muscle of accessory sex glands certainly suggests a role in the muscular control of secretory output. Stjernquist et al. (1987) found that NPY largely colocalized with D β H within axons in the guinea-pig seminal

vesicles. This observation was confirmed by Dhami and Mitchell (1994), who found that in this species many pelvic adrenergic neurons projecting to the seminal vesicles, but not to the prostate, contained this peptide. Kepper and Keast (1995), in turn, revealed that many, if not all adrenergic nerve fibres innervating the rat prostate contained NPY.

It is commonly accepted that in mammals, pelvic ganglia are the main source of nerve fibres supplying male reproductive organs (Sjostrand, 1965; Keast, 1992; Kolbeck and Steers, 1993). Kaleczyc et al. (2003) revealed that nearly all non-adrenergic neurons in the porcine APG (Anterior Pelvic Ganglion) were cholinergic in nature, i.e. expressed immunoreactivity for ChAT and VACHT, substances commonly accepted as specific markers of cholinergic nerve structures. These nerve cells co-express simultaneously immunoreactivities to other biologically active substances including NOS, VIP, NPY and SOM. Therefore it is very likely that cholinergic nerve fibres observed in the present study are processes of these neurons. Nadelhaft (2003) also demonstrated small cholinergic (VACHT-positive) neurons in the rat pelvic ganglion supplying the prostate gland.

The literature in the field contains some data dealing with the chemical coding of non-adrenergic pelvic neurons in mammals. It seems to be noteworthy that there are profound species-dependent variations with regard to the expression of biologically active substances in pelvic non-adrenergic nerve cells. Keast (1991) revealed that in the male rat two populations of cholinergic neurons in the major pelvic ganglion (MPG) could be distinguished. The first consisted of cholinergic NPY-positive neurons (stain only for NPY) and the second one was represented by cholinergic VIP-IR neurons (stain only for VIP). No SOM-IR MPG neurons have been found in this species. Thus, in contrast to the pig, NPY- and VIP-positive nerve cell bodies form separate populations of rat male pelvic cholinergic somata that do not contain SOM. In human male infants and children, non-adrenergic pelvic neurons display VIP-, NPY-, NOS-, SOM-, CGRP- and SP-immunoreactivity (Jen et al., 1996). Because the vast majority of non-adrenergic nerve cells in human pelvic ganglia contains VACHT (Dixon et al., 1999), these neurons should be considered as cholinergic.

This is the first comprehensive study dealing with the chemical coding of cholinergic nerve fibres supplying the porcine AGG. These nerve terminals are mostly associated with the tubular epithelium and

contain a variety of colocalized biologically active substances including VACHT as well as NPY, VIP or SOM. Considering a great coincidence between the chemical coding of cholinergic nerve fibres supplying the porcine AGG and that of cholinergic pelvic neurons it is further concluded that pelvic ganglia are the major source of cholinergic innervation for the porcine urogenital system.

Acknowledgements

The authors would like to thank Ms. M. Marczak, G. Greniuk and Mr. A. Penkowski for their skilful technical assistance.

REFERENCES

- Adrian T.E., Gu J., Allen J.M., Tatemoto K., Polak J.M., Bloom S.R. (1984): Neuropeptide Y in the human male genital tract. *Life Science*, 35, 2643–2648.
- Alm P., Alumets J., Hakanson R., Owman C., Sjöberg N.O., Sundler F., Walles B. (1980): Origin and distribution of VIP (vasoactive intestinal polypeptide)-nerves in the genito-urinary tract. *Cell Tissue Research*, 205, 337–347.
- Alm P., Alumets J., Hakanson R., Owman C., Sjöberg N.O., Stjernquist M., Sundler F. (1981): Enkephalin-immunoreactive nerve fibres in the feline genito-urinary tract. *Histochemistry*, 72, 351–355.
- Al-Zuhair A., Gosling J.A., Dixon J.S. (1975): Observations on the structure and autonomic innervation of the guinea-pig seminal vesicles and ductus deferens. *Journal of Anatomy*, 120, 81–93.
- Aumüller G., Jungblut T., Malek B., Konrad S., Weihe E. (1989): Regional distribution of opioidergic nerves in human and canine prostates. *The Prostate*, 14, 279–288.
- Burn J.H., Rand M.J. (1965): Acetylcholine in adrenergic transmission. *Annual Review of Pharmacology*, 5, 163–182.
- Carmena M.J., Prieto J.C. (1985a): Cyclic AMP response to vasoactive intestinal peptide and [beta]-adrenergic or cholinergic agonists in isolated epithelial cells of rat ventral prostate. *Bioscience Reports*, 5, 791–795.
- Carmena M.J., Prieto J.C. (1985b): VIP binding to epithelial cell membranes of rat ventral prostate: effect of guanine nucleotides. *Genetic Pharmacology*, 16, 495–501.
- Chapple C.R., Crowe R., Gosling J., Burnstock G. (1991): The innervation of the human prostate glands – the

- changes associated with benign enlargement. *Journal of Urology*, 146, 1637–1644.
- Chow P.H., Dockery P., Cheung A. (1997): Innervation of accessory sex glands in the adult male golden hamster and quantitative changes of nerve densities with age. *Andrologia*, 29, 331–342.
- Costa M., Furness J.B. (1973): Observations on the anatomy and amine histochemistry of the nerves and ganglia which supply the pelvic viscera and on the associated chromaffin tissue in the guinea-pig. *Zeitschrift für Anatomie Entwicklungsgeschichte*, 140, 85–108.
- Crowe R., Chapple C.R., Burnstock G. (1991): The human prostate glands: a histochemical and immunohistochemical study of neuropeptides, serotonin, dopamine beta-hydroxylase and acetylcholinesterase in autonomic nerves and ganglia. *British Journal of Urology*, 68, 53–61.
- Dail W.G. (1993): Autonomic innervation of male genitalia. In: Maggi C.A. (ed.): *Nervous Control of the Urogenital System*. Harwood, Chur, Switzerland. 69–102.
- Danuser H., Springer J.P., Katofiasc M.A., Thor K.B. (1997): Extrinsic innervation of the cat prostate gland: a combined tracing and immunohistochemical study. *Journal of Urology*, 157, 1018–1024.
- Dhami D., Mitchell B.S. (1994): Chemical coding of neurons projecting to pelvic viscera in the male guinea pig: a study by retrograde transport and immunohistochemistry. *Histochemical Journal*, 26, 262–270.
- Dixon J.S., Jen P.Y.P., Gosling J.A. (1999): Tyrosine hydroxylase and vesicular acetylcholine transporter are coexpressed in a high proportion of intramural neurons of the human neonatal and child urinary bladder. *Neuroscience Letters*, 277, 157–160.
- Dixon J.S., Jen P.Y.P., Gosling J.A. (2000): The distribution of vesicular acetylcholine transporter in human male genitourinary organs and its co-localization with neuropeptide Y and nitric oxide synthase. *Neurology and Urodynamics*, 19, 185–194.
- Dunzerdorfen U., Jonas D., Weber W. (1976): The autonomic innervation of the human prostate. *Histochemistry of acetylcholinesterase in the normal and pathologic states*. *Urological Research*, 4, 29–31.
- Gu J., Polak J.M., Probert L., Islam K.N., Marangos P.J., Mina S., Adrian T.E., McGregor G.P., O'Shaughnessy D.J., Bloom S.R. (1983): Peptidergic innervation of the human male genital tract. *Journal of Urology*, 130, 386–391.
- Hedlund P., Larrson B., Alm P., Andersson K.E. (1996): Nitric oxide synthase-containing nerves and ganglia in the dog prostate: a comparison with other transmitters. *Histochemical Journal*, 28, 635–642.
- Higgins J.R.A., Gosling J.A. (1989): Studies on the structure and intrinsic innervation of the normal human prostate. *The Prostate*, 2 (Suppl.), 5–16.
- Irvani M.M., Zar M.A. (1994): Presence of neuropeptide Y in the seminal vesicles and its effects on noradrenaline- and nerve-induced contractions. *British Journal of Pharmacology*, 113, 877–882.
- Jacobovitz D., Koelle G.B. (1965): Histochemical correlations of acetylcholinesterase and catecholamines in postganglionic autonomic nerves of the cat, rabbit and guinea-pig. *Journal of Pharmacology and Experimental Therapeutics*, 148, 225–237.
- Jen P.Y.P., Dixon J.S., Gosling J.A. (1995): Development of peptide-containing nerves in the human fetal vas deferens and seminal vesicles. *British Journal of Urology*, 75, 378–385.
- Jen P.Y.P., Dixon J.S., Gosling J.A. (1996): Co-localisation of tyrosine hydroxylase, nitric oxide synthase and neuropeptides in neurons of the human postnatal male pelvic ganglia. *Journal of the Autonomic Nervous System*, 59, 41–50.
- Kaleczyc J., Majewski M., Lakomy M. (1993): Immunohistochemical characterization of nerve fibres and pelvic neurons innervating the porcine ductus deferens. *Verhandlungen der Anatomischen Gesellschaft*, 88, 232.
- Kaleczyc J., Timmermans J.-P., Majewski M., Lakomy M., Scheuermann D.W. (1994): NO-synthase-containing neurons of the pig inferior mesenteric ganglion, part of them innervating the ductus deferens. *Acta Anatomica*, 151, 62–67.
- Kaleczyc J., Timmermans J.-P., Majewski M., Lakomy M., Scheuermann D.W. (1997): Immunohistochemical characteristics of nerve fibres supplying the porcine vas deferens. A colocalization study. *Histochemistry and Cell Biology*, 107, 229–241.
- Kaleczyc J., Timmermans J.-P., Majewski M., Lakomy M., Scheuermann D.W. (1999): Immunohistochemical properties of nerve fibres supplying accessory male genital glands in the pig. A colocalisation study. *Histochemistry and Cell Biology*, 111, 217–228.
- Kaleczyc J., Klimczuk M., Wasowicz K., Majewski M., Calka J., Schemann M., Lakomy M. (2000): Immunohistochemical characterization of choline acetyltransferase-containing nerve fibres supplying the vas deferens in the pig. *Polish Journal of Veterinary Sciences*, 3, 111–118.
- Kaleczyc J., Scheuermann D.W., Pidsudko Z., Majewski M., Lakomy M., Timmermans J.-P. (2002): Distribution, immunohistochemical characteristics and nerve pathways of primary sensory neurons supplying the porcine vas deferens. *Cell and Tissue Research*, 310, 9–17.
- Kaleczyc J., Wasowicz K., Klimczuk M., Czaja K., Lakomy M. (2003): Immunohistochemical characterisation of

- cholinergic neurons in the anterior pelvic ganglion of the male pig. *Folia Histochemica et Cytobiologica*, 41, 65–72.
- Keast J. (1991): Patterns of co-existence of peptides and differences of nerve fibre types associated with noradrenergic and non-noradrenergic (putative cholinergic) neurons in the major pelvic ganglion of the male rat. *Cell and Tissue Research*, 266, 405–415.
- Keast J. (1992): Localization and peptide content of pelvic neurons supplying the muscle and *lamina propria* of the rat vas deferens. *Journal of the Autonomic Nervous System*, 40, 1–12.
- Kepper M., Keast J. (1995): Immunohistochemical properties and spinal connections of pelvic autonomic neurons that innervate the rat prostate glands. *Cell and Tissue Research*, 281, 533–542.
- Koelle G.B. (1955): The histochemical identification of acetylcholinesterase in cholinergic, adrenergic and sensory neurons. *Journal of Pharmacology and Experimental Therapeutics*, 114, 167–184.
- Kolbeck S.C., Steers W.D. (1993): Origin of neurons supplying the vas deferens of the rat. *Journal of Urology*, 149, 918–921.
- Kujat R., Rose Ch., Wrobel K.-H. (1993): The innervation of the bovine ductus deferens: comparison of a modified acetylcholinesterase-reaction with immunoreactivities of cholinacetyltransferase and panneuronal markers. *Histochemistry*, 99, 231–239.
- Lamano Carvalho T.L., Hodson N.P., Blank M.A., Watson P.F., Mulderry P.K., Bishop A.E., Gu J., Bloom S.R., Polak J.M. (1986): Occurrence, distribution and origin of peptide-containing nerves of guinea-pig and rat male genitalia and the effects of denervation on sperm characteristics. *Journal of Anatomy*, 149, 121–141.
- Lange W., Unger J. (1990): Peptidergic innervation of within the prostate glands and seminal vesicles. *Urological Research*, 18, 337–340.
- Larsen J.J., Ottesen B., Fahrenkrug J., Fahrenkrug L. (1981): Vasoactive intestinal polypeptide (VIP) in the male genitourinary tract: concentration and motor effects. *Investigative Urology*, 19, 211–213.
- Mirabella N., Squillacioti C., Varricchio E., Genovese A., Paino G. (2003): Innervation of vas deferens and accessory male genital glands in the water buffalo (*Bubalus bubalis*). *Neurochemical characteristics and relationships to the reproductive activity. Theriogenology*, 59, 1999–2016.
- Moss H.E., Crowe R., Burnstock G. (1987): The seminal vesicle in eight week streptozotocin-induced diabetic rats: Adrenergic, cholinergic and peptidergic innervation. *Journal of Urology*, 138, 1273–1278.
- Nadelhaft I. (2003): Cholinergic axons in the rat prostate and neurons in the pelvic ganglion. *Brain Research*, 989, 52–57.
- Ottesen B., Fahrenkrug M.D. (1995): Vasoactive intestinal polypeptide and other preprovasoactive intestinal polypeptide-derived peptides in the female and male genital tract: Localization, biosynthesis, and functional and clinical significance. *American Journal of Obstetrics and Gynecology*, 172, 1615–1631.
- Pinho M.S., Sebastiao A.M., Rodrigues G., Barroso C.P., Ribeiro J.A., Mata L.R., Gulbenkian S. (1994): Vasoactive intestinal peptide in the hamster seminal vesicle: distribution, binding sites and possible functions. *Neuroscience*, 598, 1083–1091.
- Polak J.M., Bloom S.R. (1984): Localization and measurement of VIP in the genitourinary system of man and animals. *Peptides*, 5, 225–230.
- Properzi G., Cordeschi G., Francavilla S. (1992): Post-natal development and distribution of peptide-containing nerves in the genital system of the male rat. *Histochemistry*, 97, 61–68.
- Schafer M.K.H., Eiden L.E., Weihe E. (1998): Cholinergic neurons and terminal fields revealed by immunohistochemistry for the vesicular acetylcholine transporter. II. The peripheral nervous system. *Neuroscience*, 84, 361–376.
- Sjostrand N.O. (1965): The adrenergic innervation of the vas deferens and the accessory male genital glands. *Acta Physiologica Scandinavica*, 65 (Suppl. 257), 1–82.
- Sjostrand N.O., Ehren I., Eldh J., Wiklund N.P. (1998): NADPH-diaphorase in glandular cells and nerve fibres and its relation to acetylcholinesterase-positive nerves in the male reproductive tract of man and guinea-pig. *Urological Research*, 26, 181–188.
- Stjernquist M., Hakanson R., Leander S., Owman C., Sundler F., Uddman R. (1983): Immunohistochemical localization of substance P, vasoactive intestinal polypeptide and gastrin-releasing peptide in the vas deferens and seminal vesicle, and the effect of these and eight other neuropeptides on resting tension and neurally evoked contractile activity. *Regulatory Peptides*, 7, 67–86.
- Stjernquist M., Owman C., Sjoberg N.O., Sundler F. (1987): Coexistence and cooperation between neuropeptide Y and norepinephrine in nerve fibres of guinea-pig vas deferens and seminal vesicles. *Biology of Reproduction*, 36, 149–155.
- Tainio H. (1995): Peptidergic innervation of the human prostate, seminal vesicles and vas deferens. *Acta Histochemica*, 97, 113–119.
- Vaalahti A., Hervonen A. (1979): Innervation of the ven-

- tral prostate of the rat. American Journal of Anatomy, 154, 231–244.
- Vaalahti A., Linnoila I., Hervonen A. (1980): Immunohistochemical demonstration of VIP, [Met5]- and [Leu5]-enkephalin immunoreactive nerve fibres in the human prostate and seminal vesicles. Histochemistry, 66, 89–98.
- Vaalahti A., Tainio H., Pelto-Huikko M., Hervonen A. (1986): Light and electron microscope demonstration of VIP- and enkephalin-immunoreactive nerves in the human male genitourinary tract. Anatomical Record, 215, 21–27.
- Vega Alvarez J.A., Martinez Telleira A., Coalla C., Bengoechea M.E., Perez Casas A. (1989): Cholinergic innervation of the prostate in the white rat (*Rattus norvegicus*). Archivos Espanoles de Urologia, 42, 297–304.
- Yuri K. (1990): Immunohistochemical and enzyme histochemical localization of peptidergic, aminergic and cholinergic nerve fibres in the rat seminal vesicles. Journal of Urology, 143, 194–198.

Received: 03–12–11

Accepted after corrections: 05–01–31

Corresponding Author

Magdalena Klimczuk, Department of Animal Anatomy, Faculty of Veterinary Medicine, University of Warmia and Mazury, Oczapowskiego 13, 10-719 Olsztyn, Poland
Tel. +48 895 233 733, fax+48 895 234 986, e-mail: magdak@moskit.uwm.edu.pl
