

Genetic basis of resistance to quaternary ammonium compounds – the *qac* genes and their role: a review

Z. JAGLIC, D. CERVINKOVA

Veterinary Research Institute, Brno, Czech Republic

ABSTRACT: Although the *qac* genes are named after one of their main substrates (i.e., quaternary ammonium compounds), these genes also code for resistance to a broad spectrum of other cationic compounds such as intercalating dyes, diamidines and biguanides. The various Qac proteins are involved in relatively low specific efflux-based multidrug pumps and belong to a family of small multidrug resistance proteins. Even though the practical significance of *qac*-mediated resistance lies mainly in resistance to antiseptics, contradictory findings on this issue are still reported. Therefore, the aim of this review is to summarise the current knowledge on *qac*-mediated resistance with special emphasis on resistance to antiseptics and its relevance for practice.

Keywords: antimicrobial; disinfectant; biocide; benzalkonium; chlorhexidine; cation; susceptibility

Contents

1. Introduction
2. The *qac* genes and their distribution
3. *Qac*-mediated resistance
 - 3.1. Mechanisms of *qac*-mediated resistance
 - 3.2. Substrates of *qac*-mediated resistance
 - 3.3. Adaptive response to antiseptics
4. The *qac* genes and resistance to antibiotics
5. Conclusions
6. References

1. Introduction

Resistance to intercalating dyes (i.e. acriflavine and ethidium) was associated with particular genetic elements, namely staphylococcal β -lactamase plasmids, already more than 40 years ago (Ericson 1969; Johnston and Dyke 1969). Thereafter, various *Staphylococcus aureus* aminoglycoside-resistance plasmids were linked with resistance to quaternary ammonium compounds (QACs) and ethidium bromide (i.e. Archer and Johnston 1983; McDonnell et al. 1983). Furthermore, the commonly detected *S. aureus* plasmid pSK1 coding for resistance to aminoglycosides and trimethoprim was also found to encode resistance to intercalating dyes, QACs and diamidines (Lyon et al. 1984; Gillespie et al. 1986; Lyon and Skurray 1987). The

first described genetic determinant of resistance to antiseptics was the *qacA* gene found on pSK1 and β -lactamase/heavy metal resistance plasmids (Gillespie et al. 1986; Lyon and Skurray 1987). A range of other *qac* genes linked with particular plasmids (Littlejohn et al. 1992) were further reported as summarised below.

2. The *qac* genes and their distribution

So far, a range of various *qac* genes have been described. The *qacA* and *qacB* genes were found to share a high homology (Alam et al. 2003) and are now often designated as the *qacA/B* genes (McGann et al. 2011). A high homology was also observed between the *qacC* and *qacD* genes, i.e., *qacC* is pre-

Supported by the Ministry of Agriculture of the Czech Republic (Grants No. MZE 0002716202 and QJ1210301) and by the Ministry of Education, Youth and Sports of the Czech Republic (AdmireVet; Grant No. CZ.1.05/2.1.00/01.0006-ED 0006/01/01).

dicted to have evolved from *qacD* (Littlejohn et al. 1991). These two genes are now usually reported as *smr* (Bischoff et al. 2012). A high degree of similarity between another two *qac* genes (*qacE* and *qacF*) was shown by Ploy et al. (1998). The remaining *qac* genes include *qacG* (Heir et al. 1999), *qacH* (Heir et al. 1998), *qacJ* (Bjorland et al. 2003) and *qacZ* (Braga et al. 2011).

Although the *qac* genes are widely spread among clinical and environmental bacteria, it is obvious that their distribution is generally linked with a particular bacterial species. Among Gram-positive bacteria, the *qac* genes clearly predominate in staphylococci, in which the *qacA/B* genes were most frequently reported followed by the *qacC/D* genes (e.g. Longtin et al. 2011; Zmantar et al. 2012). Other *qac* genes, such as *qacG*, *qacH* and *qacJ*, have been less frequently observed (Ye et al. 2012). Besides staphylococci, the *qac* genes were also detected in enterococci. For instance, the *qacA/B* genes (which are highly prevalent in staphylococci) and the most recently identified *qac* gene, *qacZ*, were recently found in *Enterococcus faecalis* (Braga et al. 2011; Bischoff et al. 2012).

On the other hand, the *qacE* gene (including its attenuated variant *qacEΔ1*) is widely spread in Gram-negative bacteria, mainly in *Enterobacteriaceae* and *Pseudomonas* spp. but also in a range of other species such as *Aeromonas* spp., *Vibrio* spp., *Acinetobacter* spp. or those belonging to *Xanthomonadaceae* and *Helicobacteraceae* (Kazama et al. 1998; Chang et al. 2007; Wang et al. 2008a; Mak et al. 2009). This is mainly due to the high prevalence of class 1 integrons, which in Gram-negative bacteria commonly comprise *qacEΔ1* (Kazama et al. 1998; Bass et al. 1999; Rosser and Young 1999). Nevertheless, Kazama et al. (1998) reported that *qacEΔ1* may occur in Gram-positive cocci as well. Although less frequently, other *qac* genes were also found in Gram-negative bacteria. The *qacH* gene was most often described in *Enterobacteriaceae* (Soufi et al. 2009; Wannaprasat et al. 2011) whereas other *qac* genes (e.g. *qacF*) were reported only sporadically (Borgianni et al. 2011).

3. *Qac*-mediated resistance

3.1. Mechanisms of *qac*-mediated resistance

The involvement of an efflux-based system in *qac*-mediated resistance was first observed for the *qacA* and *qacE* genes (Jones and Midgley 1985; Tennent et al. 1989). In accordance with that, an en-

ergy-dependent efflux pump relying on the proton motive force was later confirmed as a mechanism of resistance conferred by the *qac* genes (Rouch et al. 1990; Littlejohn et al. 1992). Rouch et al. (1990) also demonstrated that *qac*-mediated efflux-based resistance has a common ancestry with tetracycline and sugar transport proteins. Further analysis of the Qac proteins revealed that these belong to a small multidrug resistance (SMR) protein family integrated in the cytoplasmic membrane via transmembrane segments and containing distinct subsets of amino acid residues involved in substrate recognition and binding (Rouch et al. 1990; Paulsen et al. 1995; Brown and Skurray 2001).

The Qac pumps are regulated via the transacting repressor protein QacR, which was shown to overlap with the promoter sequence for *qacA*. A range of diverse cationic lipophilic compounds are able to dissociate QacR from the operator DNA via its multidrug-binding pocket (Brown and Skurray 2001; Grkovic et al. 2003).

3.2. Substrates of *qac*-mediated resistance

Although the *qac* genes are named after one of their main substrates (QACs), the spectrum of their activity is much wider. *qac*-mediated resistance targets more than 30 lipophilic cationic compounds belonging to at least 12 different chemical classes (Hassan et al. 2006). Overall, substrates of the *qac* genes comprise various monovalent and divalent cationic compounds. For example, monovalent cations consist of intercalating dyes (e.g. acriflavine, ethidium, crystal violet) and the majority of QACs (e.g., benzalkonium, cetylpyridinium, cetrimide). Divalent cations include biguanidines (e.g., chlorhexidine), diamidines (e.g., propamidine, hexamidine, pentamidine), guanylhydrazones and some QACs (e.g., dequalinium; Tennent et al. 1989; Littlejohn et al. 1992; Mitchell et al. 1998; Brown and Skurray 2001). An interesting finding was reported by Bayer et al. (2006), who demonstrated that the *qacA* gene confers resistance to thrombin-induced platelet microbial protein 1, a cationic antimicrobial polypeptide released from thrombin-stimulated rabbit platelets.

The epidemiological and clinical significance of *qac*-mediated resistance is mainly attributed to resistance to antiseptics. In the first instance, the *qac* genes have been considered as important genetic determinants of resistance to QACs and, therefore,

have been monitored by many authors (e.g., Bjorland et al. 2005). However, their practical relevance in conferring resistance to QACs is highly questionable. Little or no increase in resistance to QACs in bacteria of various species carrying *qac* genes has been reported by several authors. It was shown in both Gram-positive and Gram-negative bacteria that minimum inhibitory concentrations of several QAC agents were comparable between different isolates regardless of the presence of *qac* genes (Kucken et al. 2000; Jaglic et al. 2012). In *S. aureus*, Smith et al. (2008) reported that differences in bactericidal concentrations of QACs between *qac*-positive and *qac*-negative isolates were statistically significant but lower than twofold. Furthermore, no overexpression among the *qac* genes was observed in *S. aureus* exposed to various QAC agents (DeMarco et al. 2007). In *Pseudomonas aeruginosa*, Romao et al. (2011) demonstrated that the *qacEΔ1* gene does not play an important role in resistance to benzalkonium chloride (BAC), a member of QACs. One explanation for this could lie in the fact that *qac*-mediated resistance is targeted to a broad spectrum of distinct cationic compounds resulting in the exclusion of a high-level specificity to QACs (Grkovic et al. 2003; Bay and Turner 2009). Moreover, *qac*-mediated resistance could be affected by the specific conditions, under which bacterial cells are exposed to the *qac* substrates. For example, it was reported that non-specific chemical and physical stressors may induce *qacA* transcription (Galluzzi et al. 2003).

On the other hand, there is a range of other studies reporting on a close association between the increased resistance to various cationic compounds and the presence of the *qac* genes. A high-level resistance to diamidines together with increased resistance to intercalating dyes, QACs and biguanides was described in staphylococci carrying the *qacA* and/or *qacC* genes (Leelaporn et al. 1994). A high correlation between the presence of *qacA* and resistance to BAC, hexamidine, chlorhexidine, acriflavine and ethidium bromide was observed in *S. aureus* (Behr et al. 1994). Heir et al. (1995) reported that 80% of QAC-resistant staphylococci harboured *qac* genes. Similarly, a high prevalence of *qacA/B* in isolates showing increased minimum inhibitory concentrations of acriflavine, QACs and chlorhexidine was demonstrated by Noguchi et al. (2005). In another study, the *qacA/B* gene was found in 94.6% of QAC-tolerant *S. aureus* isolates (Liu et al. 2009). Moreover, recently, the *qac* genes have been increasingly associated with resistance to

chlorhexidine (Smith et al. 2008; Wang et al. 2008b; Sheng et al. 2009). Based on the aforementioned findings, it could be generally assumed that the *qac* genes may increase resistance to various cationic antiseptics. However, their practical relevance is disputable because the commonly used concentrations (recommended by the manufacturer) of such antiseptics are usually higher than those tolerated due to action of the *qac* genes (Smith et al. 2008).

3.3. Adaptive response to antiseptics

It should be further mentioned that the *qac* genes may play an important role in adaption to various cationic compounds. A stepwise adaption to twofold increasing concentrations of QACs was recently described in *S. aureus* carrying the *qac* genes (Smith et al. 2008). Sundheim et al. (1998) formerly observed that although resistance to QACs is generally low among *qac*-positive staphylococci, resistant strains were isolated after exposure to a QAC disinfectant. Furthermore, Heir et al. (1999) showed that under the pressure of sublethal concentrations of BAC, *qacG*-harbouring staphylococci adapted to this antiseptic. Similarly, increased post-exposure resistance to chlorhexidine was observed in *S. aureus* carrying the *qac* genes (Vali et al. 2008).

However, adaption to BAC, chlorhexidine and some other antiseptics was also described in staphylococci negative for the *qac* genes (Heir et al. 1999; Vali et al. 2008). Therefore, the question as to whether the adaption to cationic compounds is driven by the *qac* genes or rather by other mechanisms remains open. Modifications in the cell wall resulting in increased biocide tolerance have already been described by McDonnell and Russell (1999). Moreover, an increased cell-wall thickness (rather than the action of the *qac* genes) has been recently demonstrated to be responsible for the post-exposure resistance to acriflavine (Kawai et al. 2009).

4. The *qac* genes and resistance to antibiotics

A certain degree of association over time between bacterial resistance to antiseptics and antibiotics has been reported. It has been observed that some bacteria which express increased resistance to antiseptics are generally less susceptible to antibiotics. Outer membrane changes have been believed to be one of the mechanisms responsible for such

increased non-specific cross-resistance (Russell 2000). Whether the *qac* genes are able to directly confer resistance to antibiotics remains to be determined. Fuentes et al. (2005) recently demonstrated that *qacC* confers resistance to a number of β -lactam antibiotics. In addition, the *nor* genes, coding for resistance to fluoroquinolones, were also found to encode resistance to antiseptics and further, to have similar substrates to those of the *qac* genes (DeMarco et al. 2007; Theis et al. 2007). This indicates that resistance to antibiotics and antiseptics may closely interface (Langsrud et al. 2004).

Nevertheless, a close association between resistance to antiseptics and antibiotics could be primarily explained by the fact that genetic determinants of resistance to these agents are commonly linked with each other, i.e., the *qac* genes are typically present on plasmids together with a range of other resistance genes. For instance, the linkage between resistance to trimethoprim (*dfrA*), β -lactams (*blaZ*), aminoglycosides (*aacA-aphD*) and antiseptics (*qacC*) mediated by a multi-resistance plasmid was reported in staphylococci (Weigel et al. 2003). In *S. aureus*, it is known that the *qacA/B* genes frequently occur on the pSK1 and β -lactamase/heavy metal-resistance plasmids which also confer resistance to a range of antibiotics (Mayer et al. 2001). In Gram-negative bacteria, the *qac* genes are often linked with plasmid-mediated class 1 integrons which harbour a variety of antibiotic resistance genes (Zhao et al. 2012). In *Enterobacteriaceae*, the *qac* genes have been most frequently found in combination with genes coding for resistance to aminoglycosides, chloramphenicol, sulphonamides, trimethoprim and β -lactams (Poirel et al. 2000; Riano et al. 2006; Espedido et al. 2008; Zhao et al. 2012). Similar findings were also reported in *Pseudomonas aeruginosa* (Jeong et al. 2009; Colinon et al. 2010). The link between the *qac* genes and macrolide inactivation genes was revealed in *Aeromonas hydrophila* (Poole et al. 2006) as well as in microflora from a wastewater treatment plant (Szczepanowski et al. 2004). Therefore, the use of various cationic biocides may be also responsible for the selection of bacteria resistant to antibiotics (Russell 2000; Hegstad et al. 2010).

5. CONCLUSIONS

The *qac* genes may increase resistance to various cationic compounds and, therefore, they are of

epidemiological and clinical significance. However, their specificity to particular substrates is relatively low and, compared to some antibiotic resistance genes they confer a relatively low level of resistance that can be positively or negatively affected by specific environmental conditions. It seems that the mere presence of the *qac* genes does not necessarily imply increased resistance to antiseptics that could be relevant for practice. Nevertheless, these genes together with other mechanisms (such as various multidrug efflux genes or modifications in the cell wall) probably contribute to the development of resistance to cationic antiseptics and survival of bacteria in toxic environments.

6. REFERENCES

- Alam MM, Kobayashi N, Uehara N, Watanabe N (2003): Analysis on distribution and genomic diversity of high-level antiseptic resistance genes *qacA* and *qacB* in human clinical isolates of *Staphylococcus aureus*. *Microbial Drug Resistance-Mechanisms Epidemiology and Disease* 9, 109–121.
- Archer GL, Johnston JL (1983): Self-transmissible plasmids in staphylococci that encode resistance to aminoglycosides. *Antimicrobial Agents and Chemotherapy* 24, 70–77.
- Bass L, Liebert CA, Lee MD, Summers AO, White DG, Thayer SG, Maurer JJ (1999): Incidence and characterization of integrons, genetic elements mediating multiple-drug resistance, in avian *Escherichia coli*. *Antimicrobial Agents and Chemotherapy* 43, 2925–2929.
- Bay DC, Turner RJ (2009): Diversity and evolution of the small multidrug resistance protein family. *BMC Evolutionary Biology* 9, 140.
- Bayer AS, Kupferwasser LI, Brown MH, Skurray RA, Grkovic S, Jones T, Mukhopadhyay K, Yeaman MR (2006): Low-level resistance of *Staphylococcus aureus* to thrombin-induced platelet microbicidal protein 1 in vitro associated with *qacA* gene carriage is independent of multidrug efflux pump activity. *Antimicrobial Agents and Chemotherapy* 50, 2448–2454.
- Behr H, Reverdy ME, Mabilat C, Freney J, Fleurette J (1994): Relationship between the MIC levels of 5 antiseptics and presence of the *qacA* gene in *Staphylococcus aureus* (in French). *Pathologie et Biologie* 42, 438–444.
- Bischoff M, Bauer J, Preikschat P, Schwaiger K, Molle G, Holz C (2012): First detection of the antiseptic resistance gene *qacA/B* in *Enterococcus faecalis*. *Microbial Drug Resistance* 18, 7–12.

- Bjorland J, Steinum T, Sunde M, Waage S, Heir E (2003): Novel plasmid-borne gene *qacJ* mediates resistance to quaternary ammonium compounds in equine *Staphylococcus aureus*, *Staphylococcus simulans*, and *Staphylococcus intermedius*. *Antimicrobial Agents and Chemotherapy* 47, 3046–3052.
- Bjorland J, Steinum T, Kvitle B, Waage S, Sunde M, Heir E (2005): Widespread distribution of disinfectant resistance genes among staphylococci of bovine and caprine origin in Norway. *Journal of Clinical Microbiology* 43, 4363–4368.
- Borgianni L, Prandi S, Salden L, Santella G, Hanson ND, Rossolini GM, Docquier JD (2011): Genetic context and biochemical characterization of the IMP-18 metallo-beta-lactamase identified in a *Pseudomonas aeruginosa* isolate from the United States. *Antimicrobial Agents and Chemotherapy* 55, 140–145.
- Braga TM, Marujo PE, Pomba C, Lopes MFS (2011): Involvement, and dissemination, of the enterococcal small multidrug resistance transporter QacZ in resistance to quaternary ammonium compounds. *Journal of Antimicrobial Chemotherapy* 66, 283–286.
- Brown MH, Skurray RA (2001): Staphylococcal multidrug efflux protein QacA. *Journal of Molecular Microbiology and Biotechnology* 3, 163–170.
- Chang YC, Shih DY, Wang JY, Yang SS (2007): Molecular characterization of class 1 integrons and antimicrobial resistance in *Aeromonas* strains from foodborne outbreak-suspect samples and environmental sources in Taiwan. *Diagnostic Microbiology and Infectious Disease* 59, 191–197.
- Colinon C, Jocktane D, Brothier E, Rossolini GM, Cournoyer B, Nazaret S (2010): Genetic analyses of *Pseudomonas aeruginosa* isolated from healthy captive snakes: evidence of high inter- and intrasite dissemination and occurrence of antibiotic resistance genes. *Environmental Microbiology* 12, 716–729.
- DeMarco CE, Cushing LA, Frempong-Manso E, Seo SM, Jaravaza TAA, Kaatz GW (2007): Efflux-related resistance to norfloxacin, dyes, and biocides in bloodstream isolates of *Staphylococcus aureus*. *Antimicrobial Agents and Chemotherapy* 51, 3235–3239.
- Ericson C (1969): Resistance to acriflavine and cadmium and changed phage reactions – markers of a new staphylococcal penicillinase plasmid. *Acta Pathologica et Microbiologica Scandinavica* 76, 333.
- Espedido BA, Partridge SR, Iredell JR (2008): bla(IMP-4) in different genetic contexts in *Enterobacteriaceae* isolates from Australia. *Antimicrobial Agents and Chemotherapy* 52, 2984–2987.
- Fuentes DE, Navarro CA, Tantalean JC, Araya MA, Saavedra CP, Perez JM, Calderon IL, Youderian PA, Mora GC, Vasquez CC (2005): The product of the *qacC* gene of *Staphylococcus epidermidis* CH mediates resistance to beta-lactam antibiotics in Gram-positive and Gram-negative bacteria. *Research in Microbiology* 156, 472–477.
- Galluzzi L, Virtanen P, Karp M (2003): A real-time analysis of QacR-regulated multidrug resistance in *Staphylococcus aureus*. *Biochemical and Biophysical Research Communications* 301, 24–30.
- Gillespie MT, May JW, Skurray RA (1986): Plasmid-encoded resistance to acriflavine and quaternary ammonium compounds in methicillin-resistant *Staphylococcus aureus*. *Fems Microbiology Letters* 34, 47–51.
- Grkovic S, Hardie KM, Brown MH, Skurray RA (2003): Interactions of the QacR multidrug-binding protein with structurally diverse ligands: implications for the evolution of the binding pocket. *Biochemistry* 42, 15226–15236.
- Hassan KA, Galea M, Wu JQ, Mitchell BA, Skurray RA, Brown MH (2006): Functional effects of intramembranous proline substitutions in the staphylococcal multidrug transporter QacA. *Fems Microbiology Letters* 263, 76–85.
- Hegstad K, Langsrud S, Lunestad BT, Scheie AA, Sunde M, Yazdankhah SP (2010): Does the wide use of quaternary ammonium compounds enhance the selection and spread of antimicrobial resistance and thus threaten our health? *Microbial Drug Resistance* 16, 91–104.
- Heir E, Sundheim G, Holck AL (1995): Resistance to quaternary ammonium compounds in *Staphylococcus* spp. isolated from the food industry and nucleotide sequence of the resistance plasmid pST827. *Journal of Applied Bacteriology*, 79, 149–156.
- Heir E, Sundheim G, Holck AL (1998): The *Staphylococcus qacH* gene product: a new member of the SMR family encoding multidrug resistance. *FEMS Microbiology Letters* 163, 49–56.
- Heir E, Sundheim G, Holck AL (1999): The *qacG* gene on plasmid pST94 confers resistance to quaternary ammonium compounds in staphylococci isolated from the food industry. *Journal of Applied Microbiology* 86, 378–388.
- Jaglic Z, Cervinkova D, Vlkova H, Michu E, Kunova G, Babak V (2012): Bacterial biofilms resist oxidising agents due to the presence of organic matter. *Czech Journal of Food Sciences* 30, 178–187.
- Jeong JH, Shin KS, Lee JW, Park EJ, Son SY (2009): Analysis of a novel class 1 integron containing metallo-beta-lactamase gene VIM-2 in *Pseudomonas aeruginosa*. *Journal of Microbiology* 47, 753–759.
- Johnston LH, Dyke KGH (1969): Ethidium bromide resistance, a new marker on the staphylococcal penicillinase plasmid. *Journal of Bacteriology* 100, 1413–1414.

- Jones IG, Midgley M (1985): Expression of a plasmid borne ethidium resistance determinant from *Staphylococcus* in *Escherichia coli* – evidence for an efflux system. FEMS Microbiology Letters 28, 355–358.
- Kawai M, Yamada S, Ishidoshio A, Oyamada Y, Ito H, Yamagishi J (2009): Cell-wall thickness: possible mechanism of acriflavine resistance in methicillin-resistant *Staphylococcus aureus*. Journal of Medical Microbiology 58, 331–336.
- Kazama H, Hamashima H, Sasatsu M, Arai T (1998): Distribution of the antiseptic-resistance gene *qacE* delta 1 in Gram-positive bacteria. FEMS Microbiology Letters 165, 295–299.
- Kucken D, Feucht HH, Kaulfers PM (2000): Association of *qacE* and *qacE* delta 1 with multiple resistance to antibiotics and antiseptics in clinical isolates of Gram-negative bacteria. FEMS Microbiology Letters 183, 95–98.
- Langsrud S, Sundheim G, Holck AL (2004): Cross-resistance to antibiotics of *Escherichia coli* adapted to benzalkonium chloride or exposed to stress-inducers. Journal of Applied Microbiology 96, 201–208.
- Leelaporn A, Paulsen IT, Tennent JM, Littlejohn TG, Skurray RA (1994): Multidrug resistance to antiseptics and disinfectants in coagulase-negative staphylococci. Journal of Medical Microbiology 40, 214–220.
- Littlejohn TG, DiBerardino D, Messerotti LJ, Spiers SJ, Skurray RA (1991): Structure and evolution of a family of genes encoding antiseptic and disinfectant resistance in *Staphylococcus aureus*. Gene 101, 59–66.
- Littlejohn TG, Paulsen IT, Gillespie MT, Tennent JM, Midgley M, Jones IG, Purewal AS, Skurray RA (1992): Substrate specificity and energetics of antiseptic and disinfectant resistance in *Staphylococcus aureus*. FEMS Microbiology Letters 95, 259–266.
- Liu QZ, Liu MN, Wu Q, Li C, Zhou TL, Ni YX (2009): Sensitivities to biocides and distribution of biocide resistance genes in quaternary ammonium compound tolerant *Staphylococcus aureus* isolated in a teaching hospital. Scandinavian Journal of Infectious Diseases 41, 403–409.
- Longtin J, Seah C, Siebert K, McGeer A, Simor A, Longtin Y, Low DE, Melano RG (2011): Distribution of antiseptic resistance genes *qacA*, *qacB*, and *smr* in methicillin-resistant *Staphylococcus aureus* isolated in Toronto, Canada, from 2005 to 2009. Antimicrobial Agents and Chemotherapy 55, 2999–3001.
- Lyon BR, Skurray R (1987): Antimicrobial resistance of *Staphylococcus aureus* – genetic-basis. Microbiological Reviews 51, 88–134.
- Lyon BR, May JW, Skurray RA (1984): Tn4001 – a gentamicin and kanamycin resistance transposon in *Staphylococcus aureus*. Molecular and General Genetics 193, 554–556.
- Mak JK, Kim MJ, Pham J, Tapsall J, White PA (2009): Antibiotic resistance determinants in nosocomial strains of multidrug-resistant *Acinetobacter baumannii*. Journal of Antimicrobial Chemotherapy 63, 47–54.
- Mayer S, Boos M, Beyer A, Fluit AC, Schmitz FJ (2001): Distribution of the antiseptic resistance genes *qacA*, *qacB* and *qacC* in 497 methicillin-resistant and -susceptible European isolates of *Staphylococcus aureus*. Journal of Antimicrobial Chemotherapy 47, 896–897.
- McDonnell G, Russell AD (1999): Antiseptics and disinfectants: Activity, action, and resistance. Clinical Microbiology Reviews 12, 147–179.
- McDonnell RW, Sweeney HM, Cohen S (1983): Conjugational transfer of gentamicin resistance plasmids intraspecifically and interspecifically in *Staphylococcus aureus* and *Staphylococcus epidermidis*. Antimicrobial Agents and Chemotherapy 23, 151–160.
- McGann P, Kwak YI, Summers A, Cummings JF, Waterman PE, Lesho EP (2011): Detection of *qacA/B* in clinical isolates of methicillin-resistant *Staphylococcus aureus* from a regional healthcare network in the eastern United States. Infection Control and Hospital Epidemiology 32, 1116–1119.
- Mitchell BA, Brown MH, Skurray RA (1998): QacA multidrug efflux pump from *Staphylococcus aureus*: comparative analysis of resistance to diamidines, biguanidines, and guanilylhydrazones. Antimicrobial Agents and Chemotherapy 42, 475–477.
- Noguchi N, Suwa J, Narui K, Sasatsu M, Ito T, Hiramatsu K, Song JH (2005): Susceptibilities to antiseptic agents and distribution of antiseptic-resistance genes *qacA/B* and *smr* of methicillin-resistant *Staphylococcus aureus* isolated in Asia during 1998 and 1999. Journal of Medical Microbiology 54, 557–565.
- Paulsen IT, Brown MH, Dunstan SJ, Skurray RA (1995): Molecular characterization of the staphylococcal multidrug-resistance export protein QacC. Journal of Bacteriology 177, 2827–2833.
- Ploy MC, Courvalin P, Lambert T (1998): Characterization of In40 of *Enterobacter aerogenes* BM2688, a class 1 integron with two new gene cassettes, *cmlA2* and *qacF*. Antimicrobial Agents and Chemotherapy 42, 2557–2563.
- Poirel L, Le Thomas I, Naas T, Karim A, Nordmann P (2000): Biochemical sequence analyses of GES-1, a novel class A extended-spectrum beta-lactamase, and the class 1 integron In52 from *Klebsiella pneumoniae*. Antimicrobial Agents and Chemotherapy 44, 622–632.
- Poole TL, Callaway TR, Bischoff KM, Warnes CE, Nisbet DJ (2006): Macrolide inactivation gene cluster mphA-mrx-mphR adjacent to a class 1 integron in *Aeromonas hydrophila* isolated from a diarrhoeic pig in Oklahoma. Journal of Antimicrobial Chemotherapy 57, 31–38.

- Riano I, Moreno MA, Teshager T, Saenz Y, Dominguez L, Torres C (2006): Detection and characterization of extended-spectrum beta-lactamases in *Salmonella enterica* strains of healthy food animals in Spain. *Journal of Antimicrobial Chemotherapy* 58, 844–847.
- Romao C, Miranda CA, Silva J, Clementino MM, de Filippis I, Asensi M (2011): Presence of *qacE* delta 1 gene and susceptibility to a hospital biocide in clinical isolates of *Pseudomonas aeruginosa* resistant to antibiotics. *Current Microbiology* 63, 16–21.
- Rosser SJ, Young HK (1999): Identification and characterization of class 1 integrons in bacteria from an aquatic environment. *Journal of Antimicrobial Chemotherapy* 44, 11–18.
- Rouch DA, Cram DS, Diberardino D, Littlejohn TG, Skurray RA (1990): Efflux-mediated antiseptic resistance gene *qacA* from *Staphylococcus aureus* – common ancestry with tetracycline-transport and sugar-transport proteins. *Molecular Microbiology* 4, 2051–2062.
- Russell AD (2000): Do biocides select for antibiotic resistance? *Journal of Pharmacy and Pharmacology* 52, 227–233.
- Sheng WH, Wang JT, Lauderdale TL, Weng CM, Chen DL, Chang SC (2009): Epidemiology and susceptibilities of methicillin-resistant *Staphylococcus aureus* in Taiwan: emphasis on chlorhexidine susceptibility. *Diagnostic Microbiology and Infectious Disease* 63, 309–313.
- Smith K, Gemmell CG, Hunter IS (2008): The association between biocide tolerance and the presence or absence of *qac* genes among hospital-acquired and community-acquired MRSA isolates. *Journal of Antimicrobial Chemotherapy* 61, 78–84.
- Soufi L, Abbassi MS, Saenz Y, Vinue L, Somalo S, Zarazaga M, Abbas A, Dbaya R, Khanfir L, Ben Hassen A, Hammami S, Torres C (2009): Prevalence and diversity of integrons and associated resistance genes in *Escherichia coli* isolates from poultry meat in Tunisia. *Foodborne Pathogens and Disease* 6, 1067–1073.
- Sundheim G, Langsrud S, Heir E, Holck AL (1998): Bacterial resistance to disinfectants containing quaternary ammonium compounds. *International Biodeterioration and Biodegradation* 41, 235–239.
- Szczepanowski R, Krahn I, Linke B, Goesmann A, Puhler A, Schluter A (2004): Antibiotic multiresistance plasmid pRSB101 isolated from a wastewater treatment plant is related to plasmids residing in phytopathogenic bacteria and carries eight different resistance determinants including a multidrug transport system. *Microbiology-SGM* 150, 3613–3630.
- Tennent JM, Lyon BR, Midgley M, Jones IG, Purewal AS, Skurray RA (1989): Physical and biochemical characterization of the *qacA* gene encoding antiseptic and disinfectant resistance in *Staphylococcus aureus*. *Journal of General Microbiology* 135, 1–10.
- Theis T, Skurray RA, Brown MH (2007): Identification of suitable internal controls to study expression of a *Staphylococcus aureus* multidrug resistance system by quantitative real-time PCR. *Journal of Microbiological Methods* 70, 355–362.
- Vali L, Davies SE, Lai LLG, Dave J, Amyes SGB (2008): Frequency of biocide resistance genes, antibiotic resistance and the effect of chlorhexidine exposure on clinical methicillin-resistant *Staphylococcus aureus* isolates. *Journal of Antimicrobial Chemotherapy* 61, 524–532.
- Wang C, Zhan Q, Mi Z, Huang Z, Chen G (2008a): Distribution of the antiseptic-resistance gene *qacE* delta 1 in 283 clinical isolates of Gram-negative bacteria in China. *Journal of Hospital Infection* 69, 394–396.
- Wang JT, Sheng WH, Wang JL, Chen D, Chen ML, Chen YC, Chang SC (2008b): Longitudinal analysis of chlorhexidine susceptibilities of nosocomial methicillin-resistant *Staphylococcus aureus* isolates at a teaching hospital in Taiwan. *Journal of Antimicrobial Chemotherapy* 62, 514–517.
- Wannaprasat W, Padungtod P, Chuanchuen R (2011): Class 1 integrons and virulence genes in *Salmonella enterica* isolates from pork and humans. *International Journal of Antimicrobial Agents* 37, 457–461.
- Weigel LM, Clewell DB, Gill SR, Clark NC, McDougal LK, Flannagan SE, Kolonay JE, Shetty J, Killgore GE, Tenover FC (2003): Genetic analysis of a high-level vancomycin-resistant isolate of *Staphylococcus aureus*. *Science* 302, 1569–1571.
- Ye HF, Zhang M, O'Donoghue M, Boost M (2012): Are *qacG*, *qacH* and *qacJ* genes transferring from food isolates to carriage isolates of staphylococci? *Journal of Hospital Infection* 80, 95–96.
- Zhao WH, Chen GL, Ito R, Kimura S, Hu ZQ (2012): Identification of a plasmid-borne bla(IMP-11) gene in clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae*. *Journal of Medical Microbiology* 61, 246–251.
- Zmantar T, Kouidhi B, Hentati H, Bakhrouf A (2012): Detection of disinfectant and antibiotic resistance genes in *Staphylococcus aureus* isolated from the oral cavity of Tunisian children. *Annals of Microbiology* 62, 123–128.

Received: 2012–05–28

Accepted after corrections: 2012–06–06

Corresponding Author:

Zoran Jaglic, Veterinary Research Institute, Hudcova 70, 621 00 Brno, Czech Republic
Tel. +420 533 331 619, E-mail: jaglic@vri.cz