

Impact of atmospheric ammonia on growth, C and N accumulation and photosynthesis of two maize cultivars with different N root supply

L.X. Zhang^{1,2}, H. Qiang^{1,2,3}, S.Q. Li^{1,2,3}, X.L. Chen^{1,2,3}

¹State Key Laboratory of Soil Erosion and Dryland Farming and College of Life Sciences, Northwest A & F University, Yangling, Shaanxi, P.R. China

²State Key Laboratory of Soil Erosion and Dryland Farming, Institute of Soil and Water Conservation, Chinese Academy of Sciences and Northwest A & F University, Yangling, Shaanxi, P.R. China

³College of Resource and Environment, Northwest A & F University, Yangling, Shaanxi, P.R. China

ABSTRACT

Impact of enriched atmospheric NH₃ in combination with low and high N medium on growth, total C and N accumulation (C_{tot}A and N_{tot}A) and photosynthetic characteristics of two maize cultivars i.e. SD19 (cult. 1) and NE5 (cult. 2) with low N and N high use efficiency, respectively, was investigated. Plants were exposed to 10 nl/L and 1000 nl/L NH₃ fumigation, respectively, for 30 days in open-top chambers (OTCs). Under exposure to the low N medium, increase of the atmospheric NH₃ concentration to 1000 nl/L from the ambient level significantly ($P < 0.05$) increased dry matter (DM) (by 18% in cult. 1 and 14% in cult. 2 respectively), C_{tot}A, N_{tot}A, net photosynthetic rate (P_n), stomatal conductance (G_s) and apparent quantum yield (AQY) but decreased intercellular CO₂ concentration (C_i) in both cultivars. These effects were more pronounced in cult. 1 as compared to those in cult. 2. In contrast, in the high N solution, enriched atmospheric NH₃ led to a decrease in DM, C_{tot}A, N_{tot}A, P_n, G_s and AQY but an increase in C_i of cult. 2 only. Dark respiration rate remained unaffected by enrichment of NH₃ in each treatment. Therefore, it is concluded that appropriately enriched atmospheric NH₃ can improve plant growth of maize by enhancing C_{tot}A, N_{tot}A, and photosynthesis in the low N medium, especially for low N use efficiency cultivars.

Keywords: atmospheric ammonia enrichment; dry matter; carbon accumulation; nitrogen accumulation; photosynthetic characteristic

Among the atmospheric N species, ammonia (NH₃) is not only an important atmospheric pollutant for semi-natural vegetation, but one of the key N sources for N-deficiency plants in atmosphere. Annual NH₃-N deposition in each of the 4 years 2003–2006 was estimated to increase from 3.0 ± 0.2 kg N/ha year in ambient air, with the NH₃ concentration at 0.5 m above the canopy of 0.7 µg/m³, to 50–70 kg N/ha year where annual

average air concentrations were 70–90 µg/m³ and concentrations during fumigation were up to 1600 µg/m³ (Cape et al. 2008). Factors contributing to this significant increase in atmosphere NH₃ concentration include: vicinity to location of significant NH₃ emissions, climatic conditions, and agriculture activities. The latter contributes to 50% of the global NH₃ emissions (Fangmeier et al. 1994). Maize (*Zea mays* L.) is a staple food

Supported by the National Natural Science Foundation of China, Project No. 30571116, by the National Basic Research Program of China, Project No. 2009CB118604; by the Governmental Public Industry Research Special Funds, Project No. 200803030; by the China Postdoctoral Science Foundation and Chinese Universities Scientific Fund, Project No. QN2009069, and by the Foundation of State Key Laboratory of Soil Erosion and Dryland Farming, Project No. 10501-J-3.

crop in northern China. This region is usually subjected to increasing atmospheric NH_3 emission to 0.30–1.60 mg/m^3 from regular mean air NH_3 concentration of 0.01–0.20 mg/m^3 due to high rate application of NH_3 forming fertilizers (Li et al. 2004). Atmospheric NH_3 is deposited to soil and water, either by dry deposition of NH_3 or by dry and wet deposition of ammonium (NH_4^+) (Fangmeier et al. 1994). As a plant nutrient, plants can utilize atmospheric NH_3 to improve plant growth and/or production if the mole fraction of NH_3 in the atmosphere is greater than the mole fraction of gaseous NH_3 in the substomatal cavity. Foliar NH_3 uptake by the leaves of Italian ryegrass (*Lolium multiflorum*) at 0.52 mg/m^3 supplied 47.3% and 35.2% of total plant N at fertilization levels of 100 and 200 $\text{mg } ^{15}\text{NO}_3^- \text{-N}/\text{kg}$ dry soil, respectively (Fangmeier et al. 1994). It was found that atmospheric NH_3 input is closely correlated with the morphology and metabolism of crops (Bohme et al. 2003, Li et al. 2009).

Exposure of plants to enriched atmospheric NH_3 may result in a significant impact on plant growth, total C and N (C_{tot} and N_{tot}) accumulation as well as photosynthesis, whose responses were dependent on plant species, growth stages and N availability for root uptake (Van der Eerden et al. 1991, 1992, Sommer et al. 1993, Tatsuro et al. 2001, Li et al. 2004, 2009). Li et al. (2009) concluded that enriched atmospheric NH_3 concentration increased shoot dry matter of wheat (*Triticum aestivum* L.) in the low N treatment but reduced the plant biomass in the high N treatment. In most cases, exposure to NH_3 generally results in an increase of soluble proteins concentrations in spruce (*Picea abies*) trees, N_{tot} and chlorophyll concentrations of Scots pine needles as well as photosynthesis of *Pinus sylvestris*, poplar trees and Douglas fir and dark respiration rate (R_d) of *Populus euramericana* as long as the tissue is not injured (Van Hove et al. 1989, Van der Eerden et al. 1992, Van der Eerden and Pérez-Soba 1992). Castro et al. (2005) pointed out that atmospheric NH_3 up to 4 $\mu\text{l}/\text{L}$ can be regarded as a nutrient for the fast growing of *B. oleracea*. The concentration at which NH_3 changes from being a nutrient to a toxin is not clear-cut, since NH_3 can still be metabolized when plant growth is already affected (Fangmeier et al. 1994).

However, above studies mainly focus on forest and grassland vegetation. Much less attention has been paid to the effects of atmospheric NH_3 in agro-ecosystems. Additionally, evaluation of responses of maize crop to varying concentration


of atmospheric NH_3 are not fully investigated. Keeping in view the above facts, we hypothesize that higher constitutive accumulation of C_{tot} and N_{tot} and photosynthesis induced by enriched atmospheric NH_3 may provide a mechanism of the improvement of plant growth of different maize cultivars under two levels of N in solution medium. Therefore, the objective of this study was to examine the growth response and accumulation of C_{tot} and N_{tot} as well as photosynthetic characteristic parameters of two maize cultivars, with different N use efficiency of plant, exposed to two levels of NH_3 concentration and N root supply in solution medium for 30 days.

MATERIALS AND METHODS

Plant material and experimental design.

Solution culture experiments were conducted at Institute of Soil and Water Conservation, Chinese Academy of Sciences (Yangling, China). Two maize (*Zea mays* L.) cultivars i.e. SD19 (cult. 1) and NE5 (cult. 2) with low N and high N use efficiency, respectively, were supplied for the present experiments.

The open-top chambers (OTCs) used in the experiments (Figure 1) are containers described by Paul and Bert (1993), initially designed for exposing plants to elevated CO_2 concentrations under close to natural conditions, with a square 1.2 × 1.2 m base and 1.5 m tall perpendicular glass walls topped by glass quadrilaterals inclined towards the centre in an iron frame (total volume circa 3 m^3). The chambers were each equipped with a fan and an air control system, which included a steel cylinder (inner diameter 600 mm, total length 1800 mm) containing 95% NH_3 . The NH_3 was fed from the bottom into the OTCs through a YQA-441 NH_3 decrement gauge with a pressure range of 0–4 MPa (Shanghai Shuangying Boat Decompressor Manufacture Co., Ltd., Shanghai, China) and $\Phi 8$ constant pressure oxygen pipes. The NH_3 flux was measured using an LZB-2 flux meter with anti-corrosion glass rotameter (measuring range 6–60 ml/min , rated working pressure ≤ 1 MPa) (Changzhou Shuangfa Thermal Instrument Co., Ltd., Changzhou, China). In addition, air was fed from the bottom of the OTCs using a ZB-0.10/8 air compressor (air displacement 0.1 m^3/min , rated air pressure 0.8 MPa) and $\Phi 8$ constant pressure oxygen pipes (Shanghai Luodi Air Compressor Co., Ltd., Shanghai, China). The air flux was maintained at 1.7 L/min , as measured by an LZB-2 flux meter



with an anti-corrosion glass rotameter (measuring range 0.25–2.5 m³/h, rated working pressure ≤ 1 MPa) (Changzhou Shuangfa Thermal Instrument Co., Ltd., Changzhou, China).

The temperature, humidity and NH₃ concentration inside each chamber were regulated by passing air (heated and moistened as appropriate) and NH₃ through porous pipes at the bottom of the container, while the fan (providing air speed of less than 0.5 m/s) was used to maintain close to uniform distribution of NH₃ and reduced temperature (if necessary) in the chamber. The temperature in the chambers was monitored to verify that the temperature was the same in the chamber before and after NH₃ was supplied, and thus that the effects of varying the NH₃ concentration on the growth parameters or C and N metabolism of plants were not confounded by variations in temperature.

The NH₃ concentration in each chamber was measured four times per day (at 8:00, 11:00, 14:00 and 17:00 h) by a GTL-C indoor air detector equipped with a pH618 test pen (NH₃ testing precision, ± 0.01 mg/m³) (Shanghai Minyi Electron Co., Ltd., Shanghai, China) mounted on a tripod placed in the centre of the chamber before the gas was supplied. On each sampling occasion, 5 ml of NH₃ test reagent was extracted by an injector and injected into a glass bottle for sampling. The glass bottle was immediately plugged and connected to the instrument. During the tests, the flux was adjusted to 2 L/min and the exhaust time was controlled by the auto-timing device. When the sampling time was complete, the glass bottle was removed, unplugged, the reagent in the glass bottle was poured into the test cup and the test pen was inserted into the cup to measure the

NH₃ concentration. Throughout the entire growth period, from 8:00–18:00 h every day, the NH₃ concentration in the OTCs used for the background, ambient (control) and elevated NH₃ treatments were maintained precisely at 10 nl/L and 1000 nl/L, respectively, by continuously supplying NH₃ and air at appropriate ratios.

Plant growth. Seeds of two maize cultivars were surface-sterilized in 10% H₂O₂ solution for 15 min. After rinsing in distilled water, seeds were imbibed for 12 h and then sown in porcelain trays containing quartz sand. Seeds were germinated in the dark under 23°C covering with clean wet filter papers. When the root grew to the length of 2–4 cm, the seedling was transferred and fixed in the holes of styrofoam boards by using absorbent cotton in deionized water in plastic trays in the growth chamber under the conditions with 25/18°C of average day/night temperature, 60–70% relative humidity, and 350 μmol/m²/s light intensity and 16/8 h of light/dark regime. The solution was replaced once a day and aerated continuously.

PVC (polyvinyl chloride) pots with a volume of 3534 cm³ (inner diameter 15 cm, height 20 cm) containing 3.5 L nutrient solution were used for plant growth. Ammonia fumigation and N treatments were initially proceeded on the 3rd day after the seedling transferred into OTCs at their three-leaf stage. The experimental design contained three factors which were NH₃ atmospheric concentration (at two levels: 10 nl/L, air background concentration and 1000 nl/L, high NH₃ concentration), N rate in medium (at two levels: 1/3 and 1/9 N concentration of complete nutrition solution i.e. 5.00 and 1.67 mmol/L nitrate) and maize cultivar (the two cultivars mentioned above), respectively, in a complete factorial design experiment with eight

treatments, and five replications per treatment. The 40 pots (one plant per pot) were randomly placed in four OTCs, and in order to reduce the experimental error the pots were exchanged in the four OTCs every 7 days. The low and high N rate in medium were served by 1/9 and 1/3 strength of the complete Hoagland nutrient solution, respectively. Desired N concentrations were maintained by irrigating sufficiently with new solution. The nutrient solution was amended with higher alcohol to make emulsion in order to restrain NH₃ exchange between two phase of gas and liquid. The pH of solution was adjusted to 6.2 (± 0.1). The top of pot was placed on a board with small holes for plants fixation, and the space between plants and holes were sealed with wax. The solution was aerated without NH₃ twelve hours a day. The plants of two maize cultivars were maintained for 30-day growth period under each treatment. The whole experiment was carried out twice independently under the same environmental conditions. Data presented here are means of five replicates of the two experiments (*n* = 10).

Measurements of dry matter and concentrations of total C and total N. Shoots of five plants in five pots of representing each cultivar and each of the treatments were collected from each chamber on the 30th day of N and NH₃ fumigation treatments (from 11:00 to 13:00 h), respectively. The samples were cleaned with distilled water, and were dried in a forced-ventilation oven at 65°C until constant dry weight. The dried samples were ground to pass a 1 mm screen for total C (C_{tot}) and total N (N_{tot}) assay. Concentrations of C_{tot} and N_{tot} were measured by potassium dichromate (K₂Cr₂O₇) volume method and the Kjeldahl method using an automatic N analyzer (Gerhardt Vapodest 5), respectively. Their total C and N accumulation (C_{tot}A and N_{tot}A) based on the aboveground dry matter were calculated by multiplying their C and N content and dry matter, respectively.

Determination of photosynthetic parameters. The second completely developed leaf from the top of sample plant on the 30th day of NH₃ fumigation were measured from 9:00 to 11:00 h, respectively, using LI-6400 portable photosynthesis system (LI-COR Inc., Lincoln, NE, USA). The two leaves of one plant were determined and this was repeated in three replicates. The determination covered net photosynthetic rate (P_n, μmol CO₂/m²/s), intercellular CO₂ concentration (C_i, μmol/mol), stomatal conductance (G_s, mol/m²/s) under the conditions of photosynthetically active radiation (PAR) of 1200 μmol/m²/s and ambient CO₂ concentra-

tion of 360 μmol/mol. Apparent quantum yield (AQY, mol CO₂/mol) and dark respiration rate (R_d, μmol CO₂/m²/s) were computed referring to the index model (1) of light response of maize under the setting PAR values of 2000, 1600, 1400, 1200, 1000, 800, 600, 400, 200, 150, 100, 50 and 0 μmol/m²/s in turn and ambient CO₂ concentration of 360 μmol/mol condition (Guo et al. 2005).

$$P_n = P_{\max} \left(1 - e^{-\frac{AQY \cdot PAR}{P_{\max}}} \right) - |R_d| \quad (1)$$

Where: P_n represents leaf net photosynthetic rate under different PAR, and P_{max} indicates maximum net photosynthetic rate (μmol CO₂/m²/s).

Data statistical analysis. All were subjected to the analysis of variance (ANOVA) with the SAS software package. Appropriate standard errors of the means (S.E.) were calculated for presentation with table and bar diagram. The significance of the treatment effect was determined using the *F*-test, and to determine the significance of the means, least significant differences (*LSD*) were estimated at 5% probability level, and Duncan's multiple range test was used for comparing treatments within two or three factors combinations.

RESULTS AND DISCUSSION

Plant growth. Plants can absorb moderate atmospheric NH₃ as source of N, and can thus positively benefit in term of plant growth and biomass production. Atmospheric NH₃ can enter the leaves of higher plants almost exclusively through the stomata and is dissolved in the water film of the mesophyll cells to form NH₄⁺ as long as the ambient NH₃ concentrations exceed the mesophyll concentration (compensation point) (Fangmeier et al. 1994). Since NH₃ uptake by leaves occurred directly following stomatal conductance, the plant response to atmospheric NH₃ is dependent on environment factors, including crop cultivar and soil moisture availability (Rogers and Aneja 1980), internal CO₂ and NH₃ concentration, and plant water and fertilizer availability (Tatsuro et al. 2001). At the ambient NH₃ concentrations, its uptake by stem may have limited role in N-nutrition of higher plants. Enhanced growth was observed in many experiments within a reasonable range NH₃ concentrations, not approaching toxic level (Fangmeier et al. 1994). Enriched NH₃ concentrations can contribute to N-nutrition of plants, as was shown by Faller (1972) and Li et al. (2009).

Table 1. *F*-values of the effects of NH₃ level (NH₃), cultivar (Cv), N level (N) and their interactions on dry matter production (DM), total C accumulation (C_{tot}A) and total N accumulation (N_{tot}A)

Variation (g/plant)	N	Cv	NH ₃	N × Cv	NH ₃ × N	Cv × NH ₃	N × Cv × NH ₃
DM	176.55***	138.78***	0.97	22.46***	55.94***	2.70	0.40
C _{tot} A	4478.17***	2701.51***	78.84***	396.97***	388.39***	12.01**	22.70***
N _{tot} A	7718.18***	109.58***	1088.65***	292.19***	72.12***	78.82***	93.57***

P* < 0.05; *P* < 0.01; ****P* < 0.001

The current study supports the above conclusion as is evident from the positive effects of enriched atmospheric NH₃ concentration on plant growth in the low N (LN) solution. Fumigation of 1000 nl/L NH₃ greatly increased DM production in SD19 (cult. 1) with low N use efficiency (NUE) (by 18%) than that in NE5 (cult. 2) with high NUE (by 14%) above the ambient concentration, which better counteract the negative effect of LN supply on shoot DM (Figure 1). Moreover, the single factor of N level and cultivar together with the interactions of N level and cultivar or NH₃ level had significant effects on DM production only (Table 1). Nitrate-deprived plants and low N use efficient cultivar can benefit from the atmospheric N source, since shoot biomass production recovered from nitrate deprivation (Castro et al. 2005). Atmospheric NH₃ at a concentration of 4 µmol/L was therefore able to replace nitrate as nutrient to a considerable extent. This is in agreement with previous findings (Castro et al. 2005) and theoretical calculations on the possible contribution of NH₃ to the N budget of plants (Clement et al. 1997). In the high N (HN) solution, however, enriched atmospheric NH₃ reduced the DM production of cult. 2 (by 16%) with high NUE and non-significant effects of enriched atmospheric NH₃ on DM were found in cult. 1 with low NUE (*P* > 0.05) (Figure 2). Thereby, NH₃ assimilation under enriched atmospheric NH₃ under LN medium can improve plant growth. The canopy will respond to elevated atmospheric NH₃ concentration first, especially for the cultivar with low NUE in most natural environment with N limitation for plant biomass production (Fangmeier et al. 1994).

Accumulation of total C and total N (C_{tot}A and N_{tot}A). The absorption and utilization of NH₃ supplied to the plants was evaluated by calculating C_{tot}A and N_{tot}A, respectively. Ammonia (NH₃) uptake may cause an autocatalytic increase of additional NH₃ flux into the leaves by inducing stomatal opening via the internal CO₂ level (Van der Eerden et al. 1992). Long-term exposure of

plants to moderate atmospheric NH₃ concentration may stimulate increased C and N assimilation in plant, which affects the internal C and N status of the plant (Castro et al. 2005). In *Calluna vulgaris* (heather) and *Deschampsia flexuosa* (hair-grass), the N content increased four-fold after exposure to 0.1 mg/m³ NH₃ for 38 weeks. Varied increases of N_{tot}A induced by enriched atmospheric NH₃ were found in other plants such as conifers, tomato (*Lycopersicon esculentum*), *Arnica montana* L. and *Viola canina* L. (Van der Eerden et al. 1991, 1992, Dueck and Elderson 1992, Clement et al. 1997, Li et al. 2009). Our study showed that enriched atmospheric NH₃ concentration increased C_{tot}A and N_{tot}A in both cultivars in the LN solution, relative to counterparts exposed to the ambient concentration. The above increases in biomass C_{tot}A and N_{tot}A were greater for cult. 1 (by 16% and 38%, respectively) than that for cult. 2 (by 12% and 32%, respectively). In contrast, in the




Figure 2. Differential effects of enriched atmospheric NH₃ on dry matter production (g/plant) of SD19 (cult. 1) and NE5 (cult. 2) in the low and high N (LN and HN) medium. Data represent mean of ten replicates (*n* = 10). At the top of each column, different letters indicate significant differences between enriched atmospheric NH₃ (EAN-1000 nl/L) and control treatment (10 nl/L) with the same cultivar and N level




Figure 3. Differential effects of enriched atmospheric NH₃ on total C accumulation (C_{totA}) (g/plant) of SD19 (cult. 1) and NE5 (cult. 2) in the low and high N (LN and HN) medium. Data represent mean of ten replicates (*n* = 10). At the top of each column, different letters indicate significant differences between enriched atmospheric NH₃ (EAN-1000 nl/L) and control treatment (10 nl/L) with the same cultivar and N level




Figure 4. Differential effects of enriched atmospheric NH₃ on total N accumulation (N_{totA}) (g/plant) of SD19 (cult. 1) and NE5 (cult. 2) in the low and high N (LN and HN) medium. Data represent mean of ten replicates (*n* = 10). At the top of each column, different letters indicate significant differences between enriched atmospheric NH₃ (EAN-1000 nl/L) and control treatment (10 nl/L) with the same cultivar and N level

HN solution, enriched atmospheric NH₃ more decreased C_{totA} and N_{totA} of cult. 2 (15% and 18%, respectively) than those of cult. 1 (non-significant) (Figures 2–3). Additionally, the effects of NH₃ level, cultivar, N level and their interactions on C_{totA} and N_{totA} were all significant (Table 1). Such, enhancement of C_{totA} and N_{totA} may be significant for plants in natural environments if the NH₃ concentration rises, since N is a limit-

ing factor for plant biomass production in most natural habitats, especially for low NUE cultivars (Fangmeier et al. 1994, Li et al. 2009).

Photosynthesis. Increases of CO₂ assimilation together with net photosynthetic rate (P_n) and apparent quantum yield (AQY) enhancement resulted in lower intercellular CO₂ concentration (C_i) of plants. The modulation of CO₂ concentration in plants might be increase G_s (stomatal conductance),

Table 2. Differential effects of enriched atmospheric NH₃ on photosynthesis parameters of SD19 (cult. 1) and NE5 (cult. 2) in the low and high N medium

Treatment		P _n (μmol CO ₂ /m ² /s)	C _i (μmol/mol)	G _s (mol/m ² /s)	AQY (mol CO ₂ /mol)	R _d (μmol CO ₂ /m ² /s)
Low N (1.67 mmol/L) combined with different NH ₃ concentration						
Cult. 1	control	16.18 ± 0.89 ^b	269.2 ± 10.6 ^a	0.139 ± 0.016 ^b	0.078 ± 0.003 ^b	1.273 ± 0.180 ^a
	EAN	20.63 ± 1.55 ^a	195.2 ± 10.3 ^b	0.193 ± 0.018 ^a	0.095 ± 0.004 ^a	1.290 ± 0.220 ^a
Cult. 2	control	25.02 ± 2.52 ^b	204.2 ± 8.3 ^a	0.175 ± 0.014 ^b	0.093 ± 0.004 ^b	1.638 ± 0.120 ^a
	EAN	28.70 ± 2.33 ^a	163.4 ± 7.9 ^b	0.225 ± 0.016 ^a	0.108 ± 0.005 ^a	1.636 ± 0.180 ^a
High N (5.00 mmol/L) combined with different NH ₃ concentration						
Cult. 1	control	24.02 ± 1.12 ^a	179.3 ± 8.8 ^a	0.205 ± 0.014 ^a	0.099 ± 0.002 ^a	1.312 ± 0.159 ^a
	EAN	22.96 ± 1.03 ^a	188.3 ± 8.7 ^a	0.183 ± 0.014 ^a	0.091 ± 0.003 ^a	1.327 ± 0.156 ^a
Cult. 2	control	38.24 ± 2.22 ^a	130.0 ± 7.5 ^a	0.255 ± 0.018 ^a	0.123 ± 0.002 ^a	1.750 ± 0.220 ^a
	EAN	33.17 ± 2.03 ^b	153.0 ± 8.3 ^b	0.212 ± 0.016 ^b	0.105 ± 0.005 ^b	1.744 ± 0.250 ^a

C_i – intercellular CO₂ concentration; control – ambient NH₃ concentration (10 nl/L); G_s – stomatal conductance; AQY – apparent quantum yield; EAN – enriched atmospheric NH₃ concentration (1000 nl/L); P_n – net photosynthetic rate; R_d – dark respiration rate. Data represent mean of ten replicates (*n* = 10). Mean values followed by different letters within each column indicate significant differences at *P* < 0.05 between enriched atmospheric NH₃ fumigation and control treatment with the same cultivar and N supply

Table 3. *F*-values of the effects of NH₃ level (NH₃), cultivar (Cv), N level (N) and their interactions on photosynthesis parameters

Variation	N	Cv	NH ₃	N × Cv	NH ₃ × N	Cv × NH ₃	N × Cv × NH ₃
P _n (μmol CO ₂ /m ² /s)	445.20***	980.25***	2.29	32.44***	116.64***	13.11**	6.02*
C _i (μmol/mol)	10106.2***	10416.3***	2121.1***	49.81***	6664.2***	690.9***	112.6***
G _s (mol/m ² /s)	2036.6***	2908.9***	204.75***	16.29**	3844.7***	84.1***	38.9***
AQY (mol CO ₂ /mol)	677.60***	1524.60***	12.60**	35.00***	1177.40***	50.40***	22.40***
R _d (μmol CO ₂ /m ² /s)	16.54**	123.50***	0.01	14.43**	0.01	0.01	0.04

C_i – intercellular CO₂ concentration; AQY – apparent quantum yield; G_s – stomatal conductance; P_n – net photosynthetic rate; R_d – dark respiration rate. **P* < 0.05, ***P* < 0.01, ****P* < 0.001

which is beneficial for transmittion and absorption of atmospheric NH₃ (Fangmeier et al. 1994). Van Hove et al. (1989), Van der Eerden et al. (1992) and Van der Eerden and Pe'rez-Soba (1992) pointed out that increased NH₃ concentration enhanced P_n of *Pinus sylvestris*, poplar trees and Douglas fir, respectively. However, unaffected results in P_n of sunflower (*Helianthus annuus* L.) and *Acacia auriculaeform* were obtained by Berger et al. (1986) and Zhao et al. (2003), respectively. In this study, the authors stated that enriched NH₃ concentration increased P_n, G_s and AQY but decreased C_i in both cultivars in the LN treatment, especially for low NUE cultivar (cult. 1). On the contrary, in the HN treatment, enriched atmospheric NH₃ decreased P_n, G_s, AQY but increased C_i of cult. 2 only. Dark respiration rate (R_d) of both cultivars was not affected by increased atmospheric NH₃ irrespective of N treatment (Table 2). In addition, the significant effects of NH₃ level, cultivar, N level and their interactions on C_i, G_s and AQY were all obtained. With respect for P_n, non-significant effect of NH₃ level was merely calculated (*P* > 0.05). Nitrogen level, cultivar and their interaction had a significant effect on R_d only (Table 3). Similar result was drawn out by Van Hove et al. (1989) in *Populus euramericana* under 50–100 μg/m³ NH₃ concentration for 6–8 weeks. However, increase by 76% of R_d in *Populus euramericana* was found under 240 μg/m³ NH₃ concentration for three months. Thus, the boundary factors for efficient effects of NH₃ fumigation on R_d need to further clarify. The results here show that elevated atmospheric NH₃ concentration played a distinct role in the modulation of these photosynthesis parameters except for R_d of the plants, whose responses were dependent on crop cultivar and N supply in the medium (Fangmeier et al. 1994). These consistencies of photosynthesis physiological parameters change can be useful evaluation index

for plant growth under increased atmospheric NH₃ environment. Such atmospheric NH₃ absorption by the plant canopy might offset the N deficiency from root absorption, which showed the stronger use ability of atmospheric NH₃. The possible mechanism might be as following: relative deficiency of N nutrition of plant is beneficial for absorption from atmospheric NH₃ through leaves and synthesis of Rubisco which results in the increase of P_n (Zhao et al. 2003). Under high N medium, excessive NH₃ concentrations in cell possibly restrain Rubisco activity which reduces P_n of plants (Fangmeier et al. 1994).

REFERENCES

- Berger M.G., Klaus R.E., Fock H.P. (1986): Assimilation of gaseous ammonia by sunflower leaves during photosynthesis. *Australian Journal of Plant Physiology*, 113: 211–219.
- Bohme F., Merbach I., Weigel A., Russow R. (2003): Effect of crop type and crop growth on atmospheric nitrogen deposition. *Journal of Plant Nutrition and Soil Science*, 166: 601–605.
- Cape J.N., Jones M.R., Leith I.D., Sheppard L.J., Van Dijk N., Sutton M.A., Fowler D. (2008): Estimate of annual NH₃ dry deposition to a fumigated ombrotrophic bog using concentration-dependent deposition velocities. *Atmospheric Environment*, 42: 6637–6646.
- Castro A., Stulen I., De Kok L.J. (2005): Impact of atmospheric NH₃ deposition on plant growth and functioning – a case study with *Brassica oleracea* L. In: Omasa K., Nouchi I., De Kok L.J. (eds): *Plant Responses to Air Pollution and Global Change*. Springer-Verlag, Tokyo, 13–20.
- Clement J.M.A.M., Loorbach J., Meijer J., Hasselt P.R., Stulen I. (1997): The impact of atmospheric ammonia and temperature on growth and nitrogen metabolism of winter wheat. *Plant Physiology and Biochemistry*, 35: 395–404.
- Dueck T.A., Elderson J. (1992): Influence of NH₃ and SO₂ on the growth and competitive ability of *Arnica montana* L. and *Viola canina* L. *New Phytologist*, 122: 507–514.

- Guo J., Guo X.Y., Wang J.H., Zhang F.L. (2005): Characteristic parameters of light responses of corn varieties with different plant shapes. *Acta Botany Boreal-Occident Sinica*, 5: 1612–1617.
- Faller N. (1972): Schwefeldioxid, Schwefelwasserstoff, nitrose Gase und Ammoniak als ausschließliche S– bzw. N-Quellen der höheren Pflanzen. *Zeitschrift für Pflanzenernährung und Bodenkunde*, 131: 120–130.
- Fangmeier A., Hadwiger-Fangmeier A., Van der Eerden J.M. (1994): Effects of atmospheric ammonia on vegetation – a review. *Environmental Pollution*, 86: 43–82.
- Li S.Q., Zhao L., Shao M.A. (2004): Ammonia exchange between plant canopy and the atmosphere – a review. *Acta Botany Boreal-Occident Sinica*, 24: 2154–2162.
- Li J., Li S.Q., Liu Y., Chen X.L. (2009): Effects of increased ammonia on root/shoot ratio, grain yield and nitrogen use efficiency of two wheat varieties with various N supply. *Plant, Soil and Environment*, 55: 273–280.
- Paul W.L., Bert G.D. (1993): Open top chambers for exposing plant canopies to elevated CO₂ concentration and for measuring net gas exchange. In: Rozema J., Lambers H., van de Geijn S.C., Cambridge M.L. (eds): *Vegetatio*. Kluwer Academic Publishers, Belgium, 3–15.
- Rogers H.H., Aneja V.P. (1980): Uptake of atmospheric ammonia by selected plant species. *Environmental and Experimental Botany*, 20: 251–257.
- Sommer S.G., Jensen E.S., Schjrrong J.K. (1993): Leaf absorption of atmospheric ammonia emitted from pig slurry applied beneath the canopy of winter wheat. *Acta Agriculturae Scandinavica. Section B, Plant Soil Science*, 43: 21–24.
- Tatsuro N., Motohiro F., Yukiko D., Takeshi I. (2001): Effect of nitrogen load on growth, photosynthesis and nutrient status of *Cryptomeria japonica* and *Pinus densiflora* seedlings. *Trees*, 15: 453–461.
- Van der Eerden L.J.M., Dueck T.A., Berdowski J.J.M., Greven H., Van Dobben H.F. (1991): Influence of NH₃ and (NH₄)₂SO₄ on heathland vegetation. *Acta Botany Gallica*, 40: 281–297.
- Van der Eerden L.J., Lekkerkerk L.J., Smeulders S.M., Jansen A.E. (1992): Effects of atmospheric ammonia and ammonium sulphate on Douglas fir (*Pseudotsuga menziesii*). *Environmental Pollution*, 76: 1–9.
- Van der Eerden L.J.M., Pe'rez-Soba M. (1992): Physiological responses of *Pinus sylvestris* to atmospheric ammonia. *Trees*, 6: 48–53.
- Van Hove L.W.A., Van Kooten O., Adema E.H., Vredenberg W.J., Pieters G.A. (1989): Physiological effects of long term exposure to low and moderate concentrations of atmospheric NH₃ on poplar trees. *Plant, Cell and Environment*, 12: 899–908.
- Zhao P., Sun G.C., Zeng X.P., Cai X.A., Peng S.L. (2003): Variations of photosynthetic parameters in leaves of *Acacia auriculaeformis* grown in different nitrogen sources under increased atmospheric NH₃. *Acta Ecologica Sinica*, 23: 1386–1393.

Received on January 6, 2010

Corresponding author:

Li Shi-Qing, Northwest A & F University, State Key Laboratory of Soil Erosion and Dryland Farming, 712100 Yangling, Shannxi Province, P.R. China
 fax: + 86 29 870 16 171, e-mail: lishiqing.8888@yahoo.com.cn

Zhang Lixin and Qiang Hong contributed to this work equally