

## Forest management and snag characteristics in Northern Iran lowland forests

E. TAVANKAR<sup>1</sup>, R. PICCHIO<sup>2</sup>, A. LO MONACO<sup>2</sup>, A.E. BONYAD<sup>3</sup>

<sup>1</sup>*Department of Forestry, Khalkhal Branch, Islamic Azad University, Khalkhal, Iran*

<sup>2</sup>*Department of Agriculture, Forests, Nature and Energy, University of Tuscia, Viterbo, Italy*

<sup>3</sup>*Department of Forestry, Faculty of Natural Resources, University of Guilan, Soumesara, Iran*

**ABSTRACT:** Some snag characteristics were studied in three different Hyrcanian lowland forests. The highest snag density (38.4 stem·ha<sup>-1</sup>) was found in fully protected forests, and it was significantly higher than in selectively logged (23.7 stem·ha<sup>-1</sup>) and open access forests (8.8 stem·ha<sup>-1</sup>). The snag volume, diameter and height were also significantly higher in the fully protected forests. The number of snag species was recorded in fully protected (9), selectively logged (5) and open access forests (4). The snags were more evenly distributed among diameter, height and decay classes in the fully protected forest. The larger diameter snags (> 90 cm) and snags higher than 15 m in height were not found in the open access forest. The snags of decay class 5 had a density of 6.8 stem·ha<sup>-1</sup> in the fully protected forest, while they had a low density (1.5 stem·ha<sup>-1</sup>) in the selectively logged forest and they were not found in the open access forest. Target values of snags for managed lowland forests in Iran were defined in relation to management influences.

**Keywords:** biodiversity; broadleaved forests; Caspian forests; dead wood; Hyrcanian forest

Deadwood has a wide range of ecological values in forest ecosystems, offering habitats for many living organisms (BURSELL 2002, HUMPHREY et al. 2002; BRITZKE et al. 2003, LONSDALE et al. 2008; LUČAN et al. 2009; HANBERRY et al. 2012), providing carbon sequestration (ALLARD, PARK 2013; MATSUZAKI et al. 2013) and forest productivity preservation and also contributing to soil development and to nutrient cycles (LAIHO, PRESCOTT 1999; KIM et al. 2006; STRUKELJ et al. 2013). Decaying logs retain moisture and nutrients and play an important role for forest regeneration and for the maintenance of microhabitats (SZEWCZYK, SZWAGRZYK 1996). Logs also store energy and fix nitrogen (BRUNNER, KIMMINS 2003; YATSKOV et al. 2003; CREED et al. 2004). Furthermore, deadwood could reduce soil erosion in slope areas. The research indicates that many forest insects are kept at low levels by insectivorous birds and small mammals that eat insects during all or part of their life cycle (RAFFERTY et al. 1996). Deadwood

is also considered an important indicator, becoming relevant in the National Forest Inventories, in forest certification schemes and in some forest management plans (MCPFE 2007; RONDEUX, SANCHEZ 2010). The best practice guidelines for sustainable management often encourage the retention of a higher volume of deadwood in managed forests (BÖHL, BRÄNDLI 2007).

Snags are the standing trees dead for a natural process. They are an important environmental element and are essential for maintaining biodiversity in forest ecosystems (FERRIS, HUMPHREY 1999). Snags are originated by any possible factor that contributes to tree mortality, such as lightning, storm breakage, fire, disease, insects, drought, flooding, forestry practices, and so on (RAFFERTY et al. 1996; WOLF et al. 2004; BENDIX, COWELL 2010). Snags are not only the base of a food-chain but also they provide microhabitats for many living organisms, including fungi, epixylic lichens, bryophytes, invertebrates, birds,

mammals, reptiles, and amphibians (RUSSELL et al. 2006; WISDOM, BATE 2008; LARRIEU, CABANNETTES 2012; NASCIMBENE et al. 2013).

Wildlife use snags and downed logs for nesting, roosting, foraging, perching, or territorial displays (RABE et al. 1998; WISDOM, BATE 2008; LUČAN et al. 2009). Cavities in trees and snags provide suitable nesting sites for birds, bats and other wildlife species. In forest areas, an adequate and continuous availability should be ensured for preservation purposes (FAN et al. 2003; TREMBLAY et al. 2010; RUSSO et al. 2011; ULYSHEN 2011; GHADIRI KHANAPOSHTANI et al. 2013; REGNERY et al. 2013; ROST et al. 2013). Snags also contribute to ecological processes and decay dynamics (LINDENMAYER et al. 1997; GANEY, VOJTA 2005; STRUKELJ et al. 2013). Numerous wildlife functions are attributed to decaying wood as a source of food, nutrients, and cover for organisms at diverse trophic levels (SPIES et al. 1988; NASCIMBENE et al. 2013).

The potential benefits to wildlife from deadwood are dependent on several factors, size, species, level of decay, and location. Increasingly, snags have been studied in managed forests to determine snag dynamics (CHAMBERS, MAST 2005; RUSSELL, WEISKITTEL 2012), snag abundance and snag recruitment (BULL et al. 1990; GANEY 1999; HARRIS 1999; STEPHENS 2004; MORONI, HARRIS 2010), effects of orography (GALE 2000; CLARK et al. 2002) and effects of shelterwood cut (KENEFIG, NYLAND 2007). Snag and deadwood size and abundance are highly variable among regions and are dependent on forest type, successional stage, climate and forest management regimes (FAN et al. 2003, 2004; BÖHL, BRÄNDLI 2007; NAGAIKE 2009). Different management regimes affect snag richness (PEDLAR et al. 2002; GANEY, VOJTA 2005; STEPHENS, MOGHADDAS 2005; KENEFIG, NYLAND 2007). Forest management, timber harvest and human access can have substantial effects on snag density (WISDOM, BATE 2008; LARRIEU et al. 2012; PERRY, THILL 2013). The snags have become a major conservation issue in managed forest ecosystems. Managing for quality saw timber with the single tree selection system often reduces the number of cavity trees and snags, because they are removed under an intensive timber management regime (LARRIEU et al. 2012; PERRY, TILL 2013).

The total forest area of Iran is approximately 12.4 mil. ha, which makes only 7.3% of the total national surface (FAO 2005). The Hyrcanian forests of Iran (also known as Caspian forests) are located in the north of Iran and on the south coast of the Caspian Sea (Fig. 1). These forests cover



Fig. 1. The study area

2 mil. ha of land area and are broadleaf forests while the Iranian part is only commercial forests (FAO 2005). Approximately 60% of these forests are used for commercial purposes and the rest of them are more or less degraded (MARVIE MOHADJER 2006). These are the most valuable forests in Iran, both in commercial and in natural terms. The Hyrcanian forests are generally managed by selection cutting, with different logging methodologies. Data on existing densities and composition of snag populations are scarce for the Iranian lowland forests. The purpose of this study was to determine the density, size, level of decay and species of snags in different managing scenarios and the effect of forest management method on snag characteristics in the Hyrcanian lowland forests. The species, the volume, the distribution of species and diameter of snags in relatively undisturbed forests could provide a valuable baseline for sustainable management goals in managed forests and constitute a reference for restoring degraded woodland.

## MATERIAL AND METHODS

**Study area.** This study was carried out in the Hyrcanian lowland forest in Guilan province in the north of Iran. The study area is located between 37°37'0" to 37°39'0"N and 49°0'10" to 49°0'40"E and the altitude ranges between –20 to 50 m a.s.l. (Fig. 1). The climate is temperate based on the Demarton classification and it is very wet, with the mean annual temperature of 15.7°C and mean annual precipitation of 1,306 mm. Three forest stands with different

Table 1. Management characteristics of the study area

Characteristics	Fully protected	Selectively logged	Open access
Surface	52 ha	44 ha	53 ha
	no cutting		no cutting (because stand volume is low and area is degraded)
	no hunting	selection cutting (in each 10-year period – about 7–10% of stand volume – ground-based logging – skid trails)	
	no grassing		a lot of grassing
	no tourism		tourism
Management	no rural use for firewood, fuel	low grazing	rural use for firewood (fuel)
	protected by barbed wire	low rural use for firewood protected by barbed wire	protected by barbed wire (but many parts are open)

management methods (Table 1) were selected: (1) fully protected forest (the Forest Management Plans were prepared for these forests in 1963, from that year up to now no timber harvesting has been performed), (2) selectively logged forest, where selective harvest is allowed, the woodland was and it is protected from grazing, trampling and other forms of disturbance and (3) open access forest, where people have unrestricted access to forest resources.

**Data collection.** Data were collected by a systematic sampling design (STEPHENS 2004; KENEFIC, NYLAND 2007; HESSBURG et al. 2010). In each of the three managed forest areas (treatments) 30 circular plots (replicate) with a surface of 1,000 m<sup>2</sup> were taken at regular distances (100 m) from each other.

The number and the diameter at breast height (DBH) of all living trees and standing dead trees (snags) were recorded. The height of two living trees was measured selecting the nearest to the centre of the plot and the maximum DBH of the tree.

All snags ≥ 10 cm diameter at breast height and ≥ 2 m in height within plot boundaries were sampled. For all snags sampled, species, DBH, height, volume, percent bark cover, and decay class were recorded. The species of a snag was determined from bark characteristics. The DBH was recorded to the nearest centimetre using a DBH tape. Height was estimated to the nearest meter using a Suunto clinometer. Volume was calculated by Huber's Equation (1):

$$V = A_m H \quad (1)$$

where:

$V$  – volume (m<sup>3</sup>),

$A_m$  – mid-point cross-sectional area (m<sup>2</sup>)

$H$  – height (m).

Bark coverage was visually estimated to the nearest 5%.

Decay class of snag was determined in 5 classes (HOLLOWAY et al. 2007; CORACE et al. 2010):

DC1 – recently dead tree with intact tops and the majority of fine branching present;

DC2 – trees with loose bark, intact tops, and most of the fine branches;

DC3 – trees with < 50% of coarse branches and < 50% bark;

DC4 – trees with broken tops and few or no coarse branches;

DC5 – trees with broken tops and no coarse branches.

**Data analysis.** After checking for the normality (Kolmogorov-Smirnov test) and homogeneity of variance (Levene's test), distribution of tree species in the studied forests was compared using the Kruskal-Wallis test, which was also applied for the distribution of tree snag species; the averages of stand structure variables (DBH and height) were compared using a two-way ANOVA to test differences between the plots within each forest stand and differences between the three forest stands with different management methods (treatments); the averages of snag density and snag characteristics (volume, DBH and height) were compared using a two-way ANOVA to test differences between the plots within each forest stand and differences between the three forest stands with different management methods (treatments). Multiple comparisons were made by Duncan's test considered for this data typology as one of the most powerful tests (DAY, QUINN 1989) (significance at  $\alpha < 0.05$ ). Correspondence analysis was done considering the distribution of snags by DBH classes, height classes and decay classes for the three forest stand management. SPSS 19.0 software (IBM, New York, USA) was used for statistical analysis; also the results of the analysis were presented using descriptive statistics.

## RESULTS

The presence of boxwood (*Buxus hyrcana*) live trees is relevant only in the fully protected forest (Table 2), while it is absent in the open access forest. *Alnus glutinosa* is the most frequent species followed by *Parrotia persica* and *Quercus castaneifolia* both in selectively logged and in open access forest.

The species number decreases if disturbances increase. There are 18 species in the fully protected forest, 14 in the selected logged area and 11 in the open access forest (Table 2). The main dendrometric characteristics of the three treatments (DBH

and height) were analyzed to differences between the plots of every single treatment and between the three treatments. The ANOVA test shows any difference between the plots for each treatment ( $P > 0.05$ ,  $df = 2.27$ ) and significant differences between the three treatments ( $P < 0.05$ ,  $df = 2.27$ ) (Table 3). The same trend was observed in tree density, seedling density and canopy cover (Table 3). In the selectively logged forest, basal area and height are greater.

The species richness of the snags mirrored those of living trees in the different forests. The number of snag species in the fully protected, selectively logged and open access forest was 9, 5

Table 2. Distribution of tree species in the studied forests. From the Kruskal-Wallis test between the three areas there are significant differences in the distribution ( $P < 0.05$ )

	Fully protected		Selectively logged		Open access	
	(indd·ha <sup>-1</sup> )	(%)	(indd·ha <sup>-1</sup> )	(%)	(indd·ha <sup>-1</sup> )	(%)
<b>Tree species</b>						
<i>Buxus hyrcana</i> Pojark.	91.6	47.9	6.4	3.8	0	0
<i>Parrotia persica</i> C.A. Meyer	27.9	14.6	31.5	18.6	22.4	18.5
<i>Quercus castaneifolia</i> Gled.	26.3	13.7	31.4	18.5	21.4	17.7
<i>Alnus glutinosa</i> L.	21.1	11.0	36.0	21.2	30.7	25.3
<i>Pterocarya fraxinifolia</i> Lam.	7.0	3.7	12.1	7.1	14.1	11.6
<i>Carpinus betulus</i> L.	5.2	2.7	22.5	13.3	14.8	12.2
<i>Acer insigne</i> Boiss.	2.2	1.2	14.0	8.3	12.1	10.0
<i>Acer cappadocicum</i> Gled.	2.2	1.2	7.6	4.5	0	0
<i>Zelkova caprinifolia</i> (Pall.) Dipp.	2.1	1.1	6.1	3.6	0	0
<i>Mespilus germanica</i> L.	2.0	1.0	0.5	0.3	2.4	2.0
<i>Diospyrus lotus</i> L.	1.1	0.6	0.5	0.3	0	0
<i>Ulmus minor</i> Miller.	0.7	0.4	0.4	0.2	0	0
<i>Gleditsia caspica</i> Desf.	0.6	0.3	0	0	1.6	1.3
<i>Albizia julibrissin</i> Durazz.	0.5	0.3	0	0	1.1	0.9
<i>Prunus avium</i> L.	0.3	0.2	0.2	0.1	0.6	0.5
<i>Populus nigra</i> L.	0.2	0.1	0	0	0	0
<i>Fraxinus excelsior</i> L.	0.2	0.1	0.2	0.1	0	0
<i>Ficus carica</i> L.	0.1	0.1	0	0	0.5	0.4
<b>Snag species</b>						
<i>Buxus hyrcana</i> Pojark.	17.8	46.3	–	–	–	–
<i>Quercus castaneifolia</i> Gled.	7.1	18.5	3.6	15.2	–	–
<i>Carpinus betulus</i> L.	4.1	10.7	10.8	45.6	–	–
<i>Alnus glutinosa</i> L.	3.0	7.8	3.0	12.7	1.5	17.0
<i>Acer insigne</i> Boiss.	2.7	7.0	–	–	–	–
<i>Parrotia persica</i> C.A. Meyer	1.8	4.7	4.5	18.9	2.0	22.7
<i>Pterocarya fraxinifolia</i> (Lam.	1.0	2.6	–	–	–	–
<i>Mespilus germanica</i> L.	0.8	2.1	1.8	7.6	2.6	29.6
<i>Gleditsia caspica</i> Desf.	0.1	0.3	–	–	2.7	30.7

Table 3. Stand structure variables (mean  $\pm$  standard deviation)

Stand characteristics	Fully protected	Selectively logged	Open access
Stand basal area ( $\text{m}^2\cdot\text{ha}^{-1}$ )	13.3 $\pm$ 4.2	17.9 $\pm$ 2.9	12.5 $\pm$ 3.2
Tree density ( $\text{stem}\cdot\text{ha}^{-1}$ )	191.3 $\pm$ 22.2	169.4 $\pm$ 8.7	121.1 $\pm$ 16.2
Seedling density ( $\text{stem}\cdot\text{ha}^{-1}$ )	447.1 $\pm$ 32.7	322.3 $\pm$ 38.1	86.5 $\pm$ 19.5
Tree height (m)	16.1 $\pm$ 4.1	21.2 $\pm$ 4.2	15.5 $\pm$ 5.4
Canopy cover (%)	83.1 $\pm$ 8.4	78.0 $\pm$ 7.9	59.9 $\pm$ 6.9

and 4, respectively (Table 2). *Buxus hyrcana* had the highest snag density ( $17.8 \text{ stem}\cdot\text{ha}^{-1}$ ) in the fully protected forest, where it is the most representative species, while it was not found in the selectively logged and open access forest, where boxwood was sporadic or absent among the living trees. Three species (*Buxus hyrcana*, *Quercus castaneifolia* and *Carpinus betulus*) had the greater spread representing 75.5% of snags; these species also accounted for nearly 65% of living trees in the fully protected forest. In particular boxwood showed the same frequency both in the living trees and in snags.

*Carpinus betulus* had the highest snag density ( $10.8 \text{ stem}\cdot\text{ha}^{-1}$ ) in the selectively logged forest. This species represents 45.6% of the snags while only 13.3% of the living trees. *Gleditsia caspica* had the highest snag density ( $2.7 \text{ stem}\cdot\text{ha}^{-1}$ ) in the open access forest although the frequency in living trees was only 1.3%.

The density of snags was 38.4, 23.7 and  $8.8 \text{ stem}\cdot\text{ha}^{-1}$  in the fully protected, selectively logged and open access forest, respectively (Table 4). The snags represented 20, 14 and 7% of the living trees, respectively. The average volume of snags in the fully protected, selectively logged and open access forest was found to be 17.3, 9.4 and  $1.1 \text{ m}^3\cdot\text{ha}^{-1}$  (Table 4). The ANOVA and Duncan's test showed that the snag density and the snag volume are significantly higher in fully protected forest than in selectively logged forest ( $P < 0.01$ ), and

they are significantly higher in selectively logged forest than in open access forest ( $P < 0.01$ ).

The snag diameter in fully protected forest ( $37.7 \pm 9.9 \text{ cm}$ ) was significantly higher than in selectively logged ( $22.2 \pm 6.0 \text{ cm}$ ) and in open access ( $13.5 \pm 3.2 \text{ cm}$ ) forests (Table 4 and Fig. 2). Moreover diameter, snag height and snag volume in fully protected forest were significantly higher than in selectively logged and open access forests (Table 4 and Fig. 2). The mean of snag height in fully protected, selectively logged and open access forests measured 14.5, 10.1 and 6.3 m, respectively (Table 4). The average volume of snag in fully protected, selectively logged and open access forests measured 0.47, 0.36 and  $0.09 \text{ m}^3$  (Table 4).

The distribution of snags by diameter classes is shown in Table 5. The results indicated that the snags were more evenly distributed among diameter classes in the fully protected forest than in the selectively logged and open access forests (Fig. 2). The large and very large diameter snags ( $\text{DBH} > 60 \text{ cm}$ ) had the highest density ( $15.9 \text{ stem}\cdot\text{ha}^{-1}$ ) in fully protected forest,  $2.3 \text{ stem}\cdot\text{ha}^{-1}$  in selectively logged forest and were not observed in open access forest (Table 5 and Fig. 2). Small snags ( $\text{DBH} < 30 \text{ cm}$ ) had the higher frequency (82.9%) in the open access forest (Fig. 2). Even in selectively logged forest the small snags were more abundant than the other diameter classes. Snags  $< 30 \text{ cm DBH}$  are less valuable as wildlife habitats than larger snags. The very large diameter snags ( $\text{DBH} > 90 \text{ cm}$ ) were not found

Table 4. Density, volume and average parameters of snags (mean  $\pm$  standard deviation) in the studied forests (different letters indicate statistically significant differences by Duncan's test at  $\alpha = 0.05$ )

	Fully protected	Selectively logged	Open access
<b>Snag parameters</b>			
Density ( $\text{stem}\cdot\text{ha}^{-1}$ )	38.4 $\pm$ 7.2 <sup>a</sup>	23.7 $\pm$ 5.6 <sup>b</sup>	8.8 $\pm$ 2.5 <sup>c</sup>
Volume ( $\text{m}^3\cdot\text{ha}^{-1}$ )	17.3 $\pm$ 5.1 <sup>a</sup>	9.4 $\pm$ 2.9 <sup>b</sup>	1.1 $\pm$ 0.3 <sup>c</sup>
<b>Snag average parameters</b>			
Diameter (cm)	37.7 $\pm$ 9.9 <sup>a</sup>	22.2 $\pm$ 6.0 <sup>b</sup>	13.5 $\pm$ 3.2 <sup>c</sup>
Height (m)	14.5 $\pm$ 3.4 <sup>a</sup>	10.1 $\pm$ 2.2 <sup>b</sup>	6.3 $\pm$ 1.3 <sup>c</sup>
Volume ( $\text{m}^3$ )	0.47 $\pm$ 0.13 <sup>a</sup>	0.36 $\pm$ 0.10 <sup>b</sup>	0.09 $\pm$ 0.02 <sup>c</sup>




Fig. 2. Correspondence analysis considering the distribution of snags by DBH classes (d), height classes (h) and decay classes (DC) (solid line) for the three types of forest management (dotted line)

in selectively logged and open access forests, while these snags had the density of  $6.1 \text{ stem} \cdot \text{ha}^{-1}$  in the fully protected forest (Fig. 2).

The distribution of snags by height classes is shown in Table 5 and Fig. 2. The snag height was more evenly distributed among classes in the fully protected forest than in the selectively logged and open access forests. The snags higher than 15 m had the highest density ( $4.5 \text{ stem} \cdot \text{ha}^{-1}$ ) in the fully protected forest (Fig. 2). The snags higher than 10 m

were not found in the open access forest where the shorter class showed a frequency of more than 70%.

The higher snags (height > 10 m) were found in fully protected and selectively logged forests. The density was  $13.2$  and  $3.9 \text{ stem} \cdot \text{ha}^{-1}$  in fully protected and selectively logged forest, respectively, but with relevant difference in frequency (Fig. 2). It was about double in the fully protected forest.

The distribution of snags by decay classes is shown in Table 6 and Fig. 2. The level of snag decay

Table 5. Distribution of snags by parameters in the studied forests

	Fully protected		Selectively logged		Open access	
	(indd·ha <sup>-1</sup> )	(%)	(indd·ha <sup>-1</sup> )	(%)	(indd·ha <sup>-1</sup> )	(%)
<b>Snag diameter (cm)</b>						
< 30 (d <sub>0</sub> )	8.0	20.8	16.3	68.8	7.3	82.9
31 – 60 (d <sub>1</sub> )	14.5	37.8	5.1	21.5	1.5	17.1
61 – 90 (d <sub>2</sub> )	9.8	25.5	2.3	9.7	–	–
> 91 (d <sub>3</sub> )	6.1	15.9	–	–	–	–
<b>Snag height (m)</b>						
< 5 (h <sub>0</sub> )	13.2	34.4	4.4	18.6	6.2	70.4
5 – 10 (h <sub>1</sub> )	12.0	31.2	15.4	65.0	2.6	29.6
10 – 15 (h <sub>2</sub> )	8.7	22.7	2.5	10.5	–	–
> 15 (h <sub>3</sub> )	4.5	11.7	1.4	5.9	–	–
<b>Snag decay class</b>						
DC1	9.8	25.5	5.4	22.8	5.1	57.9
DC2	7.7	20.1	8.3	35.0	3.7	42.1
DC3	7.0	18.2	5.9	24.9	–	–
DC4	7.1	18.5	2.6	11.0	–	–
DC5	6.8	17.7	1.5	6.3	–	–

is another important factor to consider in management decisions. Snags were more evenly distributed among the decay classes in the fully protected and in the selectively logged forests than in the open access forest.

Snag decay classes DC3, DC4 and DC5 were not found in the open access forest, while these classes had 20.9 and 10.0 stem·ha<sup>-1</sup> in the fully protected and in the selectively logged forest (Fig. 2). The snags in decay class DC5 (very decayed snags) had a density of 6.8 stem·ha<sup>-1</sup> in the fully protected forest, while their density in the selectively logged forest was 1.5 stem·ha<sup>-1</sup> (Fig. 2).

## DISCUSSION

Dead wood and snags particularly may be considered a proxy of biodiversity although the knowledge acquired on the abundance and attributes of snags is not directly transferable to other forest contexts for the well-known differences due to climate, site, forest types and history. Forest managers need biodiversity indicators that are easy to use. The effect of forest management was surveyed focusing on snags. The snag attributes (species, density, size, and level of decay) were studied in three different forests (fully protected, selectively logged and open access) in the Hyrcanian lowland forests. Forest management has caused changes in stand composition, tree density, basal area, height, seedling density, and canopy cover. Forest management has also influenced the attributes of snags. Snags were more diffused in fully protected forest than in forests managed in another way (SEFIDI et al. 2009). The reduction of snag density was higher in the open access forest, where cutting of snags is allowed. Generally, snags were removed by rural people for fuelwood. The removal of snags can negatively impact wildlife populations that are dependent on them as essential habitat components. In addition, our results on the snag density and human access are similar to those of WISDOM and BATE (2008) on pine and larch in the Rocky Mountains and to those of BATE et al. (2007) obtained in the Northeast Oregon. RUSSELL et al. (2012) found that snag density was lower in intensively managed stands and that height and DBH were greater in unmanaged stands. The results indicated that the range of snag species depends on the specific composition of the stand, although the frequency does not. SEFIDI and MARVIE MOHADJER (2009) found similar results in a beech stand in Caspian forest in the north of Iran. The snag density, especially of large-size snags,

was significantly higher in the fully protected forest than in the selectively logged and open access forests. RUSSEL and WEISKITTEL (2012) found that the probability of snag survival was dependent on species and DBH. LÖHMUS et al. (2013) noted that snags of deciduous trees had a higher probability of trunk breakage during the time, particularly the higher and the thinner ones, with the consequence of decay acceleration due to the ground contact. From an ecological point of view, snags of larger diameter are particularly important, due to the richness of microhabitats per tree (REGNERY et al. 2013). In addition, the diameter at breast height and height of snags determine which species will use the snag for nesting (THOMAS et al. 1979). GHADIRI KHANAPOSHTANI et al. (2012) underlined that high volumes of coarse woody debris, especially large snags, and dense canopy cover are habitat requirements of the black woodpecker, *Dryocopus martius*, in Hyrcanian Forests.

The level of snag decay is another important factor driving management decisions (JONSSON et al. 2005; NASCIMBENE et al. 2013). Suitable density and different stages of decay of snags are critical to the preservation of biodiversity and the constant functionality of forest ecosystems (DE LONG et al. 2008). For example, soft snags are most often used for nesting whereas hard snags are most often used for foraging (FAN et al. 2003). In order to maximize the wildlife benefits, a variety of decay snag classes is necessary: snags from harder trees newly died still with bark and old decayed snags debarked.

Defects and indicators of wood decay are considered when trees are selected for cutting, due to the maximization of the future value of forest (TRIMBLE et al. 1974). Standing dead or dying trees are usually harvested to avoid pest problems and fire hazards and to maximize the commercial value of the harvest. Intensively managed forest or productive forests are subjected to disturbances caused by logging operations. Damage to residual trees may increase the stand mortality both during the selection cutting operations (TAVANKAR et al. 2011, 2013) and during thinning (PICCHIO et al. 2011, 2012; MARCHI et al. 2014) with a consequent increase of dying or standing dead trees, which take on the configuration of snags during time.

Snags are a significant structural component of forest ecosystems. They preserve biodiversity, as many dead wood dependent organisms are confined to snags during their life cycle (NILSSON et al. 2002). Although density or volume thresholds of snags are not yet established for Iranian for-

est types, some authors have addressed this issue. MARVIE MOHADJER et al. (2009) found a density of 19.3 snags (DBH  $\geq$  7.5cm) per hectare for a selectively logged stand in the northern forests of Iran. It is important to provide snags of various sizes. A selectively logged stand had a density of 2.1 to 2.5 snag·ha<sup>-1</sup> (DBH  $\geq$  50 cm) in the Hyrcanian forests (SEFIDI et al. 2009). In Caspian Hyrcanian mixed forest within the Alborz mountains GHADIRI KHANAPOSHTANI et al. (2013), studying the effect of forest logging on avian communities, found 6.66 (+ 0.37 SD) snag·ha<sup>-1</sup> in a never harvested area and 4.32 (+ 0.42 SD) snag·ha<sup>-1</sup> in a logged area. In Mazandaran Province, SEFIDI and MARVIE MOHADJER (2010) examined the amount of dead wood in mixed beech forests in late, middle and early successional stages. The late successional forest had a larger amount of logs, snags and stumps than the other two forests. The snag volume differed between the late (15.62 m<sup>3</sup>·ha<sup>-1</sup>) and middle successional forest (0.48 m<sup>3</sup>·ha<sup>-1</sup>), and it was 16.60 m<sup>3</sup>·ha<sup>-1</sup> in the early successional stand.

The results of this study showed that the characteristics of snags were affected by forest management, these results were confirmed by KENEFIC and NYLAND (2007) and by WISDOM and BATE (2008). Timber harvest and human access can have substantial effects on snag density (WISDOM, BATE 2008). It is now widely accepted that forests should be managed in an ecologically sustainable manner (KOHM, FRANKLIN 1997). For the first time this study contributes to the definition of the target values of snags for managed lowland forests in Iran and highlights the management influences on snag density, species frequency and distribution in decay classes. The fully protected forest could represent a guide for snag management in selectively logged and open access forests, in the lowland Hyrcanian forest.

## CONCLUSION

Today, the Hyrcanian lowland forests of Iran are depleting rapidly due to the population growth and associated socio-economic problems, industrial development and urbanism (POORZADY, BAKHTIARI 2009). Consequently, the reduction in snag density would likely decrease the biodiversity in forest ecosystems. Snags provide habitats for foraging, nesting, resting or cover for many species of wildlife, as well as a source of coarse woody debris important in forest succession. Characteristics of snags, including the stage of decay, density, size, and spe-

cies influence their use by wildlife. A reduction in the density of snags in selectively logged and open access forests may have negative consequences for wildlife. A continuous supply of snags and downed logs must be preserved to sustain wildlife populations that depend on these resources. Recommendations for “management of snags” include long rotations of forest stands and retaining of dying trees, dead standing trees and snags during harvest.

## References

- ALLARD J., PARK A. (2013): Woody debris volumes and carbon accumulation differ across a chronosequence of boreal red pine and jack pine stands. *Canadian Journal of Forest Research*, **43**: 768–775.
- BATE L.J., WISDOM M.J., WALES B.C. (2007): Snag densities in relation to human access and associated management factors in forests of northeastern Oregon, USA. *Landscape and Urban Planning*, **80**: 278–291.
- BENDIX J., COWELL C.M. (2010): Fire, floods and woody debris: interactions between biotic and geomorphic processes. *Geomorphology*, **116**: 297–304.
- BÖHL J., BRÄNDLI U.B. (2007): Deadwood volume assessment in the third swiss national forest inventory: methods and first results. *European Journal of Forest Research*, **126**: 449–457.
- BRITZKE E.R., HARVEY M.J., LOEB S.C. (2003): Indiana bat, *Myotis sodalis*, maternity roosts in the southern United States. *Southeastern Naturalist*, **2**: 235–242.
- BRUNNER A., KIMMINS J.P. (2003): Nitrogen fixation in coarse woody debris of *Thuja plicata* and *Tsuga heterophylla* forests on northern Vancouver Island. *Canadian Journal of Forest Research*, **33**: 1670–1682.
- BULL E.L., HOLTHAUSEN R.S., MARX D.B. (1990): How to determine snag density? *Western Journal of Applied Forestry*, **5**: 56–58.
- BULL E.L., PARKS C.G., TORGERSEN T.R. (1997): Trees and Logs Important to Wildlife in the interior Columbia River Basin. Portland, USDA Forest Service: 391.
- BURSELL J. (2002): Winter abundance of hole-nesting birds in natural and managed woods of Zealand (Denmark). *Acta Ornithologica*, **37**: 67–74.
- CHAMBERS C.L., MAST J.N. (2005): Ponderosa pine snag dynamics and cavity excavation following wildlife in northern Arizona. *Forest Ecology and Management*, **216**: 227–240.
- CLARK D.B., CLARK D.A., BROWN S., OBERBAUER S.F., VELDKAMP E. (2002): Stocks and flows of coarse woody debris across a tropical rain forest nutrient and topography gradient. *Forest Ecology and Management*, **164**: 237–248.
- CORACE R.G., SEEFFELT N.E., GOEBEL P.C., SHAW H.L. (2010): Snag longevity and decay class development in a recent jack pine clearcut in Michigan. *Northern Journal of Applied Forestry*, **27**: 125–131.



- CREED I.F., MORRISON D.L., NICHOLAS N.S. (2004): Is coarse woody debris a net sink or source of nitrogen in the red spruce–Fraser fir forest of the southern Appalachians, U.S.A.? *Canadian Journal of Forest Research*, **34**: 716–727.
- DAY R.W., QUINN G.P. (1989): Comparisons of treatments after an analysis of variance. *Ecological Monograph*, **54**: 433–463.
- DE LONG S.C., SUTHERLAND G.D., DANIELS L.D., HEEMSKERK B.H., STORAUNET K.O. (2008): Temporal dynamics of snags and development of snag habitats in wet spruce–fir stands in east-central British Columbia. *Forest Ecology and Management*, **255**: 3613–3620.
- FAN Z., LARSEN D.R., SHILLEY S.R., THOMPSON F.R. (2003): Estimating cavity tree abundance by stand age and basal area, Missouri, USA. *Forest Ecology and Management*, **179**: 231–242.
- FAN Z., SHIFLEY S.R., THOMPSON F.R., LARSEN D.R. (2004): Simulated cavity tree dynamics under alternative timber harvest regimes. *Forest Ecology and Management*, **193**: 399–412.
- FAO (2005): Global Forest Resources Assessment, Country Reports, Islamic Republic of Iran. Rome, FRA: 42.
- FERRIS R., HUMPHREY J.W. (1999): A review of potential biodiversity indicators for application in British forests. *Forestry*, **72**: 313–328.
- GALE N. (2000): The aftermath of tree death: coarse woody debris and the topography in four tropical rain forests. *Canadian Journal of Forest Research*, **30**: 1489–1493.
- GANEY J.L. (1999): Snag density and composition of snag populations on two National Forests in northern Arizona. *Forest Ecology and Management*, **117**: 169–178.
- GANEY J.L., VOJTA S.C. (2005): Changes in snag populations in northern Arizona mixed-conifer and Ponderosa pine forests 1997–2002. *Forest Science*, **51**: 396–405.
- GHADIRI KHANAPOSHTANI M., KABOLI M., KARAMI M., ETEMAD V. (2012): Effect of habitat complexity on richness, abundance and distributional pattern of forest birds. *Environmental Management*, **50**: 296–303.
- GHADIRI KHANAPOSHTANI M., KABOLI M., KARAMI M., ETEMAD V., BANIASADI S. (2013): Effects of logged and unlogged forest patches on avifaunal diversity. *Environmental Management*, **51**: 750–758.
- HANBERRY B.B., HANBERRY P., DEMARAIS S., JONES J.C. (2012): Importance of residual trees to birds in regenerating pine plantations. *IForest*, **5**: 108–112.
- HARRIS R.B. (1999): Abundance and characteristics of snags in western montana forests. Moscow, USDA Forest Service: 31.
- HESSBURG P.F., POVAK N.A., SALTER R.B. (2010): Thinning and prescribed fire effects on snag abundance and spatial pattern in an eastern cascade range dry forest, Washington, USA. *Forest Science*, **56**: 74–87.
- HOLLOWAY G.L., CASPERSEN J.P., VANDERWEL M.C., NAYLOR B.J. (2007): Cavity tree occurrence in hardwood forests of central Ontario. *Forest Ecology and Management*, **239**: 191–199.
- HUMPHREY J.W., DAVEY S., PEACE A.J., FERRIS R., HARDING K. (2002): Lichens and bryophyte communities of planted and semi-natural forests in Britain: the influence of site type, stand structure and deadwood. *Biological Conservation*, **107**: 165–180.
- JONSSON B.G., KRUYSS N., RANIUS T. (2005): Ecology of species living on dead wood – lessons for dead wood management. *Silva Fennica*, **39**: 289–309.
- KENEFIC L.S., NYLAND R.D. (2007): Cavity trees, snags, and selection cutting: a northern hardwood case study. *Northern Journal of Applied Forestry*, **24**: 192–196.
- KIM R., SON Y., LIM J.H., LEE I.K., SEO K.W., KOO J.W., NOH N.J., RYU S., HONG S.K., IHM B.S. (2006): Coarse woody debris mass and nutrients in forest ecosystems of Korea. *Ecological Research*, **21**: 819–827.
- KOHM K., FRANKLIN J.F. (1997): Creating a Forestry in 21<sup>st</sup> century. Covelo, Island Press: 491.
- LAIHO R., PRESCOTT C.E. (1999): The contribution of coarse woody debris to carbon, nitrogen, and phosphorous cycles in three Rocky Mountain coniferous forests. *Canadian Journal of Forest Research*, **29**: 1592–1603.
- LARRIERUE L., CABANNETTES A. (2012): Species, live status, and diameter are important tree features for diversity and abundance of tree microhabitats in subnatural montane beech–fir forests. *Canadian Journal of Forest Research*, **42**: 1433–1445.
- LARRIERUE L., CABANNETTES A., DELARUE A. (2012): Impact of silviculture on dead wood and on the distribution and frequency of tree microhabitats in montane beech–fir forests of the Pyrenees. *European Journal of Forest Research*, **131**: 773–786.
- LINDENMAYER D.B., CUNNINGHAM R.B., DONNELLY C.F. (1997): Decay and collapse of trees with hollows in Eastern Australian forests: impacts on arboreal marsupials. *Ecological Application*, **7**: 625–641.
- LONSDALE D., PAUTASSO M., HOLDENRIEDE O. (2008): Wood-decaying fungi in the forest: conservation needs and management options. *European Journal of Forest Research*, **127**: 1–22.
- LÖHMUS A., KRAUT A., ROSENVALD R. (2013): Dead wood in clearcuts of semi-natural forests in Estonia: sitetype variation, degradation, and the influences of tree retention and slash harvest. *European Journal of Forest Research*, **132**: 335–349.
- LUČAN R.K., HANÁK V., HORÁČEK I. (2009): Long-term reuse of tree roosts by European forest bats. *Forest Ecology and Management*, **258**: 1301–1306.
- MARCHI E., PICCHIO R., SPINELLI R., VERANI S., VENANZI R., CERTINI G. (2014): Environmental impact assessment of different logging methods in pine forests thinning. *Ecological Engineering*, **70**: 429–436.
- MARVIE MOHADJER M. (2006): Silviculture in Iran. Teheran, Tehran University Press: 387.
- MARVIE MOHADJER M., ZOBEIRI M., ETEMAD V., JOUR GHOLAMI M. (2009): Performing the single selection method at compartment level and necessity for full in-

- ventory of tree species (Case study: Gorazbon district in Kheyroud Forest). *Journal of Iranian Natural Resources*, **61**: 889–908.
- MATSUZAKI E., SANBORN P., FREDEEN A.L., SHAW C.H., HAWKINS C. (2013): Carbon stocks in managed and unmanaged old-growth western redcedar and western hemlock stands of Canada's inland temperate rainforests. *Forest Ecology and Management*, **297**: 108–119.
- MCPFE (2007): State of Europe's forests 2007 – The MCPFE report on sustainable forest management in Europe. In: 5<sup>th</sup> Ministerial Conference on the Protection of Forests in Europe. Forests for Quality of Life. Warsaw, 5.–7. November 2007. Warsaw, Ministerial Conference on the Protection of Forests in Europe: 247.
- MORONI M.T., HARRIS D.D. (2010): Snag frequency, diameter and species distribution and input rate in Newfoundland boreal forests. *Forestry*, **83**: 229–244.
- NAGAIKE T. (2009): Snag abundance and species composition in a managed forest landscape in central Japan composed of *Larix kaempferi* plantation and secondary broadleaf forests. *Silva Fennica*, **43**: 755–766.
- NASCIMBENE J., THOR G., NIMIS P.L. (2013): Effects of forest management on epiphytic lichens in temperate deciduous forests of Europe – a review. *Forest Ecology and Management*, **298**: 27–38.
- NILSSON S.G., NIKLASSON M., HEDIN J., ARONSSON G., GUTOWSKI M., LINDER P., LJUNGBERG H., MIKUSINSKI G., RANIUS T. (2002): Densities of large living and dead trees in old-growth temperate and boreal forests. *Forest Ecology and Management*, **161**: 189–204.
- PEDLAR J.H., PEARCE J.L., VENIER L.A., MCKENNEY D.W. (2002): Coarse woody debris in relation to disturbance and forest type in boreal Canada. *Forest Ecology and Management*, **158**: 189–194.
- PERRY R.W., THILL R.E. (2013): Comparison of snag densities among regeneration treatments in mixed pine-hardwood forests. *Canadian Journal of Forest Research*, **43**: 619–626.
- PICCHIO R., MAGAGNOTTI N., SIRNA A., SPINELLI R. (2012): Improved winching technique to reduce logging damage. *Ecological Engineering*, **47**: 83–86.
- PICCHIO R., NERI F., MAESANO M., SAVELLI S., SIRNA A., BLASI S., BALDINI S., MARCHI E. (2011): Growth effects of thinning damage in a Corsican pine (*Pinus laricio* Poir.) stand in central Italy. *Forest Ecology and Management*, **262**: 237–243.
- POORZADY M., BAKHTIARI F. (2009): Spatial and temporal changes of Hyrcanian forest in Iran. *IForest*, **2**: 198–206.
- RABE M.J., MORRELL T.E., GREEN H., DEVOS J.C., MILLER C.R. (1998): Characteristics of ponderosa pine snag roosts used by reproductive bats in northern Arizona. *Journal of Wildlife Management*, **62**: 612–621.
- RAFFERTY D., MASTERS R., GREEN C. (1996): Snags, cavity trees and downed logs. *Wildlife Management Notes*, **4**: 1–8.
- REGNERY B., PAILLET Y., COUVET D., KERBIRIOU C. (2013): which factors influence the occurrence and density of tree microhabitats in mediterranean oak forests? *Forest Ecology and Management*, **295**: 118–125.
- RONDEUX J., SANCHEZ C. (2010): Review of indicators and field methods for monitoring biodiversity within national forest inventories. Core variable: Deadwood. *Environmental Monitoring and Assessment*, **164**: 617–630.
- ROST J., HUTTO R.L., BROTONS L., PONS P. (2013): Comparing the effect of salvage logging on birds in the Mediterranean Basin and the Rocky Mountains: common patterns, different conservation implications. *Biological Conservation*, **158**: 7–13.
- RUSSELL R.E., SAAB V.A., DUDLEY J.G., ROTELLA J.J. (2006): Snag longevity in relation to wildfire and post fire salvage logging. *Forest Ecology and Management*, **232**: 179–187.
- RUSSELL M.B., KENEFIC L.S., WEISKITTE A.R., PUHLICK J.J., BRISSETTE J.C. (2012): assessing and modeling standing deadwood attributes under alternative silvicultural regimes in the acadian forest region of Maine, USA. *Canadian Journal of Forest Research*, **42**: 1873–1883.
- RUSSELL M.B., WEISKITTE A.R. (2012): Assessing and modeling snag survival and decay dynamics for the primary species in the Acadian forest of Maine, USA. *Forest Ecology and Management*, **284**: 230–240.
- RUSO D., CISTRONE L., GARONNA A.P. (2011): Habitat selection by the highly endangered long-horned beetle *Rosalia alpina* in Southern Europe: a multiple spatial scale assessment. *Journal of Insect Conservation*, **15**: 685–693.
- SEFIDI K., MARVI MOHADJER M.R. (2009): Amount and quality of dead trees (snag and logs) in a mixed beech forest with different management histories. *Journal of Forest and Wood Product*, **62**: 191–202.
- SEFIDI K., MARVIE MOHADJER M.R. (2010): Characteristics of coarse woody debris in successional stages of natural beech (*Fagus orientalis*) forests of Northern Iran. *Journal of Forest Science*, **56**: 7–17.
- SPIES T.A., FRANKLIN J.F., THOMAS T.B. (1988): Coarse woody debris in Douglas-fir forests of western Oregon and Washington. *Ecology*, **69**: 689–702.
- STEPHENS S.L. (2004): Fuel loads, snag abundance, and snag recruitment in an unmanaged Jeffrey pine mixed conifer forest in Northwestern Mexico. *Forest Ecology and Management*, **199**: 103–113.
- STEPHENS S.L., MOGHADDAS J.J. (2005): Fuel treatment effects on snags and coarse woody debris in a Sierra Nevada mixed conifer forest. *Forest Ecology and Management*, **214**: 53–64.
- STRUKEIJ M., BRAIS S., QUIDEAUA S.A., ANGERS V.A., KEBLI H., DRAPEAU P., OH.S. (2013): Chemical transformations in downed logs and snags of mixed boreal species during decomposition. *Canadian Journal of Forest Research*, **43**: 785–798.
- SZEWCZYK J., SZWAGRZYK J. (1996): Tree regeneration on rotten wood and on soil in old-growth stand. *Vegetatio*, **122**: 37–46.

- TAVANKAR F., BONYAD A.E., MAJNOUNIAN B. (2011): Investigation of damages to stand caused by selection cutting using skidding system in the Asalem-Nav Forest, Iran. *Journal of Environmental Studies*, **37**: 89–98.
- TAVANKAR F., MAJNOUNIAN B., BONYAD A.E. (2013): Felling and skidding damage to residual trees following selection cutting in caspian forests of Iran. *Journal of Forest Science*, **59**: 196–203.
- THOMAS J.W., ANDERSON R.G., MASER C., BULL E.L. (1979): Snags, Wildlife Habitats in Managed Forests: the Blue Mountains of Oregon and Washington. Agricultural Handbook No. 553. Portland, USDA Forest Service: 510
- TREMBLAY J.A., IBARZABAL J., SAVARD J.P.L. (2010): Foraging ecology of black-backed woodpeckers (*Picoides arcticus*) in unburned eastern boreal forest stands. *Canadian Journal of Forest Research*, **40**: 991–999.
- TRIMBLE G.R., MENDEL J.R., KENNEL R.A. (1974): A Procedure for Selection Marking in Hardwoods Combining Silvicultural Consideration with Economic Guidelines. Research Paper NE-292. Newtown Square, USDA Forest Services: 13.
- ULYSHEN M.D. (2011): Arthropod vertical stratification in temperate deciduous forests: Implications for conservation-oriented management. *Forest Ecology and Management*, **261**: 1479–1489.
- WISDOM M.J., BATE L.J. (2008): Snag density varies with intensity of timber harvest and human access. *Forest Ecology and Management*, **255**: 2085–2093.
- WOLF A., MØLLER P.F., BRADSHAW R.H.W., BIGLER J. (2004): Storm damage and long-term mortality in a semi-natural, temperate deciduous forest. *Forest Ecology and Management*, **188**: 197–210.
- YATSKOV M., HARMON M.E., KRANKINA O.N. (2003): A chronosequence of wood decomposition in the boreal forests of Russia. *Canadian Journal of Forest Research*, **33**: 1211–1226.

Received for publication July 7, 2014

Accepted after corrections September 23, 2014

---

*Corresponding author:*

Prof. ANGELA LO MONACO, University of Tuscia, Department of Agriculture, Forests, Nature and Energy,  
Via San Camillo de Lellis, 01100 Viterbo, Italy; e-mail: lomonaco@unitus.it

---