

## Changes in the threat of spruce stand disintegration in the Beskid Śląski and Żywiecki Mts. in the years 2007–2010

S. MAŁEK<sup>1</sup>, J. BARSZCZ<sup>1</sup>, K. MAJSTERKIEWICZ<sup>2</sup>

<sup>1</sup>*Department of Forest Ecology, Faculty of Forestry, University of Agriculture in Krakow, Krakow, Poland*

<sup>2</sup>*Regional Directorate of the State Forests in Katowice, Katowice, Poland*

**ABSTRACT:** The present paper describes the dynamics of changes in spruce stand disintegration in the Beskid Śląski and Żywiecki Mts. on the basis of the results of inventory performed in 2010 as compared to those in 2009. The research was done in six forest districts in the Beskid Mts. The area of spruce stands in degrees of threat of disintegration on the day of 30<sup>th</sup> September in the research period was determined for forest districts Bielsko, Ustroń, Wisła, Węgierska Górka, Jeleśnia and Ujsoły, on the basis of data in the State Forests Electronic Database (SILP) in degrees of threat of stand disintegration as well as in climate and vegetation zones. A comparison of the results of inventory of the stands threatened with disintegration in six forest districts in 2007 (altogether 24,446 ha) and in 2010 (altogether 16,886 ha) indicates a decrease in the rate of spruce stand disintegration. This may point to the slowing down of the rate of spruce stand disintegration as well as to considerable intensification of reconstruction work.

**Keywords:** spruce stands; degrees of reconstruction urgency; inventory; climate and vegetation zones; Beskid Śląski and Żywiecki Mts.

The spruce stands of the Beskid Śląski and Beskid Żywiecki Mts. are affected by a forest system disease. The problem concerns spruce in its whole European range (ŠRÁMEK et al. 2010). What contributes to the disintegration of the stands of this species are unfavourable climatic changes, anthropogenic factors and historical reasons related to the establishment of spruce monocultures, usually of foreign origin, without regard to site conditions (BRODA 1965; MACH, BARSZCZ 1983; SZUJECKI 1992; SIEROTA 1995; BARSZCZ, MAŁEK 2003; GRODZKI 2007; BARKA et al. 2010; HLÁSNY et al. 2010a, b; ŠRÁMEK et al. 2010). The need to reconstruct these spruce stands was indicated already in the past (MACH, BARSZCZ 1983). No one expected that their disease would assume such a rapid and

mass character. During the last few years there has occurred rapid acceleration of tree dieback and stand disintegration, which leads to the degradation of the entire forest environment, threat to forest biocoenosis stability, lowering of the production potential and changes in forest management. The process is further intensified by violent, extreme weather phenomena, such as windstorms, winters that are either long and severe or too mild, droughts, etc. These phenomena have been making the condition of these forests increasingly worse in terms of both area and quality. In the last two years there has occurred a distinct decrease in the area of the spruce stands threatened by disintegration, which may point to a decline of this phenomenon, due to such factors as a fall in the population of

---

Supported by NFOŚiGW in Warsaw as a part of the project Improvement of the Reconstruction Methods of Spruce Stand Disintegration in the Beskid Śląski and Żywiecki Mts. based on the close to nature silviculture and increased biological biodiversity – part I.

the bark beetle as well as a fall in the area of dying spruce stands, which are gradually removed.

The aim of the present study was to assess changes in the degree of threat of spruce stand disintegration in the Beskid Śląski and Żywiecki Mts. in the years 2009–2010 on the basis of the instructions, developed and improved by the present authors, of data acquisition from the State Forests Electronic Database (SILP). The present study is a response to an urgent need in forest practice for fast and objective periodical assessment of the dynamically changing area of disintegrating spruce stands in the analysed regions.

## MATERIAL AND METHODS

The research was done in six forest districts in the Beskidy Mts., belonging to the Regional Directorate of the State Forests in Katowice. The area of spruce stands in degrees of disintegration threat on the day of 30<sup>th</sup> September 2010 was determined by K. Majsterkiewicz for all the forest districts under assessment (Bielsko, Ustroń, Wisła, Węgierska Górka, Jeleśnia and Ujsoły). Due to a limited possibility of data acquisition from the State Forests Electronic Database (SILP), the following data contained in this database were considered: the year of developing the forest management plan; forest management year; the index of increase of sanitary deadwood (NPS) and of increase of active deadwood (NPC); stand density in particular forest layers; the degree of area being covered by the regeneration layer, with a division into classes (0.1–0.3; 0.4–0.5; > 0.5) for the year of the forest management plan; the record of recognized natural regeneration and underplanting in a given year; altitude: minimum, maximum and medium for a given division; blanks and irregularly stocked open stands; for spruce which occurs in the first forest layer: initial stock (for the year of developing the forest management plan) and stock for the current year.

As the above-mentioned forest districts do not have the area division corresponding to climate and vegetation zones yet, the stand divisions were included on the basis of medium altitude into one of the four altitude zones: foothills, lower forest zone, middle forest zone and upper forest zone, assuming that in the Beskid Śląski and Mały Mts. the upper borders of the first three zones will be roughly delimited by the following contour lines: 500, 800 and 1,000 m a.s.l., while in the Beskid Żywiecki Mts. they will be: 600, 900 and 1,100 m a.s.l., respec-

tively. The Beskid Śląski Mts. include the following forest districts: Ustroń, Bielsko and Wisła as well as the whole division of Lipowa belonging to the forest district of Węgierska Górka and the forest sub-districts of Kamesznica, Sikorzany and Zielona from the division of Węgierska Górka. The Beskid Żywiecki Mts. include the remaining forest sub-districts of the division of Węgierska Górka and the forest districts: Jeleśnia and Ujsoły.

In the form of the algorithm applied here the researchers used the joined area of regeneration (natural and planted) because of the lack of precise data on its origin in the SILP database. In-field recognition of the possibility to obtain data from SILP according to the above criteria allowed for adopting the following rules of qualification of stands for reconstruction in accordance with degrees of its urgency:

- (1) very urgent reconstruction within a very short period (up to a few years – max. 10 years), which should include the following stands: dying ones, where the harvesting of sanitary deadwood (NPS) is above 50 m<sup>3</sup>·ha<sup>-1</sup>, which corresponds to the active deadwood increase index (NPC) over 16, tree coverage in the stand of 0.1–0.3 and degree of coverage with a young generation of 0.1–0.3;
- (2) urgent reconstruction within a relatively short period (up to several years – 10 to max. 20 years), which should include the following stands: weakened ones, where the harvesting of sanitary deadwood (NPS) is 30–50 m<sup>3</sup>·ha<sup>-1</sup>, which corresponds to the active deadwood increase index (NPC) of 10–16, stand tree coverage of 0.4–0.5 and degree of coverage with a young generation of 0.4–0.5;
- (3) relatively urgent reconstruction within a longer period (within 20–30 years), which should include the following stands: weakened ones, where the harvesting of sanitary deadwood (NPS) is below 30 m<sup>3</sup>·ha<sup>-1</sup>, which corresponds to the active deadwood increase index (NPC) below 10, stand tree coverage of 0.5–0.7 and degree of coverage with a young generation above 0.5;

As in their previous research (BARSZCZ, MAŁEK 2005, 2007, 2009, 2010; BARSZCZ et al. 2009), the present authors recommend introducing the 4<sup>th</sup> degree of stand reconstruction urgency in order to include the remaining spruce stands, mostly cultures, with tree coverage above 0.7, as well as a few spruce stands which should undergo reconstruction within a long period (over 30 years).

The data obtained from SILP allowed for an attempt at visualization of the results of the current inventory in the form of digital maps showing dif-

ferent degrees of stand reconstruction urgency in accordance with the divisions within each of the six forest districts under analysis (Figs. 1 and 2).

## RESULTS AND DISCUSSION

The inventory of the stands threatened with disintegration (lacking sufficient regeneration) in degrees of their reconstruction urgency 1–3, performed in 2005, showed that in the six forest districts under analysis the total area of these stands amounted to 14,609 ha (BARSZCZ, MAŁEK 2005), in 2007 it was already 24,446 ha (BARSZCZ, MAŁEK 2007; BARSZCZ et al. 2009), whereas in 2009 their area decreased to 20,557 ha, i.e. by about 14% (BARSZCZ, MAŁEK 2009), and in 2010 to 16,886 ha (BARSZCZ, MAŁEK 2010). The decrease in the area of the threatened stands in the last period under analysis amounts to 3,671 ha, i.e. about 19.0%, which indicates a slowing down of the rate of spruce stand disintegration in the research area (Table 1). It must be remembered, however, that this slower rate is also caused by a decrease in the area of the threatened stands due to their removal during sanitary cutting (which was the most intensive in Węgierska Górką Forest District) as well

as a lower rate of tree dying caused by a fall in the gradation of the spruce bark beetle (GRODZKI 2007).

The area that is the most threatened in this respect is Ujsoły Forest District and, to a smaller degree, Jeleśnia and Wisła Forest Districts. Against this background, the forest districts of Bielsko and Ustroń face a relatively lower threat. This indicates constant dieback of spruce stands in the southern and south-eastern part of the research area (Table 1, Figs. 1 and 2), which is where it began the latest.

In 2010, with regard to altitude, the largest area of stands threatened with disintegration in the six forest districts (including Jeleśnia Forest District) occurred in the lower forest zone (Tables 2 and 3), particularly in the following forest districts: Ujsoły (about 6,127 ha), Wisła (about 3,490 ha) and Ustroń (about 2,231 ha) – cf. Table 2.

Reconstruction of stands in the lower forest zone and those in foothills should not pose any large problems thanks to good site conditions, possibility to use numerous tree species and a considerable area of stands producing seeds of the main forest-creating species. The next group (in terms of area) consists of threatened spruce stands in the middle forest zone, especially in Ujsoły Forest District (approx. 2,945 ha), and next in Jeleśnia (approx. 1,345 ha) – cf. Table 2.

Table 1. The area (in ha) of stands threatened with disintegration, with regard to natural and artificial regeneration, in degrees of their reconstruction urgency in the mountain areas of forest districts of the Regional Directorate of the State Forests in Katowice in 2009–2010

| Type of area and degree of reconstruction urgency |                 | Forest districts |        |       |                 |          |        | Total  |        |
|---|-----------------|------------------|--------|-------|-----------------|----------|--------|--------|--------|
|   |                 | Bielsko          | Ustroń | Wisła | Węgierska Górką | Jeleśnia | Ujsoły |        |        |
| Forest area in forest district                    | 2009            | 9.966            | 11.226 | 8.506 | 9.107           | 11.539   | 13.225 | 63.569 |        |
|   | 2010            | 9.964            | 11.219 | 8.493 | 9.091           | 11.539   | 13.225 | 63.531 |        |
| Area of stands threatened with disintegration     | 2009            | 2.114            | 3.360  | 4.489 | 4.222           | 3.993    | 9.341  | 2.7519 |        |
|   | 2010            | 1.767            | 3.279  | 3.950 | 3.739           | 3.730    | 9.350  | 2.2785 |        |
| area with regeneration                            | 2009            | 331              | 1.135  | 834   | 1.237           | 838      | 2.586  | 6.961  |        |
|   | 2010            | 184              | 1.059  | 1.119 | 233             | 470      | 2.832  | 5.897  |        |
| Including<br>for reconstruction                   | degree 1        | 2009             | 491    | 422   | 456             | 859      | 181    | 296    | 2.705  |
|   |                 | 2010             | 349    | 364   | 247             | 149      | 199    | 406    | 1.713  |
|   | degree 2        | 2009             | 283    | 287   | 672             | 759      | 223    | 505    | 2.729  |
|   |                 | 2010             | 217    | 193   | 400             | 228      | 240    | 317    | 1.595  |
|   | degree 3        | 2009             | 1.009  | 1.515 | 2.527           | 1.367    | 2.751  | 5.953  | 15.122 |
|   |                 | 2010             | 1.017  | 1.632 | 2.184           | 129      | 2.821  | 5.755  | 13.538 |
|   | all degrees 1–3 | 2009             | 1.783  | 2.224 | 3.655           | 2.985    | 3.155  | 6.755  | 20.557 |
|   |                 | 2010             | 1.583  | 2.188 | 2.831           | 506      | 3.260  | 6.518  | 16.886 |

degrees of reconstruction urgency: 1 – very urgent, 2 – urgent, 3 – relatively urgent




Fig. 1. The distribution of spruce stands for reconstruction (in degrees of reconstruction urgency: 1 – very urgent, 2 – urgent, 3 – relatively urgent, 4 – other spruce stands) in the forest districts: (a) Bielsko, (b) Ustroń and (c) Wisła, according to data obtained from SILP (on 30.08.2010)




Fig. 2. The distribution of spruce stands for reconstruction (in degrees of reconstruction urgency: 1 – very urgent, 2 – urgent, 3 – relatively urgent, 4 – other spruce stands) in the forest districts: (a) Węgierska Góra, (b) Jeleśnia and (c) Ujszoły, according to data obtained from SILP (on 30.08.2010)

Table 2. The area of divisions threatened with disintegration (in ha) divided into areas with regeneration and those for reconstruction in climate and vegetation zones in the mountain stands of forest districts of the Regional Directorate of the State Forests in Katowice in 2009–2010

| Climate and vegetation zones          |      | Area of divisions threatened with disintegration (ha) |              |                    |
|---------------------------------------|------|---|--------------|--------------------|
|                                       |      | total   | regeneration | for reconstruction |
| <b>Bielsko Forest District</b>        |      |   |              |                    |
| Foothills                             | 2009 | 268   | 1            | 267                |
|                                       | 2010 | 417   | 5            | 412                |
| Lower forest zone                     | 2009 | 1134  | 177          | 957                |
|                                       | 2010 | 760   | 42           | 718                |
| Middle forest zone                    | 2009 | 520   | 77           | 443                |
|                                       | 2010 | 395   | 54           | 341                |
| Upper forest zone                     | 2009 | 192   | 76           | 116                |
|                                       | 2010 | 195   | 83           | 112                |
| <b>Ustroń Forest District</b>         |      |   |              |                    |
| Foothills                             | 2009 | 932   | 234          | 698                |
|                                       | 2010 | 1.030   | 181          | 849                |
| Lower forest zone                     | 2009 | 2.231   | 781          | 1.450              |
|                                       | 2010 | 1.934   | 756          | 1.178              |
| Middle forest zone                    | 2009 | 197   | 120          | 76                 |
|                                       | 2010 | 285   | 122          | 161                |
| Upper forest zone                     | 2009 | 0   | 0            | 0                  |
|                                       | 2010 | 0   | 0            | 0                  |
| <b>Wisła Forest District</b>          |      |   |              |                    |
| Foothills                             | 2009 | 0   | 0            | 0                  |
|                                       | 2010 | 56  | 0            | 56                 |
| Lower forest zone                     | 2009 | 3.490   | 613          | 2.877              |
|                                       | 2010 | 3.006   | 858          | 2.148              |
| Middle forest zone                    | 2009 | 792   | 202          | 590                |
|                                       | 2010 | 677   | 187          | 490                |
| Upper forest zone                     | 2009 | 207   | 19           | 188                |
|                                       | 2010 | 211   | 74           | 137                |
| <b>Węgerska Górka Forest District</b> |      |   |              |                    |
| Foothills                             | 2009 | 95  | 18           | 77                 |
|                                       | 2010 | 41  | 0            | 41                 |
| Lower forest zone                     | 2009 | 2.892   | 842          | 2.050              |
|                                       | 2010 | 275   | 57           | 218                |
| Middle forest zone                    | 2009 | 1.156   | 345          | 811                |
|                                       | 2010 | 220   | 87           | 133                |
| Upper forest zone                     | 2009 | 79  | 32           | 47                 |
|                                       | 2010 | 203   | 89           | 114                |

Table 2 to be continued

| Climate and vegetation zones     | Area of divisions threatened with disintegration (ha) |              |                    |       |
|----------------------------------|---|--------------|--------------------|-------|
|                                  | total   | regeneration | for reconstruction |       |
| <b>Jeleśnia Forest District*</b> |   |              |                    |       |
| Foothills                        | 277   | 51           | 226                |       |
| Lower forest zone                | 1.789   | 213          | 1.576              |       |
| Middle forest zone               | 1.345   | 172          | 1.173              |       |
| Upper forest zone                | 319   | 34           | 285                |       |
| <b>Ujsoły Forest District</b>    |   |              |                    |       |
| Foothills                        | 2009  | 234          | 50                 | 184   |
|                                  | 2010  | 175          | 18                 | 157   |
| Lower forest zone                | 2009  | 6.127        | 1.852              | 4.275 |
|                                  | 2010  | 6.163        | 1.895              | 4.268 |
| Middle forest zone               | 2009  | 2.945        | 684                | 2.261 |
|                                  | 2010  | 2.912        | 857                | 2.055 |
| Upper forest zone                | 2009  | 35           | 0                  | 35    |
|                                  | 2010  | 100          | 62                 | 38    |

\*for Jeleśnia Forest District in 2009 there were no data in the SILP-LAS base that would make it possible to ascribe divisions to climate and vegetation zones

The total area of threatened stands in this zone amounts to about 6,000 ha. In the middle forest zone, the problem (apart from a large area of stands for reconstruction) is a smaller area of spruce seed-bearing stands, a small area of such beech stands and a lack of such a base for fir and admixture species. Threatened stands in the upper forest zone (altogether about 1,000 ha, mostly in Jeleśnia Forest District) are going to, despite a relatively small area (Table 2), cause the largest reconstruction problems due to a lack of recognized seed-bearing stands, a small number of species and very difficult site conditions, especially at mountain tops and ridges and because of the need for different silvicultural measures in such areas.

The above data (only for five forest districts, except Jeleśnia) show that between 2009 and 2010 there occurred a decrease in the area of threatened spruce stands in the lower and middle forest zones, a distinct increase in the foothills and a slight increase in the upper forest zone (Table 2). A comparison between the areas of the recognized natural and artificial regeneration in regions threatened with disintegration shows that in the years 2009–2010 there were about 5,900 ha that were considered as reconstructed (Table 1). With regard to altitude, the largest number of areas included in this group occurs in

the lower and middle forest zones, particularly in the forest districts of Ujsoły, Wisła and Ustroń.

A comparison of the shares of areas of particular degrees of reconstruction urgency in the total area of stands threatened with disintegration shows a decrease, which is the largest in degree 1 – cf. Table 1. Despite these changes, relative proportions between the area for reconstruction in particular climate and vegetation zones and the total area in the whole region under analysis are similar to those in previous record periods (Fig. 1, Table 3).

At high altitudes, mountain areas will have to undergo artificial regeneration and in certain cases also measures stimulating natural regeneration (BARSZCZ, MAŁEK 2003, 2005).

## CONCLUSIONS

Comparison of the results of inventory (from the year 2007: 24,446 ha, 2009: 20,557 ha and 2010: 16,886 ha) of stands threatened with disintegration in six forest districts of the Beskidy Mts: Bielsko, Ustroń, Wisła, Węgierska Górką, Jeleśnia and Ujsoły shows a decrease in the threat of spruce stand disintegration. In total, the decrease in such stands in the period 2009–2010 amounted to 3,671 ha (i.e. nearly

Table 3. The area of fragments of divisions for reconstruction (in ha) according to degrees of its urgency in climate and vegetation zones in the mountain stands of forest districts of the Regional Directorate of the State Forests in Katowice in 2009–2010

| Climate and vegetation zones          |      | Degree of reconstruction urgency |     |       | Total |
|---------------------------------------|------|----------------------------------|-----|-------|-------|
|                                       |      | 1                                | 2   | 3     |       |
| <b>Bielsko Forest District</b>        |      |                                  |     |       |       |
| Foothills                             | 2009 | 111                              | 55  | 101   | 267   |
|                                       | 2010 | 176                              | 73  | 163   | 412   |
| Lower forest zone                     | 2009 | 156                              | 113 | 688   | 957   |
|                                       | 2010 | 47                               | 67  | 604   | 718   |
| Middle forest zone                    | 2009 | 165                              | 103 | 175   | 443   |
|                                       | 2010 | 92                               | 51  | 198   | 341   |
| Upper forest zone                     | 2009 | 59                               | 12  | 45    | 116   |
|                                       | 2010 | 34                               | 26  | 52    | 112   |
| <b>Ustroń Forest District</b>         |      |                                  |     |       |       |
| Foothills                             | 2009 | 256                              | 53  | 389   | 698   |
|                                       | 2010 | 281                              | 80  | 488   | 849   |
| Lower forest zone                     | 2009 | 165                              | 214 | 1.071 | 1.450 |
|                                       | 2010 | 60                               | 85  | 1.033 | 1.178 |
| Middle forest zone                    | 2009 | 1                                | 20  | 55    | 76    |
|                                       | 2010 | 22                               | 28  | 111   | 161   |
| Upper forest zone                     |      |                                  |     |       |       |
| <b>Wisła Forest District</b>          |      |                                  |     |       |       |
| Foothills                             | 2009 | 0                                | 0   | 0     | 0     |
|                                       | 2010 | 11                               | 0   | 45    | 56    |
| Lower forest zone                     | 2009 | 331                              | 506 | 2.040 | 2.877 |
|                                       | 2010 | 171                              | 258 | 1.719 | 2.148 |
| Middle forest zone                    | 2009 | 53                               | 136 | 401   | 590   |
|                                       | 2010 | 45                               | 89  | 356   | 490   |
| Upper forest zone                     | 2009 | 72                               | 30  | 86    | 188   |
|                                       | 2010 | 20                               | 53  | 64    | 137   |
| <b>Węgerska Górka Forest District</b> |      |                                  |     |       |       |
| Foothills                             | 2009 | 21                               | 6   | 50    | 77    |
|                                       | 2010 | 41                               | 0   | 0     | 41    |
| Lower forest zone                     | 2009 | 448                              | 523 | 1.079 | 2.050 |
|                                       | 2010 | 34                               | 119 | 65    | 218   |
| Middle forest zone                    | 2009 | 381                              | 211 | 219   | 811   |
|                                       | 2010 | 22                               | 64  | 47    | 133   |
| Upper forest zone                     | 2009 | 9                                | 19  | 19    | 47    |
|                                       | 2010 | 52                               | 45  | 17    | 114   |



Table 3 to be continued

| Climate and vegetation zones     | Degree of reconstruction urgency |     |       | Total |       |
|----------------------------------|----------------------------------|-----|-------|-------|-------|
|                                  | 1                                | 2   | 3     |       |       |
| <b>Jeleśnia Forest District*</b> |                                  |     |       |       |       |
| foothills                        | 93                               | 25  | 108   | 226   |       |
| lower forest zone                | 83                               | 170 | 1323  | 1.576 |       |
| middle forest zone               | 23                               | 45  | 1.105 | 1.173 |       |
| upper forest zone                | 0                                | 0   | 285   | 285   |       |
| <b>Ujsoły Forest District</b>    |                                  |     |       |       |       |
| Foothills                        | 2009                             | 7   | 1     | 176   | 184   |
|                                  | 2010                             | 38  | 16    | 103   | 157   |
| Lower forest zone                | 2009                             | 204 | 429   | 3.641 | 4.275 |
|                                  | 2010                             | 336 | 238   | 3.694 | 4.268 |
| Middle forest zone               | 2009                             | 85  | 75    | 2.101 | 2.261 |
|                                  | 2010                             | 29  | 63    | 1.963 | 2.055 |
| Upper forest zone                | 2009                             | 0   | 0     | 35    | 35    |
|                                  | 2010                             | 3   | 0     | 35    | 38    |

\*for Jeleśnia Forest District in 2009 there were no data in the SILP-LAS base that would make it possible to ascribe divisions to climate and vegetation zones; 1 – very urgent, 2 – urgent, 3 – relatively urgent

20%). This indicates a slowing down of the rate of spruce stand disintegration, and increase in the area of stands with recognized regeneration, but also a decrease in the total area of spruce stands due to their removal during sanitary cutting (especially in Węgierska Górka Forest District).

At present, spruce stands which are the most threatened with disintegration are found in Ujsoły Forest District, and to smaller degrees in the forest districts of Jeleśnia and Wisła. It indicates a constant rate of dieback of stands in the same, southern and south-eastern part of the analysed region. Moreover, the dieback has extended considerably in the foothills and slightly in the upper forest zone, whereas it decreased in the lower and middle forest zones.

The reconstruction of stands in the lower forest zone and those in the foothills should not pose any large problems except for their size, thanks to good site conditions, the possibility to use a large number of tree species and a considerable area of the existing seed-bearing stands of the main forest-creating species.

The second group with regard to area are threatened spruce stands in the middle forest zone, mainly in Ujsoły Forest District, and then in Jeleśnia. In this zone, the problem is due to a smaller area of seed-

bearing spruce stands, only a small one for beech and practically none for fir and admixture species.

Threatened stands in the upper forest zone (most of which are in Jeleśnia) are going to cause the greatest problems with their reconstruction because of a lack of recognized seed-bearing stands, a small number of species and very difficult site conditions, especially in top and ridge mountain locations and because of the necessity to change silvicultural measures in these areas.

Reconstruction or conversion should also be applied in the remaining, artificial spruce stands in the mountains, with the relatively least visible signs of damage. The above-mentioned measures should be conducted in areas of all degrees of reconstruction urgency, with the intensity suitable for the adopted reconstruction periods.

## References

- BARKA I., BUCHA T., SITKOVÁ Z., HLÁSNY T., KULLA L. (2010): Remote sensing view on declining spruce forest. In: HLÁSNY T., SITKOVÁ Z. (eds): Spruce Forest Decline in the Beskids. Zvolen, National Forest Centre – Forest Research Institute Zvolen and Czech University of Life

- Sciences Prague and Forestry and Game Management Research Institute Jiloviště – Strnady: 33–47.
- BARSZCZ J., MAŁEK S. (2003): Perspektywy wzrostu świerka w wyższych położeniach Beskidu Śląskiego na obszarach zagrożenia trwałości lasu w świetle oceny jego odnowień. [Prospects of the growth of spruce at higher altitudes of the Beskid Śląski Mts in areas of threat to forest sustainability in the light of evaluation of its regeneration.] In: Drzewostany świerkowe: stan, problemy, perspektywy rozwojowe. [Spruce stands: status, problems and development perspectives.] PTL, Jaszowiec: 141–159.
- BARSZCZ J., MAŁEK S. (2005): Opracowanie wytycznych hodowlanych do przebudowy zagrożonych drzewostanów świerkowych o różnym stopniu zagrożenia rozpadem i ustalenie dla nich celów hodowlanych uwzględniających skład gatunkowy, pochodzenie, możliwości wykorzystania odnowienia naturalnego, osłony i zabiegów melioracyjnych (w tym fitomelioracyjnych) i nawożeniowych. [Development of management guidelines for the redevelopment of endangered spruce stands with different degrees of the threat of disintegration and determination of their management objectives taking into account their composition, origin, the possibility of using natural regeneration, protection as well as drainage and fertilization measures (including phytomelioration).] In: BARSZCZ J., MAŁEK S. (eds): Doskonalenie rewitalizacji siedlisk i przebudowy drzewostanów górskich w RDLP Katowice z uwzględnieniem poprawy stosunków wodnych i selekcji genetycznej drzew – etap III. [Improving habitat revitalization and reconstruction of the mountain stands in the Regional Directorate of the State Forests in Katowice including improvement of water relations and genetic selection of trees – stage III.] Kraków, Department of Forest Ecology: 49–71.
- BARSZCZ J., MAŁEK S. (2007): Dynamika zmian zagrożenia rozpadem drzewostanów świerkowych w Beskidzie Śląskim i Żywieckim [The dynamics of changes in the threat of spruce stand disintegration in the Beskid Śląski and Żywiecki Mts.] In: BARSZCZ J., MAŁEK S. (eds): Doskonalenie rewitalizacji siedlisk i przebudowy drzewostanów górskich w RDLP Katowice z uwzględnieniem selekcji genetycznej drzew – kontynuacja badań i doświadczeń oraz wdrażanie zaleceń gospodarczych – etap I. [Improving habitat revitalization and reconstruction of the mountain stands in the Regional Directorate of the State Forests in Katowice including genetic selection of trees – continuation of research and experiments and implementation of management guidelines – stage I.] Kraków, Department of Forest Ecology: 5–23.
- BARSZCZ J., MAŁEK S. (2009): Doskonalenie rewitalizacji siedlisk i przebudowy drzewostanów górskich w RDLP Katowice – podsumowanie wyników badań i zalecenia gospodarcze w zakresie zadania 1. [Improving habitat revitalization and reconstruction of the mountain stands in the Regional Directorate of the State Forests in Katowice – research conclusions and management recommendations within task 1.] In: BARSZCZ J., MAŁEK S. (eds): Doskonalenie rewitalizacji siedlisk i przebudowy drzewostanów górskich w RDLP Katowice z uwzględnieniem selekcji genetycznej drzew – kontynuacja badań i doświadczeń oraz wdrażanie zaleceń gospodarczych – etap III. [Improving habitat revitalization and reconstruction of the mountain stands in the Regional Directorate of the State Forests in Katowice including genetic selection of trees – continuation of research and experiments and implementation of management guidelines – stage III.] Kraków, Department of Forest Ecology: 47–76.
- BARSZCZ J., MAŁEK S. (2010): Doskonalenie metody oceny stopnia zagrożenia rozpadem świerczyn beskidzkich w oparciu o dane z SILP i ocenę ich stanu w 2010 roku. [Improvement of the methods of reconstruction of threatened spruce stands in the Beskid Śląski and Żywiecki Mts. on the basis of data from the State Forests Electronic Database and assessment of their condition in 2010] In: BARSZCZ J., MAŁEK S. (eds): Określenie budowy, struktury i dynamiki lasów naturalnych z Beskidu Śląskiego i Żywieckiego jako podstawa przebudowy świerczyn beskidzkich – Doskonalenie metod przebudowy zagrożonych świerczyn w Beskidach: Śląskim i Żywieckim w oparciu o hodowlę lasu bliską naturze i wzmoczoną różnorodność biologiczną – etap I. [Determination of the composition, structure and dynamics of natural forests in the Beskid Śląski and Żywiecki Mts. as a basis for reconstruction of the Beskid spruce stands – Improvement of the methods of reconstruction of threatened spruce stands in the Beskid Śląski and Żywiecki Mts. on the basis of close to nature forest management and intensified biodiversity – stage I.] Kraków, Department of Forest Ecology: 64–74
- BARSZCZ J., MAŁEK S., MAJSTERKIEWICZ K. (2009): Dynamika zmian zagrożenia rozpadem świerczyn Beskidu Śląskiego i Żywieckiego. [The dynamics of changes in the threat of disintegration of spruce stands in the Beskid Śląski and Żywiecki Mts.] Prace Komisji Nauk Rolniczych, Leśnych i Weterynaryjnych PAU, **11**: 93–113.
- BRODA J. (1965): Puszcze Karpackie i Sudeckie. [Carpathian and Sudeten forests.] In: Dzieje lasów, leśnictwa i drzewnictwa w Polsce. [The History of Forests, Forestry and Wood Science in Poland.] Warszawa, PWRiL: 685–715.
- GRODZKI W. (2007): Spatio-temporal patterns of the Norway spruce decline in the Western Beskidy mountains in Poland. *Journal of Forest Science*, **53** (Special Issue): 38–44.
- HLÁSNY T., KULLA L., BARKA L., TURČÁNI M., SITKOVÁ Z., KOREŇ M. (2010a): The proposal of biotic hazard zones in selected spruce dominant Regions in Slovakia. *Journal of Forest Science*, **56**: 236–242
- HLÁSNY T., GRODZKI W., ŠRÁMEK V., HOLUŠA J., KULLA L., SITKOVÁ Z., TURČÁNI M., RĄCZKA G., STRZELIŃSKI P., WĘGIEL A. (2010b): Spruce forest decline in the Beskids. In: HLÁSNY T., SITKOVÁ Z. (eds): Spruce Forest Decline in the Beskids. Zvolen, National Forest Centre – Forest

- Research Institute Zvolen and Czech University of Life Sciences Prague and Forestry and Game Management Research Institute Jíloviště – Strnady: 33–47.
- MACH J., BARSZCZ J. (1983): Ekologiczne uwarunkowania produktywności lasów Beskidu Śląskiego. [Ecological determinants of the productivity of the Beskid Śląski forests.] In: Zagospodarowanie lasów górskich na przykładzie Beskidu Śląskiego. Jaszowiec 24.–25. September 1983. Warszawa, PTL: 14–42.
- SIEROTA Z. (1995): Możliwości zmniejszenia predyspozycji chorobowej lasów metodami gospodarki leśnej. [The possibilities of reducing the disease predisposition of forests by means of forest management methods in Poland.] Prace IBL, **22**: 1–55.
- ŠRÁMEK V., SITKOVÁ Z., MAŁEK S., PAVLENDÁ P., HLÁSNÝ T. (2010): Air pollution and forest nutrition – what are their roles in spruce forest decline? In: HLÁSNÝ T., SITKOVÁ Z. (eds): Spruce forest decline in the Beskids. Zvolen, National Forest Centre – Forest Research Institute Zvolen and Czech University of Life Sciences Prague and Forestry and Game Management Research Institute Jíloviště – Strnady: 33–47.
- SZUJECKI A. (1992): Czy lasy muszą zginąć? [Do forests have to die?] Warszawa, Wiedza Powszechna: 126.

Received for publication November 2, 2011

Accepted after corrections November 6, 2012

---

*Corresponding author:*

STANISŁAW MAŁEK, PhD, University of Agriculture in Krakow, Faculty of Forestry, Department of Forest Ecology, Aleja 29 Listopada 46, 31-425 Krakow, Poland  
e-mail: rlmalek@cyf-kr.edu.pl

---