

# Leaf growth under temperature and light control

J. Repková, M. Brestič, K. Olšovská

*Faculty of Agrobiological and Food Resources, Slovak Agricultural University in Nitra, Nitra, Slovak Republic*

## ABSTRACT

Dynamics of crop growth and photosynthesis are two main processes that are of major importance for adaptation of plants to their environment. Two experiments were carried out during 2005 and 2006 with sun and shaded barley plants. The results showed that leaf area increased with leaf position on the main stem up to leaf position 5 for sun and leaf position 6 for shaded plants, and then declined towards flag leaves. Air temperature affected leaf appearance, mainly at the beginning of the growing season. A positive correlation between leaf expansion duration (LED) and accumulated temperature sum was measured for both variants, LED linearly increased with temperature sum. Leaf expansion rate (LER) showed a similar dependence on accumulated air temperatures in both light variants. A strong relation was found between soil temperature and LER for the first four leaves of the main stem of shaded plants in 2006. In shaded environment the higher LER was associated with lower accumulated irradiance sum when light restriction supported an increase of leaf elongation and final leaf length.

**Keywords:** barley (*Hordeum vulgare* L.); leaf growth; leaf area; leaf expansion; temperature; irradiance; sun and shaded leaves

Leaf area is the main determining factor affecting light interception by crop and biomass production. Therefore, any reduction of leaf expansion rate is usually associated with reduction of photosynthesis and consequent decrease in above-ground biomass, grain yield and quality (Schurr et al. 2006, Váňová et al. 2006). In cereals, especially barley and wheat, leaf blade area is just a part of total assimilatory area. Wang et al. (2001) mentioned that non-leaf green organs consisting of ear, peduncle and flag leaf sheath have a greater importance for the grain filling of wheat than flag leaf and penultimate blades. In the field, spring cereal crops are characterized by occurrence of spacious leaf area produced in short time interval which limits considerably reaching the high proportion of absorbed light needed for driving crop productivity (Tardieu et al. 2005). More detailed study of environmental effects on leaf emergence is necessary for understanding canopy growth dynamics in fluctuated environmental conditions.

Spring barley crop development is a succession of phenological events regulated by relationship between genotype and environment. Rate of leaf appearance is a developmental trait which, together with final number of differentiated leaves, determines the length of crop cycle (Abeledo et al. 2004). Leaf area growth and thus ontogenetic and leaf area insertion changes are strongly modified by temperature, radiation and shading as well as photoperiod, air humidity, water supply and nitrogen nutrition (Tardieu et al. 2005, Schurr et al. 2006).

The rate of leaf formation on the stems depends primarily on the temperature effect on leaf expansion, especially at the zones of cell expansion. Tamaki et al. (2002) and Bartholomew and Williams (2005) showed that emergence of new leaves is a linear function of time at any temperature regime. When temperature increases, time interval of two successive leaf tips appearance is dropped (Kirby 1995, Bos and Neuteboom 1998, McMaster et al. 2003). Leaf growth rate raises with increasing tem-

---

Supported by the Scientific Grant Agency (VEGA) of the Ministry of Education of the Slovak Republic and the Slovak Academy of Sciences, Project No. 1/0803/08, and by the Slovak Research and Development Agency, Project No. LPP-0345-2006.

perature until the optimum temperature is reached, but reversely further raise of temperature could reduce leaf growth (Cao and Moss 1989, Tamaki et al. 2002). Some previous studies (Hay and Wilson 1982, McMaster et al. 2003) demonstrated that root zone temperature affects leaf growth, leaf development and physiological processes, such as carbon assimilation and transpiration of higher plants. Therefore, leaf development rate is more closely associated with soil temperature near soil surface than air temperature during early stages of crop development. As Hay and Wilson (1982) observed for wheat leaf appearance, soil temperature at the depth of 5 cm was more effective than air temperature.

Light environment can also modify leaf growth and tiller appearance in cereal crops, mainly if canopy plants compete for light (Abeledo et al. 2004). Self-shading of plants within a dense canopy is associated with an increase of leaf length and reduction of tillering. These effects are linked to altered red: far red ratio light quality due to shading at canopy basis. As a consequence of compensation for lower light interception the increase of leaf area could occur, through the increased leaf elongation rate and leaf expansion duration (Bahmani et al. 2000, Evers et al. 2006). In some cases leaf shading can increase the length of elongation zone and leaf elongation rate, but Bos and Neuteboom (1998) did not find a significant effect of light intensity on leaf elongation rate. Miralles and Richards (2000) observed that increasing light intensity is usually associated with an increase of time interval between appearance of two consecutive leaves.

In cereal crops, leaf blade expansion takes places rather via an increase in length than in width since their emergence from sheath bundle (Bos and Neuteboom 1998). For that reason the analysis

of area expansion of individual leaves requires an assessment of such parameters as leaf elongation rate (LER), leaf elongation duration (LED) and maximum leaf width. Effects of temperature on leaf appearance rates and growth are usually quantified as some form of thermal time calculated from air temperature above the crop canopy (McMaster et al. 2003). Since temperature varies considerably under field conditions, the aim of this work was to analyze effects of fluctuating air and soil temperatures on barley leaves growth under two contrasting light environments, full irradiation and simulated shade, applying the parameters mentioned.

## MATERIAL AND METHODS

**Plant material and cultivation.** Barley plants (*Hordeum vulgare* L.) were cultivated (variety Kompakt from Slovakia) in plastic pots with soil substrate in a density of 390 plants per square meter in the natural environmental conditions within the vegetation cage of Dept. of Plant Physiology, Slovak University of Agriculture in Nitra (latitude 48°18'N, longitude 18°05'E). Experiments were carried out during the 2005 and 2006 growing seasons. Plants were fertilized and watered optimally, so that no symptoms of nutrient or water deficits were observed. Two variants of light regime were applied. In the first variant, plants were cultivated under natural light environment with full irradiation. The second variant, shaded, was simulated by non-weave textile restricting direct sunlight, where daily maximum did not exceed 20% of the total solar irradiation.

**Measurements.** During each growing season the microclimate conditions (photosynthetic active radiation, air and soil temperature, relative humidity) were monitored using a LiCor

Table 1. Climatic conditions of growing seasons 2005 and 2006: accumulated air temperature sum, average daily temperature and photosynthetic photon flux density (PPFD) measured over the canopy and in the simulated shaded conditions

	Accumulated temperature sum (°C)		Average daily temperature (°C)		Sum of PPFD over the canopy (mol/m <sup>2</sup> )		Sum of PPFD in the shade (mol/m <sup>2</sup> )	
	2005	2006	2005	2006	2005	2006	2005	2006
Growing season	1787	1783	18.2	18.7	2089.90	1963.78	275.89	302.93
April (14.4.–30.4.)	221	258	13.0	15.2	300.51	328.04	49.19	68.22
May	538	505	17.4	16.3	701.13	563.82	91.25	117.63
June	609	626	20.3	20.9	697.51	662.79	88.08	84.18
July (1.7.–20.7.)	419	394	21.0	23.2	390.75	409.13	47.43	32.90





Figure 1. Average daily air and soil temperatures measured at the beginning of the growing season (April) – from the plant sowing day up to the 3<sup>rd</sup> and 4<sup>th</sup> leaf occurrences

1400 datalogger (LiCor, Nebraska, USA). The temperature and humidity sensor was 5 cm far from soil surface and the thermometer was in the depth of 10 cm in the soil. The LI-190 sensors (LiCor, USA) measuring photon flux density of PAR were located above barley canopy, at two different levels inside the canopy (at the 4<sup>th</sup> and 5<sup>th</sup> leaf level) and in the simulated shaded environment. Data of air and soil temperature were recorded each minute, photosynthetic photon flux density each second; from them daily maximum, minimum and average values were calculated. Accumulated temperature was calculated for defined time interval as a sum of average daily air or soil temperatures. Accumulated photosynthetic photon flux density (PPFD) sum was calculated in a similar way.

Leaf growth was evaluated as a leaf area increase. This measurement was done in 10 repetitions per variant, until the leaf number 8 (flag leaf for sun and for shaded variant) on the main shoot was fully elongated (during all growing season). All leaves were numbered from stem basis in order as they had appeared. All measurements were carried out in a non-destructive way.



Leaf length and width were measured with a ruler. Then leaf area was calculated as:

$$A = l \times w \times k$$

where:  $l$  is leaf length,  $w$  is leaf width and  $k$  is a multiplying factor (0.64) usually used for barley (Šesták et al. 1971).

Leaf elongation was evaluated as a rate of leaf expansion per unit time (LER, mm/day). LER was estimated as maximal individual leaf length divided by the growth period (Singh et al. 2006). Leaf expansion duration (LED, day) of individual leaves was calculated as the number of days between the appearance and full emergence of an individual leaf (Bahmani et al. 2000).

**Statistical analysis.** All measured data were evaluated by the statistical packet of Microsoft Excel.

## RESULTS AND DISCUSSION

The course of climatic factors during both growing seasons is characterized in Table 1. Results show that accumulated air temperature sum and

Figure 2. Changes in leaf area of individual main stem barley leaves. Data were measured for sun and shaded variants during both growing seasons. Each data point represents the mean leaf area calculated from 10 repetitions. The vertical bars describe the standard deviations




Figure 3. Leaf expansion duration (LED, day) as a linear function of accumulated air temperature. Data were measured for leaves growing in conditions of full sunlight (full symbols) and in shaded conditions (empty symbols) during growing season 2005 and 2006

average daily air temperature were similar for both seasons. Substantial temperature differences were found at the beginning of both seasons, when average daily air temperature in 2005 was by 3°C lower than in 2006 (Figure 1).

This fact resulted in an increase of time interval between leaf appearance and leaf expansion starting in the case of the 1<sup>st</sup> and 2<sup>nd</sup> main stem leaves according to Kirby (1995) and Tamaki et al. (2002). Also, lower temperature was a reason for higher requirements of accumulated temperature sum for the first three leaves appearance in shaded variant. The rate of leaf initiation and leaf emergence had linear relationships with temperature

as was also found by Porter and Gawith (1999). On the other hand, the accumulated sum of photosynthetic active radiation during vegetation was 2089.90 and 1963.78 mol/m<sup>2</sup> for sun variant in 2005 and 2006, respectively; in the shaded variant the values measured were 275.89 and 302.93 mol/m<sup>2</sup> in 2005 and 2006, respectively, which was 13.2 and 15.4% of daily sum. These values reflected minimal differences in the amount of light incident on the canopy between both growing seasons.

Leaves on the main stem in shaded variant had larger leaf blade area than those at similar positions in sun variant. Leaf area increased from base of the main stem up to leaf position 5 for sun leaves




Figure 4. Relationship between the leaf expansion rate (LER, mm/day) and accumulated air temperature sum in barley leaves cultivated in different light regimes (full sunlight – full symbols; shade – empty symbols) in 2005 and 2006. Plotted data were measured for all main stem leaves




Figure 5. Leaf expansion rate (LER, mm/day) in relation to accumulated soil temperature sum for sun (full symbols) and shaded leaves (empty symbols). Data were measured from the 1<sup>st</sup> leaf position to the 4<sup>th</sup> leaf position on the main stem during 2005 and 2006

and leaf position 6 for shaded leaves (Figure 2) and then declined towards flag leaves which had the smallest leaf area. In 2006, a longer time interval of leaf expansion connected with higher accumulated temperature sum was observed for 6<sup>th</sup> shaded leaves than in 2005. This pattern was strongly associated with increasing leaf elongation rate and leaf width, as it was already confirmed for cereal crop species by Kirby et al. (1982) and Bos and Neuteboom (1998). Differences in individual leaf area in shaded variant was related to increased final leaf length and leaf elongation rate as was earlier found by Bahmani et al. (2000) and Evers et al. (2006).

Hay and Porter (2006) stated that under optimal conditions, duration of leaf expansion from its

emergence at a given node tends to be constant in thermal time. Our results indicated temperature influence on LED. A positive correlation between LED and accumulated air temperature sum for sun and shaded growing conditions was measured in 2005 and 2006 (Figure 3), where LED was a linear function of increased accumulated air temperatures with  $R^2$  values not less than 0.90. For leaf positions higher than position 5 for sun leaves and 6 for shade leaves, it was measured that maximal leaf length required shorter LED with lower accumulated temperature sum. Bultynck et al. (2004) mentioned that it is due to a higher extent of leaf elongation rate and leaf width of upper leaves and in several cases it could be associated with an increase of apical dome size.




Figure 6. Leaf expansion rate (LER, mm/day) in relation to accumulated photosynthetic active radiation sum plotted for all main stem leaves of sun and shaded variant during growing seasons 2005 and 2006

At an early stage of barley development, leaf elongation increased with main stem position, as was reported earlier (Kirby et al. 1982, Natr and Natrova 1992, Bultynck et al. 2004). In comparison with duration of leaf elongation, the rate of leaf expansion is more sensitive to fluctuating environment as was also stated by Hay and Porter (2006). LER decreased exponentially with raised accumulated temperature sum (Figure 4). Only small differences in correlations were measured between the sun and shaded variant, if LER was plotted to accumulated air temperature sum; however, when LER was plotted to accumulated soil temperature sum, stronger correlation was found for the first four leaves of the main stem in 2006 in sun and shaded plants with  $R^2 = 0.86$  and  $R^2 = 0.87$ , respectively (Figure 5). Results showed that soil temperature was more important for leaves emergence than air temperature, especially at the beginning of plant development as it is also confirmed by Hay and Wilson (1982) and McMaster et al. (2003).

Although the influence of temperature on leaf expansion rate has been already well described, the effect of light intensity on leaf expansion rate is still a matter of debate (Bos and Neuteboom 1998, Bahmani et al. 2000, Evers et al. 2006). Our results pointed out that in shading conditions barley leaf elongation rate was intensive and more sensitive to temperature in both experimental years. For similar values of LER as were measured in full sunlight conditions, only small interval of accumulated irradiance sum (20–90 mol/m<sup>2</sup>) was required to achieve the maximal leaf length in shaded plants (Figure 6).

Results showed that shaded environment consistently increased final leaf length mainly through increased leaf expansion rate and duration of leaf expansion. Both leaf growth parameters were dependent on accumulated temperature sum control. This suggests that a fully developed concept of the relationship of temperature to plant growth and development is an important aspect of dynamic crop simulation models.

## REFERENCES

- Abeledo L.G., Calderini D.F., Slafer G.A. (2004): Leaf appearance, tillering and their coordination in old and modern barleys from Argentina. *Field Crop Research*, 86: 23–32.
- Bahmani I., Hazard L., Varlet-Grancher C., Betin M., Lemaire G., Mattew C., Thom E.R. (2000): Differences in tillering of long- and short-leaved perennial ryegrass genetic lines under full light and shade treatments. *Crop Science*, 40: 1095–1102.
- Bartholomew P.W., Williams R.D. (2005): Cool-season grass development response to accumulated temperature under a range of temperature regimes. *Crop Science*, 45: 529–534.
- Bos H.J., Neuteboom J.H. (1998): Growth of individual leaves of spring wheat (*Triticum aestivum* L.) as influenced by temperature and light intensity. *Annals of Botany*, 81: 141–149.
- Bultynck L., Ter Steege M.W., Schortemeyer M., Poot P., Lambers H. (2004): From individual leaf elongation to whole shoot leaf area expansion: a comparison of three *Aegilops* and two *Triticum* species. *Annals of Botany*, 94: 99–108.
- Cao W., Moss D.N. (1989): Temperature effect on leaf emergence and phyllochron in wheat and barley. *Crop Science*, 29: 1018–1021.
- Evers J.B., Vos J., Andrieu B., Struik P.C. (2006): Cessation of tillering in spring wheat in relation to light interception and red: far red ratio. *Annals of Botany*, 97: 649–658.
- Hay R.K.M., Wilson G.T. (1982): Leaf appearance and extension in field-grown winter wheat plants: the importance of soil temperature during vegetative growth. *Journal of Agricultural Science*, 99: 403–410.
- Hay R.K.M., Porter J.R. (2006): *The Physiology of Crop Yield*. Blackwell Publishing, Oxford.
- Kirby E.J.M., Appeyard M., Fellowes G. (1982): Effect of sowing date on the temperature response of leaf emergence and leaf size in barley. *Plant, Cell and Environment*, 5: 477–484.
- Kirby E.J.M. (1995): Factors affecting rate of leaf emergence in barley and wheat. *Crop Science*, 35: 11–19.
- Miralles D.J., Richards R.A. (2000): Responses of leaf and tiller emergence and primordium initiation in wheat and barley to interchanged photoperiod. *Annals of Botany*, 85: 655–663.
- McMaster G.S., Wilhelm W.W., Palic D.B., Porter J.R., Jamieson P.D. (2003): Spring wheat leaf appearance and temperature: extending the paradigm? *Annals of Botany*, 91: 697–705.
- Natr L., Natrova Z. (1992): Characteristics of leaf growth in 6 varieties of spring wheat cultivated under constant conditions. *Rostlinná Výroba*, 38: 247–251. (In Czech)
- Porter J.R., Gawith M. (1999): Temperatures and the growth and development of wheat: a review. *European Journal of Agricultural*, 10: 23–36.
- Schurr U., Walter A., Rascher U. (2006): Functional dynamics of plant growth and photosynthesis – from steady-state to dynamics – from homogeneity to heterogeneity. *Plant, Cell and Environment*, 29: 340–352.

- Singh V., Pallaghy C.K., Singh D. (2006): Phosphorus nutrition and tolerance of cotton to water stress II. Water relations, free and bound water and leaf expansion rate. *Field Crop Research*, 96: 199–206.
- Šesták Z., Čatský J., Jarvis P.G. (1971): *Plant Photosynthetic Production: Manual of Methods*. Dr. W. Junk N.V. Publishers, Den Haag.
- Tamaki M., Kondo S., Itani T., Goto Y. (2002): Temperature responses of leaf emergence and leaf growth in barley. *Journal of Agricultural Science, Cambridge*, 138: 17–20.
- Tardieu F., Reymond M., Muller B., Granier C., Simonneau T., Sadok W., Welcker C. (2005): Linking physiological and genetic analyses of the control of leaf growth under changing environmental conditions. *Australian Journal of Agricultural Research*, 56: 937–946.
- Váňová M., Palík S., Hajšlová J., Burešová I. (2006): Grain quality and yield of spring barley in field trials under variable growing conditions. *Plant, Soil and Environment*, 52: 211–219.
- Wang Z.-M., Wei A.-L., Zheng D.-M. (2001): Photosynthetic characteristic of non-leaf organs of winter wheat cultivars differing in ear type and their relationship with grain mass per ear. *Photosynthetica*, 39: 239–244.

Received on May 7, 2009

---

*Corresponding author:*

Doc. Ing. Marián Brestič, CSc., Slovenská poľnohospodárska univerzita v Nitre, Tr. A. Hlinku 2, 949 76 Nitra, Slovenská republika  
phone: + 421 376 414 448, fax: + 421 377 411 451, e-mail: marian.brestic@uniag.sk

---