

<https://doi.org/10.17221/97/2017-CJGPB>

Correlations between *SmCPS1* Promoter Polymorphism and Tanshinone Contents in *Salvia miltiorrhiza*

YING SUN, XIN CHEN*, XIAO-YAN GAN, ZHU-YUN YAN,
DI-XIU MU and QING-RONG WANG

Chengdu University of Traditional Chinese Medicine, Chengdu, P.R. China

*Corresponding author: chenxin64@hotmail.com

Abstract

Sun Y., Chen X., Gan X.-Y., Yan Z.-Y., Mu D.-X., Wang Q.-R. (2018): Correlations between *SmCPS1* promoter polymorphism and tanshinone contents in *Salvia miltiorrhiza*. Czech J. Genet. Plant Breed., 54: 177–182.

Copalyl diphosphate synthase 1 (*SmCPS1*) is the first committed enzyme in tanshinone biosynthesis. The promoter region plays an important role in the transcriptional regulation of genes. Mutations in the promoter region may affect gene expression, resulting in changes in the amount of metabolites. In this study, we investigated the *SmCPS1* gene promoter region together with the 388 bp downstream from the translation start site and the content of tanshinones of 12 different genotypes of *Salvia miltiorrhiza*. The *cis*-elements of *SmCPS1* promoter were predicted and analysed by the Plant Transcriptional Regulatory Map database. We found (1) a different correlation between the polymorphism in the promoter region and the contents of tanshinones; (2) functional polymorphic loci – four tandem repeat variations, three indels and five single nucleotide polymorphisms (SNPs) in five *cis*-elements, three SNPs in exons and two SNPs in introns; (3) the correlation coefficient was higher when only functional (informative) polymorphic loci were considered. These findings have laid the foundation for further exploring the interspecific variation of *S. miltiorrhiza* and its relationship with the contents of tanshinones.

Keywords: *cis*-elements; copalyl diphosphate synthase 1; danshen; polymorphism

Salvia miltiorrhiza Bunge (danshen) is a medicinal model plant widely distributed throughout China (SONG *et al.* 2010). Its root is used for the treatment of cardiovascular and cerebrovascular diseases (YAN 2015). The main effective components of the root of *S. miltiorrhiza* are tanshinones, including dihydrotanshinone I, cryptotanshinone, 1,2-dihydrotanshinone, tanshinone IIA, tanshinone I and so on (ZENG *et al.* 2017).

Both sexual and vegetative propagation is involved in the cultivation of *S. miltiorrhiza*. But the crude field management of some origins of *S. miltiorrhiza* resulted in quality degradation and impaired medicinal value of the herb. Modern technologies, such as DNA markers based on Amplified Fragment Length Polymorphism (AFLP), Random Amplified Polymorphic DNA (RAPD), ISSR (Inter Simple Sequence Repeat) (SONG *et al.* 2010; ZHANG *et al.* 2013) and

Sequence Related Amplified Polymorphism (SRAP) (LI *et al.* 2008; WANG *et al.* 2010) were developed to identify different genotypes of *S. miltiorrhiza*. However, none of them is associated with important pharmacological ingredients. A more reliable trait for the identification of good quality danshen is required.

Former studies revealed the metabolic pathway of tanshinones (MA *et al.* 2012; YANG *et al.* 2013). The product of the 2-C-methyl-d-erythritol 4-phosphate (MEP) pathway, geranylgeranyl diphosphate (GGPP), is catalyzed by Copalyl diphosphate synthase 1 (*SmCPS1*) and Kaurene synthase like (*SmKSL*) together into miltiradiene (CUI *et al.* 2015), and then processed by CYP76AH1 to produce various tanshinone precursors (GUO *et al.* 2013). *SmCPS1* and *SmCPS2* have been confirmed to be involved in the tanshinone biosynthesis. But *SmCPS1* rather than *SmCPS2* plays the dominant role in tanshinone biosynthesis (CUI *et al.* 2015; LI *et al.*

<https://doi.org/10.17221/97/2017-CJGPB>

2017). Therefore, the regulation of *SmCPS1* will directly impact the downstream biosynthesis of tanshinones.

The promoter carries the basic information of the spatio-temporal expression pattern of the gene, which is the main regulatory region of the gene. The promoter is a region located upstream of the transcription start site and contains short sequences which are named *cis*-elements. Ranging from 5 to 30 bp in length (STEWART *et al.* 2012) and variable both in sequence and in location, *cis*-elements are the binding sites for corresponding transcriptional factors (TFs) and can be species-specific (WITTKOPP & KALAY 2011). The combination of the *cis*-elements determines the expression pattern of a gene.

By changing the binding capacity between the promoter elements and corresponding *trans*-factors, or the distance between co-functional elements, mutations from either substitution or insertion/deletion (indel) in the promoter can alter the expression pattern of a gene and subsequently lead to a phenotypic change (PHAM *et al.* 2017). Therefore, it is tempting to speculate that the variation of tanshinone contents of different *S. miltiorrhiza* genotypes is related to the promoter variation. In this study, we investigated the *SmCPS1* gene promoter region together with the 388bp downstream from the translation start site and the content of tanshinones of 12 different genotypes of *S. miltiorrhiza* to find the correlation between promoter variations and the content of tanshinones.

MATERIAL AND METHODS

Material. Samples were collected from the major origins of *S. miltiorrhiza* in China, these areas spanned the geographical range and have different

Table 1. Coordinates of sampling sites and GenBank accessions of *SmCPS1* sequences of each *Salvia miltiorrhiza* genotype

Sample	Latitude (N)	Longitude (E)	GenBank accession
01LS	33.7	111.29	MF409403
02LY	35.39	117.64	MF409404
03LT	34.35	109.85	MF409405
04QW	35.61	111.57	MF409406
05LCY	32.61	111.96	MF409407
06GD	30.99	104.5	MF409408
07SQ	33.48	112.99	MF409409
08ZJ	31.01	105	MF409410
09RC	34.71	110.81	MF409411
10SL2	33.84	109.92	MF409412
11JF	31.02	104.59	MF409413
12LQ	35.4	118.42	MF409414

ecological environments (Figure 1, Table 1). In order to eliminate the effect of the environment on the content of tanshinones, all samples were cultivated for 18 months in the Medicinal Botanical Garden of Chengdu University of Traditional Chinese Medicine before experiment.

Detection of tanshinone content. High performance liquid chromatography (HPLC) analysis was carried out on an Agilent 1200 Series HPLC instrument with Agilent ZORBAX Eclipse Plus C18 column (Analytical 4.6 × 250 mm, 5 μm; Agilent Technologies, USA), and Diode Array Detector (G315C 1260; Agilent Technologies). A gram of powder made from the fresh root of *S. miltiorrhiza* was used to extract the tanshinones with 10 ml methanol under ultrasonic conditions for 1 h. After cooling to room




Figure 1. Sampling sites of *Salvia miltiorrhiza* genotypes in China

<https://doi.org/10.17221/97/2017-CJGPB>

temperature, methanol was added to replenish the lost weight. Then the mixture was filtered through 0.45 µl microporous membrane. A volume of 20 µl of the liquid was used for detection. All peaks were detected at the wavelength of 270 nm. The column temperature was maintained at room temperature. The flow rate was 1.0 ml/min. Acetonitrile-water (V/V) was used as the mobile phase. The program was set with a linear gradient T from 0 to 30 min: T = 0 min, acetonitrile/water (55:45, V/V); T = 30 min, acetonitrile/water (0:100, V/V).

SmCPS1 promoter amplification. DNA was extracted using the Plant Genomic DNA Kit (DP305, Tiangen Biotech Co., Beijing). PCR was adopted to amplify the promoter region together with the downstream coding areas of *SmCPS1*. The detailed information about the primer pairs and the PCR reactions is in File S1 in Electronic Supplementary Material (ESM). The sequences were aligned and truncated to the same ends for further analysis. Their GenBank accessions are in Table 1.

Promoter analysis. The Binding Site Prediction Tool of Plant Transcriptional Regulatory Map database (<http://planttfdb.cbi.pku.edu.cn/prediction.php>) (JIN *et al.* 2017) was used to identify the *cis*-elements in the species-specific datasets that it provides. For overlapped predictions, a conservative point of view that different TFs can compete for the same binding site was taken; so different TFs were retained if they shared a common binding locus. But for the same TF, if it had overlapped binding sites, only the result with minimum *P*-value was retained. The aligned consensus sequence was used for prediction. Then the local sequences that excluded gaps were used for the second prediction to eliminate the prediction bias resulting from gaps.

The *cis*-element locus on which no alternative bases existed according to the position weight matrix of a *cis*-element was defined as a key binding site. Polymorphic loci in the key binding sites, exons and introns such as tandem repeat variations (TRV), single nucleotide polymorphisms (SNP) and indels were defined as functional polymorphic loci.

Statistical analysis. Sequence alignment was conducted using clustw by MEGA 7 (KUMAR *et al.* 2016) with default settings. The genetic distance matrix was generated by GenAlex (PEAKALL & SMOUSE 2012). Chemical component content distance matrices were generated using Euclidean distance. The Mantel test was conducted by GenAlex with a setting of 999 permutations.

RESULTS

Detection of tanshinone content. The contents of five tanshinones are shown in Figure 2 and Table S1 in ESM.

Promoter analysis and functional polymorphic loci identification. In the promoter upstream of untranslated region (5' UTR, +1), 33 regulatory regions were identified resulting from 81 TF and 117 TF binding sites (Table S2 in ESM).

A few mutation sites were found to be functional polymorphic loci (Table 2, Table S3 in ESM). These mutations included four TRVs, three indels and five SNPs in five *cis*-elements, three SNPs in exons and two SNPs in introns.

Correlations between the content of tanshinones and *SmCPS1* polymorphism. The Mantel test suggested a significant ($P < 0.05$) relationship between genetic distance matrix (Table S4 in ESM)




Figure 2. Tanshinone contents in *Salvia miltiorrhiza* genotypes (in mol/g)

Table 2. Characterization of functional polymorphic loci of *SmCPS1* and their variation pattern

Location*	Regions	Variation pattern	Type
–1685 to –1650	MIKC_MADS/C2H2	#	TRV
–1458	ARR-B	A/G	SNP
–1306	CPP	A/T	SNP
–1092 to –1091	MYB	T/C, A/C	SNP
–1070 to –1065	AP2/MIKC-MAC	(A)6/–	TRV
–1015 to –1003	TCP	AGTATGAACCACA/–	Indel
–687	Dof	T/–	Indel
–553	TCP	C/T	SNP
–541	MIKC_MADS	G/A	SNP
–269 to –267	C2H2	TAC/–	Indel
–204 to –142	(ATT) <i>n</i>	<i>n</i> = 0, 9, 12, 21	TRV
–22 to –21	TATA-box	AT/–	TRV
168	exon	C/T	SNP
179	intron	C/T	SNP
287	exon	G/C	SNP
328	exon	T/A	SNP
362	intron	T/C	SNP

*5' untranslated region (+1); # see Table S3 in ESM for details; SNP – single nucleotide polymorphism; TRV – tandem repeat variation; Indel – insertion/deletion variation

and content distance matrix of the chemical components (Table S5 in ESM) except for tanshinone I and dihydrotanshinone ($P > 0.05$, data not shown) (Table 3) when either all polymorphic sites or only functional polymorphic loci were considered. But the correlation coefficient was higher when only functional polymorphic loci were used. Cryptotanshinone showed the highest values of correlation coefficients; $R = 0.440$ when all polymorphic loci were counted and $R = 0.473$ when only functional polymorphic loci were counted. 1,2-dihydrotanshinone showed the lowest values of correlation coefficients ($R = 0.285$

for all polymorphic loci and $R = 0.297$ for functional polymorphic loci).

DISCUSSION

SmCPS1 is a key gene in the metabolism pathway of tanshinones. Any change of its expression will directly influence the content of tanshinones. Since a promoter contains the basic information about the expression pattern of the gene, we speculated that the variation of tanshinone contents of different *S. miltiorrhiza* genotypes is related to the promoter

Table 3. The Mantel test between genetic distance matrix and Euclidean distance matrix of chemical component content in *Salvia miltiorrhiza*

Chemical component	Genetic distance matrix based on all polymorphic loci				Genetic distance matrix based on functional polymorphic loci			
	P^*	function	$R^{\#}$	R^2	P^*	function	$R^{\#}$	R^2
1,2-dihydrotanshinone	0.028	$y = 1E-05x + 0.0021$	0.285	0.0811	0.023	$y = 2E-05x + 0.0022$	0.297	0.0883
Tanshinone IIA	0.034	$y = 1E-05x + 0.0018$	0.345	0.1193	0.026	$y = 2E-05x + 0.0018$	0.355	0.1261
Cryptotanshinone	0.042	$y = 9E-06x + 0.0007$	0.440	0.1939	0.038	$y = 1E-05x + 0.0007$	0.473	0.2233
Tanshinones [★]	0.018	$y = 2E-05x + 0.0031$	0.368	0.1352	0.029	$y = 2E-05x + 0.0031$	0.382	0.1459
Total tanshinones [◆]	0.018	$y = 4E-05x + 0.0047$	0.374	0.1399	0.024	$y = 5E-05x + 0.0047$	0.392	0.1536

#Mantel correlation coefficient; *significant level ($P < 0.05$); ★tanshinones is the combination of all 5 tanshinones that were investigated; ◆total tanshinones is the sum of contents of all 5 tanshinones

<https://doi.org/10.17221/97/2017-CJGPB>

polymorphism of *SmCPS1*. Our results that the polymorphic sites in *SmCPS1* promoter are in correlation with the variation of tanshinone contents except for tanshinone I and dihydrotanshinone supported the hypothesis.

Functional polymorphic loci are polymorphic loci located in the regulatory regions or coding regions of a gene and may change the expression pattern or function of the gene (KAGE *et al.* 2016). Changes in exons, *cis*-elements, or introns of a gene may change the function or expression pattern of the gene (LE HIR *et al.* 2003; NITHIANANTHARAJAH & HANNAN 2007; WITTKOPP & KALAY 2011). Hence the polymorphic loci such as SNPs, indels and TRVs were considered as functional polymorphic loci. According to the position weight matrix of *cis*-element, some sites contain alternative bases. Since no information about the effect of these alternative bases on the function of a *cis*-element was available, a key binding site was restricted on a *cis*-element locus where no alternative bases exist in this study. Such polymorphic loci in the key binding sites of *cis*-elements were considered as functional polymorphic loci.

Both genetic distance matrices generated based on all polymorphic loci and functional polymorphic loci showed the same significant relationships with the tanshinone contents. But the correlation coefficient was slightly higher when only functional polymorphic loci were considered. Considering the fact that a former study based on the whole genome polymorphism failed to find significant associations between DNA markers and the content of tanshinones (ZHANG *et al.* 2013), it is obvious that all polymorphic loci included in the calculation concealed a correlation between tanshinones and truly informative (functional) polymorphic loci. Hence, it is necessary to filter out informative polymorphic loci from non-informative ones when studying DNA markers.

Different correlations between *SmCPS1* functional polymorphism and tanshinone contents of *S. miltiorrhiza* may be the result of different distance between these chemical components and *SmCPS1* in the downstream pathway (YANG *et al.* 2013). If there is a stronger correlation suggesting a closer distance or a weaker correlation suggesting a longer distance between a chemical component and *SmCPS1*, our data suggest that the distances between the different tanshinones and *SmCPS1* are as follows: cryptotanshinone < tanshinone IIA < 1,2-dihydrotanshinone < tanshinone I and dihydrotanshinone. This supports part of the hypothesis from YANG *et al.* (2013)

about the relationship: cryptotanshinone < tanshinone IIA < tanshinone I. Though dihydrotanshinone and 1,2-dihydrotanshinone were not included in this hypothesis, our results provided them a probable place in the downstream pathway of tanshinone biosynthesis (see above).

This study is the first to find out correlations between the tanshinone content and polymorphism in *SmCPS1*. It exhibited the potential of *SmCPS1* sequence polymorphism to be developed as a new kind of DNA marker and will be of great value in the breeding and cultivation of *S. miltiorrhiza*. In conclusion, exploring the relationship between sequence polymorphism and phenotype provides us a new thread in establishing a DNA marker.

References

- Cui G., Duan L., Jin B., Qian J., Xue Z., Shen G., Snyder J.H., Song J., Chen S., Huang L., Peters R.J., Qi X. (2015): Functional divergence of diterpene syntheses in the medicinal plant *Salvia miltiorrhiza*. *Plant Physiology*, 169: 1607–1618.
- Guo J., Zhou Y.J., Hillwig M.L., Shen Y., Yang L., Wang Y., Zhang X., Liu W., Peters R.J., Chen X., Zhao Z.K., Huang L. (2013): CYP76AH1 catalyzes turnover of miltiradiene in tanshinones biosynthesis and enables heterologous production of ferruginol in yeasts. *Proceedings of the National Academy of Sciences of the USA*, 110: 12108–12113.
- Jin J., Tian F., Yang D. C., Meng Y. Q., Kong L., Luo J., Gao G. (2017): PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Research*, 45: D1040–D1045.
- Kage U., Kumar A., Dhokane D., Karre S., Kushalappa A.C. (2016): Functional molecular markers for crop improvement. *Critical Reviews in Biotechnology*, 36: 917–930.
- Kumar S., Stecher G., Tamura K. (2016): MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, 33: 1870–1874.
- Le Hir H., Nott A., Moore M.J. (2003): How introns influence and enhance eukaryotic gene expression. *Trends in Biochemical Sciences*, 28: 215–220.
- Li B., Cui G., Shen G., Zhan Z., Huang L., Chen J., Qi X. (2017): Targeted mutagenesis in the medicinal plant *Salvia miltiorrhiza*. *Scientific Reports*, 7: 43320.
- Li T., Fan H., Gao Z., Zhou Y., Yang H. (2008): Studies on genetic diversity of *Salvia miltiorrhiza* bunge with a molecular marker SRAP. *Journal of Nuclear Agricultural Sciences*, 22: 576–580.

<https://doi.org/10.17221/97/2017-CJGPB>

- Ma Y., Yuan L., Wu B., Li X., Chen S., Lu S. (2012): Genome-wide identification and characterization of novel genes involved in terpenoid biosynthesis in *Salvia miltiorrhiza*. *Journal of Experimental Botany*, 63: 2809–2823.
- Nithianantharajah J., Hannan A.J. (2007): Dynamic mutations as digital genetic modulators of brain development, function and dysfunction. *Bioessays*, 29: 525–535.
- Peakall R., Smouse P.E. (2012): GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research – an update. *Bioinformatics*, 28: 2537–2539.
- Pham T., Day S.M., Glassford W.J., Williams T.M., Rebeiz M. (2017): The evolutionary origination of a novel expression pattern through an extreme heterochronic shift. *Evolution & Development*, 19: 43–55.
- Song Z., Li X., Wang H., Wang J. (2010): Genetic diversity and population structure of *Salvia miltiorrhiza* Bge in China revealed by ISSR and SRAP. *Genetica*, 138: 241–249.
- Stewart A.J., Hannenhalli S., Plotkin J.B. (2012): Why transcription factor binding sites are ten nucleotides long. *Genetics*, 192: 973–985.
- Wang W., Shan C., Ni D., Wang Z. (2010): SRAP – Analysis of genetic diversity about germplasm in *Salvia miltiorrhiza* from different sources. *Chinese Traditional and Herbal Drugs*, 41: 632–635.
- Wittkopp P.J., Kalay G. (2011): Cis-regulatory elements: molecular mechanisms and evolutionary processes underlying divergence. *Nature Reviews Genetics*, 13: 59–69.
- Yan X. (2015): *Dan Shen (Salvia miltiorrhiza) in Medicine*. Beijing, Springer Netherlands and People's Medical Publishing House.
- Yang L., Ding G., Lin H., Cheng H., Kong Y., Wei Y., Fang X., Liu R., Wang L., Chen X., Yang C. (2013): Transcriptome analysis of medicinal plant *Salvia miltiorrhiza* and identification of genes related to tanshinone biosynthesis. *PLoS ONE*, 8: e80464.
- Zeng H., Su S., Xiang X., Sha X., Zhu Z., Wang Y., Guo S., Yan H., Qian D., Duan J. (2017): Comparative analysis of the major chemical constituents in *Salvia miltiorrhiza* roots, stems, leaves and flowers during different growth periods by UPLC-TQ-MS/MS and HPLC-ELSD methods. *Molecules*, 22: 771.
- Zhang Y., Li X., Wang Z. (2013): Diversity evaluation of *Salvia miltiorrhiza* using ISSR markers. *Biochemical Genetics*, 51: 707–721.

Received for publication July 10, 2017

Accepted after corrections March 5, 2018

Published online June 13, 2018