

<https://doi.org/10.17221/98/2019-VETMED>

## Optimisation of the lymphocyte proliferation assay in rainbow trout (*Oncorhynchus mykiss*)

HANA MINAROVA<sup>1,2\*</sup>, MIROSLAVA PALIKOVA<sup>1,3</sup>, JAN MARES<sup>3</sup>, EVA SYROVA<sup>1,2</sup>, JANA BLAHOVA<sup>4</sup>, MARTIN FALDYNA<sup>2</sup>, PETRA ONDRACKOVA<sup>2</sup>

<sup>1</sup>Department of Ecology and Diseases of Game, Fish and Bees, Faculty of Veterinary Hygiene and Ecology, University of Veterinary and Pharmaceutical Sciences Brno, Brno, Czech Republic

<sup>2</sup>Department of Immunology, Veterinary Research Institute, Brno, Czech Republic

<sup>3</sup>Department of Zoology, Fisheries, Hydrobiology and Apiculture, Faculty of AgriSciences, Mendel University in Brno, Brno, Czech Republic

<sup>4</sup>Department of Animal Protection, Welfare and Behaviour, Faculty of Veterinary Hygiene and Ecology, University of Veterinary and Pharmaceutical Sciences Brno, Brno, Czech Republic

\*Corresponding author: [minarova@vri.cz](mailto:minarova@vri.cz)

**Citation:** Minarova H, Palikova M, Mares J, Syrova E, Blahova J, Faldyna M, Ondrackova P (2019): Optimisation of the lymphocyte proliferation assay in rainbow trout (*Oncorhynchus mykiss*). Veterinarni Medicina 64, 547–557.

**Abstract:** The lymphocyte proliferation assay is a valuable method used for the evaluation of the fish immune system. However, there are many variations and optimal results are not always obtained. Unification is necessary to ensure the comparability between different studies. The aim of this study was to optimise the lymphocyte proliferation assay in rainbow trout (*Oncorhynchus mykiss*). This goal included the determination of the optimal incubation length, serum type, incubation temperature, type of mitogen and its concentration, and anticoagulant. The peripheral blood and head kidney lymphocytes were isolated by density gradient centrifugation. Subsequently, the cells were incubated for 3–8 days with different mitogens (pokeweed mitogen 5, 10 and 50 µg/ml, concanavalin A 1, 10 and 20 µg/ml, phytohaemagglutinin 25, 50 and 100 µg/ml, lipopolysaccharide 1, 50 and 100 µg/ml). The use of the different serum types (foetal bovine serum, trout serum), incubation temperatures (10–20 °C) and anticoagulants (heparin, EDTA) was compared. Labelled thymidine was used to evaluate the assay. The best results were obtained after seven days of incubation at 15 °C with foetal bovine serum (FBS). The head kidney lymphocytes showed the highest proliferative response with 50 µg/ml phytohaemagglutinin. With the peripheral blood lymphocytes (heparin and EDTA), the best results were obtained with 50 µg/ml pokeweed mitogen. The highest proliferation levels were detected with heparinised blood. In conclusion, optimisation of this assay contributes to the improved assessment of the rainbow trout immune function.

**Keywords:** fish; mitogen; stimulation; thymidine; transformation

Funded by the Internal Grant Agency of the University of Veterinary and Pharmaceutical Sciences Brno (IGA VFU Brno 216/2017/FVHE and 220/2018/FVHE), the Ministry of Agriculture of the Czech Republic (MZe NAZV QJ1510077) and the Ministry of Education, Youth and Sports of the Czech Republic (No. LO1218). This study was also supported by the project PROFISH CZ.02.1.01/0.0/0.0/16\_019/0000869. The project is financed by European Regional Development Fund in the operational programme VVV MŠMT.

<https://doi.org/10.17221/98/2019-VETMED>

Due to the rapid increase in the demand and production of fish meat, intensive aquaculture systems often deal with problems caused particularly by high stocking density and increased stress in fish. These conditions lead to various problems which are also reflected in the fish immune system – its ability to react can be examined by functional immunological assays (Kollner et al. 2002).

The lymphocyte proliferation assay was described by Nowell (1960). During the assay, lymphocytes are transformed into lymphoblasts after mitogenic stimulation and, subsequently, their proliferation activity is measured. Plant mitogens providing polyclonal activation of lymphocytes are most commonly used. Similarly to mammals, T lymphocytes are stimulated by phytohaemagglutinin and concanavalin, B lymphocytes are stimulated by lipopolysaccharide, and pokeweed mitogen stimulates both cell types, although primarily T blastogenesis was observed in rainbow trout (*Oncorhynchus mykiss*) by some authors (Janossy and Greaves 1971; Chilmoneczyk 1978; Kehrer et al. 1998). The purity of the mitogens may differ and may be the cause of the different results in some cases. The activation and proliferation of leukocytes is an important part of every immune response (Rycyzyn et al. 1998). This test provides a wide range of utilisation possibilities considering that the lymphocyte proliferation can be influenced by many factors, including diseases, pollutants, rearing conditions and stress (Espelid et al. 1996; Wendelaar Bonga 1997). It has been commonly used in mammals (Divya et al. 2015; Kralova-Kovarikova et al. 2016), but in fish, there are many different methodologies for various species and optimal results are not always obtained. The LTT (Lymphocyte Transformation Test) can be applied to assess the effect of pesticides on the lymphocyte viability and proliferation (Shelley et al. 2012), and is also often used in other toxicological studies (Harford et al. 2007; Muller et al. 2009) as well as nutritional studies (Siwicki et al. 2003; Kowalska et al. 2017).

Different incubation conditions were used in the previous studies with rainbow trout lymphocytes. The incubation time usually varies from 3 to 5 days, the shorter period being more common (Tahir and Secombes 1995; Siwicki et al. 2003; Shelley et al. 2012). Incubation with foetal bovine serum (FBS) as well as trout serum was described (Verlhac et al. 1998; Siwicki et al. 2003; Hebert et al. 2008). Temperature ranged from 15 °C to 22 °C (Tahir

and Secombes 1995; Lunden and Bylund 2000; Shelley et al. 2012). The usual mitogens are phytohaemagglutinin, concanavalin A and lipopolysaccharide (Flory and Bayne 1991; Siwicki et al. 2003), stimulation with pokeweed mitogen is less common (Ottinger and Kaattari 2000). In fish, mitogen concentrations vary from 1 µg/ml to 200 µg/ml (sometimes even 1000) (Chilmoneczyk 1978; Wang et al. 1997; Kehrer et al. 1998). The most commonly used samples include peripheral blood, the head kidney and spleen (O'Halloran et al. 1998; Shelley et al. 2012).

There are many possibilities when it comes to the evaluation of the lymphocyte proliferative response. One of the traditional options is the use of methyl-<sup>3</sup>H-thymidine (Yun et al. 2006; Harford et al. 2007; Muller et al. 2009). This method requires special equipment and involves work with radiolabelled substances, but it is sensitive and reliable. Another frequently applied evaluation method is the incorporation of the thymidine analogue 5-bromo-2'-deoxyuridine (BrdU) and its measurement by an ELISA-based assay (van der Heijden et al. 1995; Gauthier et al. 2003). It has been mainly used in mammals (Walsh et al. 2005; Karami et al. 2016) and less often in birds (Finkelstein et al. 2007; Grasman et al. 2013). The MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide) colorimetric assay is another alternative (Mosmann 1983; Siwicki et al. 2003; Kowalska et al. 2017), as well as the evaluation by flow cytometry (Scharsack et al. 2001; Shelley et al. 2012). However, these assays were mostly optimised in mammals, so they are often unreliable and have limitations like the detection range and other technical issues (Gauthier et al. 2003). Therefore, the use of labelled thymidine can be a suitable option. Also, in contrast to using a commercial kit, an unlimited number of samples can be examined.

The aim of this work was to optimise the lymphocyte proliferation assay in rainbow trout. To achieve this goal, the optimal length of incubation, serum type, incubation temperature, type of mitogen and its concentration and anticoagulant had to be determined.

## MATERIAL AND METHODS

**Fish.** An all-female population of rainbow trout (*Oncorhynchus mykiss*) was kept in an experimental

<https://doi.org/10.17221/98/2019-VETMED>

recirculating system situated at Mendel University in Brno (Czech Republic). The fish (average weight  $220.46 \pm 121.11$  g, total length  $24.91 \pm 3.53$  cm) were stocked in four breeding tanks, each with a volume of approximately 1000 l. A Nexus 310 biofilter and a UV-C lamp were used for water filtration and disinfection. All fish were fed two times a day with a BioMar EFICO Enviro 920 feed (BioMar, Denmark).

The average water temperature was  $17.5$  °C, the oxygen saturation was 87.2% (8.15 mg/l), the pH was 7.2,  $\text{N-NH}_4^+$  0.39 mg/l,  $\text{N-NO}_2^-$  0.18 mg/l and  $\text{Cl}^-$  145.83 mg/l.

The experiments were performed in accordance with the EU Directive 2010/63/EU for animal experiments. The Czech national regulations for animal experimentation were also followed (Act No. 246/1992 Coll.) and the experiments were approved by the Central Commission for Animal Welfare (MŠMT-19896/2015).

**Sample collection.** Every studied part of the test (the length of incubation, the serum, the temperature, the mitogens and the anticoagulants) was examined in a separate experiment (the mitogens in two), usually with eight (up to twelve) fish sampled. The data sample varies based on the numbers of the isolated cells. At first, some incubation parameters were selected based on pre-experiments. Blood (1 ml) was collected from the caudal vein into a heparinised (Zentiva, Czech Republic) syringe or into a syringe with EDTA (0.5M; ethylenediaminetetraacetic acid; Sigma-Aldrich, USA). The non-heparinised blood for the serum preparation was also obtained at this point. Consequently, the fish were humanely euthanised by a blow to the head, the samples of the head kidney were taken and placed into DPBS (Dulbecco's phosphate-buffered saline; Lonza, Switzerland). All samples were kept on ice and processed in less than two hours after collection. All the procedures described below were carried out in a laminar flow cabinet, under sterile conditions.

**Isolation of lymphocytes.** The head kidney samples were forced through a fine nylon mesh and washed by centrifugation in DPBS for  $2 \times 10$  min at 450 g. The blood samples were diluted 1 : 2 by RPMI-1640 Medium (Sigma-Aldrich, USA). Subsequently, all samples were layered on Histopaque (1077 g/ml; Sigma-Aldrich, USA) and the mononuclear cells were isolated by density gradient centrifugation at  $15$  °C for 40 min at 800 g.

The layer of mononuclear cells formed on the interface was collected and washed by centrifugation in DPBS for 10 min at 450 g. The isolated cells were counted by the BC-2800Vet haematology analyser (Mindray, China) and resuspended in L-15 Medium (Leibovitz, Sigma-Aldrich, USA) to the concentration of  $2 \times 10^5$  cells/well.

**Lymphocyte proliferation assay.** The lymphocyte suspension and different mitogens were dispensed into 96-well plates (Techno Plastic Products, Switzerland) making a total reaction volume of 200  $\mu$ l ( $2 \times 10^5$  cells). The heat inactivated (60 °C for 30 min) and sterile filtered foetal bovine serum (FBS; Hyclone; GE Healthcare Life Sciences, UK) or the heat inactivated (44 °C for 20 min as described by Sakai (1981)) autologous or homologous trout serum was added to each well (10%). The mitogens comprised 5, 10 and 50  $\mu$ g/ml pokeweed mitogen (PWM; Sigma-Aldrich, USA), 1, 10 and 20  $\mu$ g/ml concanavalin A (ConA; Sigma-Aldrich, USA), 25, 50 and 100  $\mu$ g/ml phytohaemagglutinin (PHA; Thermo Fisher Scientific, USA) and 1, 50 and 100  $\mu$ g/ml lipopolysaccharide (LPS; prepared by phenol extraction from *Actinobacillus pleuropneumoniae*). During the preliminary experiments, two times higher than the highest and two times lower than the lowest concentration of each mitogen were used. All samples including the non-stimulated negative controls were run in triplicates. The lymphocytes were placed in an incubator (Memmert, Germany) at 10 °C, 15 °C or 20 °C. After 3–8 days, 50  $\mu$ l of  $^3\text{H}$ -thymidine (5  $\mu$ Ci/ml; American Radiolabeled Chemicals, USA) was added and incorporated into the newly formed DNA of the proliferating lymphocytes. After 20 h, the cells were harvested by the FilterMate Harvester (Packard, USA) and the proliferation activity was measured as counts per minute (cpm) by the TopCount NXT liquid scintillation counter (Packard, USA). The stimulation index (SI; the mean ratio of the stimulated and non-stimulated lymphocytes) was calculated.

**Statistical analyses.** For the statistical analyses, the Shapiro-Wilk test was used to determine if the data are normally distributed. Subsequently, the Mann-Whitney *U* test or Student's *t*-test, and the multi-sample median test or Tukey's HSD (Honestly Significant Difference) test was performed, depending on the parametricity of the data. The means and standard errors of the means (SEM) were calculated;  $P < 0.05$  was considered significant.

## RESULTS

### Length of incubation

The optimal length of incubation was investigated. Due to the different numbers of the isolated cells from each fish, the cells were primarily incubated for 5–7 days, resulting in a lower data sample when a longer time period is included in the statistics evaluation. The incubation of the head kidney lymphocytes was carried out for 8 days (with the cell proliferation measured during day 3–8) with pokeweed mitogen (10 µg/ml) and foetal bovine serum at 15 °C (the mitogen, temperature and serum were selected based on the pre-experiments). The best results were obtained when incubated for 7 days (SI 24.72). There was a significant difference between the 3- and 6-day incubation ( $P < 0.05$ ;




Figure 1. The effect of the incubation time (3–8 days) on the proliferation of the head kidney lymphocytes (A) and the peripheral blood lymphocytes collected using heparin (B). The cells were incubated at 15 °C with pokeweed mitogen (10 µg/ml) and foetal bovine serum. The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values (means ± SEM, A:  $n = 4$ ; B:  $n = 5$ ) with the different superscript letters are significantly different ( $P < 0.05$ )

Figure 1A). With the individual statistical analysis of the 6- and 7-day incubation (with a larger data sample), higher values were still obtained after 7 days (SI 32.51); however, the differences were not statistically significant (Figure 2).

With the peripheral blood lymphocytes (collected using heparin), the highest proliferation levels were detected after 8 days of incubation (SI 47.04). The cells were incubated for 8 days at 15 °C with pokeweed mitogen (10 µg/ml) and FBS. A significant difference was detected between 3 days and 7 + 8 days of incubation ( $P < 0.05$ ; Figure 1B). However, when 5–8 days of incubation were analysed separately (with a larger data sample), the highest values were obtained at 7 days (SI 61.23; Figure 3).

### Serum

The optimal serum type was investigated – incubation with FBS, autologous and homologous trout serum was compared. The head kidney lymphocytes were incubated for 7 days at 15 °C with 100 µg/ml




Figure 2. The effect of the incubation time (6–7 days) on the proliferation of the head kidney lymphocytes. The cells were incubated at 15 °C with pokeweed mitogen (10 µg/ml) and foetal bovine serum. The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values are the means ± SEM ( $n = 12$ ), no significant differences were detected

<https://doi.org/10.17221/98/2019-VETMED>




Figure 3. The effect of the incubation time (5–8 days) on the proliferation of the peripheral blood lymphocytes collected using heparin. The cells were incubated at 15 °C with pokeweed mitogen (10 µg/ml) and foetal bovine serum. The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values are the means ± SEM ( $n = 7$ ), no significant differences were detected




Figure 4. The effect of the different sera (foetal bovine serum – FBS, autologous or homologous trout serum) on the proliferation of the head kidney lymphocytes. The cells were incubated for 7 days at 15 °C with phytohaemagglutinin (100 µg/ml) and the different sera. The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values (means ± SEM) with the different superscript letters are significantly different ( $n = 6$ ;  $P < 0.01$ )

phytohaemagglutinin (the selected temperature and mitogen were based on the pre-experiments); FBS showed significantly higher proliferation levels than trout serum ( $P < 0.01$ ; Figure 4).

### Incubation temperature

The optimal incubation temperature was determined. The head kidney lymphocytes were incubated at 10 °C, 15 °C, and 20 °C for 7 days with FBS and 100 µg/ml phytohaemagglutinin (the selected mitogen was based on the pre-experiments). There were no significant differences between the different incubation temperatures (Figure 5A).




Figure 5. The effect of the incubation temperature (10, 15 and 20 °C) on the lymphocyte proliferation. The head kidney lymphocytes (A) were incubated for 7 days with foetal bovine serum and phytohaemagglutinin (100 µg/ml), the peripheral blood lymphocytes collected using heparin (B) were incubated with pokeweed mitogen (50 µg/ml). The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values are the means ± SEM ( $n = 8$ ), no significant differences were detected


<https://doi.org/10.17221/98/2019-VETMED>

Figure 6. The effect of the different mitogens on the proliferation of the head kidney lymphocytes (A) and the peripheral blood lymphocytes collected using heparin (B) – pokeweed mitogen (PWM; 5, 10, 50  $\mu\text{g/ml}$ ), phytohaemagglutinin (PHA; 25, 50, 100  $\mu\text{g/ml}$ ) and concanavalin A (ConA; 1, 10, 20  $\mu\text{g/ml}$ ). The cells were incubated for 7 days at 15  $^{\circ}\text{C}$  with foetal bovine serum and the different mitogens. The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values (means  $\pm$  SEM, A:  $n = 6$ ; B:  $n = 8$ ) with the different superscript letters are significantly different (A: no significant differences were detected; B:  $P < 0.05$ )

The peripheral blood lymphocytes collected with the use of heparin were incubated for 7 days with FBS and pokeweed mitogen (50  $\mu\text{g/ml}$ ). The highest proliferation levels were measured with the incubation carried out at 15  $^{\circ}\text{C}$  (SI 83.16; Figure 5B).

#### Mitogens and their concentrations

Subsequently, the optimal mitogens and their concentrations were determined. The cells were incubated for 7 days at 15  $^{\circ}\text{C}$  with FBS and the different mitogens (2.4.). The head kidney lymphocytes




Figure 7. The effect of the different mitogens on the proliferation of the head kidney lymphocytes (A) – pokeweed mitogen (PWM; 10  $\mu\text{g/ml}$ ) and lipopolysaccharide (LPS; 1, 50, 100  $\mu\text{g/ml}$ ). The effect of the different mitogens on the proliferation of the peripheral blood lymphocytes collected using heparin (B) – pokeweed mitogen (PWM; 50  $\mu\text{g/ml}$ ) and lipopolysaccharide (LPS; 1, 50, 100  $\mu\text{g/ml}$ ). The cells were incubated for 7 days at 15  $^{\circ}\text{C}$  with foetal bovine serum and the different mitogens. The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values (means  $\pm$  SEM,  $n = 8$ ) with the different superscript letters are significantly different (A:  $P < 0.05$ ; B:  $P < 0.01$ )

<https://doi.org/10.17221/98/2019-VETMED>

showed very low results with concanavalin A (no stimulation was observed) compared to pokeweed mitogen and phytohaemagglutinin (Figure 6A). The highest values were detected with 50 µg/ml phytohaemagglutinin. Pokeweed mitogen (10 µg/ml) also provided significantly higher proliferation levels than 1 µg/ml lipopolysaccharide ( $P < 0.05$ ; Figure 7A).

The peripheral blood lymphocytes isolated from heparinised blood showed the best results with 50 µg/ml pokeweed mitogen (SI 49.09), with a significant difference compared to its other concentrations, phytohaemagglutinin and concanavalin A ( $P < 0.05$ ). With concanavalin A, almost no stimulation was observed (Figure 6B). In comparison with 1 µg/ml lipopolysaccharide, 50 µg/ml pokeweed mitogen provided a significantly higher proliferative response (11 536.88 cpm/SI 303.68;  $P < 0.01$ ; Figure 7B).

Similarly to the heparinised blood, the peripheral blood lymphocytes collected with the use of EDTA (incubated for 6 days – the optimal incubation time based on the pre-experiments) were well stimulated by 50 µg/ml pokeweed mitogen (SI 44.25), with a significant difference compared to 50 µg/ml phytohaemagglutinin ( $P < 0.01$ ), where no stimulation was observed (Figure 8).

### Anticoagulants

The proliferation activity of the blood samples collected using heparin or EDTA was compared. The cells isolated from peripheral blood were incubated for 7 days at 15 °C with foetal bovine serum and pokeweed mitogen (50 µg/ml). The heparinised blood significantly showed the best results (SI 278.08;  $P < 0.01$ ), approximately 6 times higher than those detected with the use of EDTA (Figure 9).

### DISCUSSION

In other studies, a shorter incubation period (3–4 days) has usually been used (Chilmonczyk 1978; Kehrer et al. 1998; Gauthier et al. 2003; Siwicki et al. 2003; Muller et al. 2009; Shelley et al. 2012). In mammals, lymphocytes are also generally incubated for 2–4 days (Yaqoob et al. 1994; Kearns et al. 1999; Yun et al. 2006; Divya et al. 2015). However, our values were usually higher than those obtained in the studies using shorter (suboptimal) incuba-




Figure 8. The effect of the different mitogens on the proliferation of the peripheral blood lymphocytes collected using EDTA – pokeweed mitogen (PWM; 50 µg/ml) and phytohaemagglutinin (PHA; 50 µg/ml). The cells were incubated for 6 days at 15 °C with foetal bovine serum and the different mitogens. The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values (means ± SEM) with the different superscript letters are significantly different ( $n = 10$ ;  $P < 0.01$ )




Figure 9. The effect of the different anticoagulants (heparin – PBL<sub>H</sub> or EDTA – PBL<sub>E</sub>) on the proliferation of the peripheral blood lymphocytes. The peripheral blood lymphocytes were incubated for 7 days at 15 °C with foetal bovine serum and pokeweed mitogen (50 µg/ml). The stimulation index (SI) is the mean ratio of the stimulated and non-stimulated lymphocytes (measured as counts per minute). The values (means ± SEM) with the different superscript letters are significantly different ( $n = 9$ ;  $P < 0.01$ )

<https://doi.org/10.17221/98/2019-VETMED>

tion periods (Chilmonczyk 1978; Flory and Bayne 1991; Tahir and Secombes 1995; Lin et al. 1999; Gauthier et al. 2003; Leonardi and Klempau 2003; Hebert et al. 2008). Similar observations to ours were described by DeKoning and Kaattari (1991) and Reitan and Thuvander (1991). In their studies, longer incubation periods (5–6 days) provided higher stimulation levels.

As for the mitogens, the best results were acquired with pokeweed mitogen (11 536.88 cpm/SI 303.68). These values are comparable to the ones observed in mammals (Yaqoob et al. 1994; Wagner et al. 1999). Since pokeweed mitogen stimulates both B and T lymphocytes, the highest proliferation activity was expected. In our study, the T cell mitogen phytohaemagglutinin showed better results than the B cell mitogen lipopolysaccharide. This observation indicates that, in rainbow trout, T cells are the predominant leucocyte population stimulated by pokeweed mitogen, as observed by other authors (Kehrer et al. 1998). The use of concanavalin A resulted in a very low lymphocyte stimulation compared to the other mitogens. Different studies obtained similar proliferative responses using this mitogen (Gauthier et al. 2003; Leonardi and Klempau 2003). In contrast to our findings, Reitan and Thuvander (1991) observed no mitogenic effect of higher concentrations of PHA on peripheral blood lymphocytes. According to Etlinger et al. (1976), head kidney cells could be stimulated by LPS but not by ConA. The differences between the lymphocyte mitogenesis detected in the peripheral blood and head kidney lymphocytes were also described by Daly et al. (1995). The lymphocytes collected with the use of EDTA (inhibitor of cell activity) showed no proliferation with phytohaemagglutinin – the effect which has also been observed by other authors (Alford 1970). Generally, high concentrations of mitogens were usually more effective than the low concentrations in our study.

No stimulation was observed with autologous or homologous serum, although DeKoning and Kaattari (1991) describe different findings with homologous rainbow trout plasma. However, the values detected in the studies using homologous serum (Chilmonczyk 1978) were markedly lower compared to ours. Therefore, FBS (commonly used in mammals) seems to provide better proliferation conditions for the rainbow trout lymphocytes. It was used in many studies (Tillitt et al. 1988; Kehrer et al. 1998; Gauthier et al. 2003;

Muller et al. 2009; Shelley et al. 2012), although some authors preferred autologous or homologous serum (Daly et al. 1995; Rycyzyn et al. 1998).

The most commonly used incubation temperature is 15–20 °C for rainbow trout lymphocytes (Kehrer et al. 1998; Gauthier et al. 2003; Muller et al. 2009; Shelley et al. 2012). Similarly, our data yielded the best results at 15 °C, with a decrease in the proliferation activity at 10 °C. Some authors use even higher incubation temperatures in rainbow trout (Chilmonczyk 1978; Siwicki et al. 1996; Verlhac et al. 1998; Lunden and Bylund 2000; Siwicki et al. 2003; Hebert et al. 2008). Because temperature is one of the most important factors in fish, its maintenance can be crucial (Bly and Clem 1992; Hardie et al. 1994).

The samples most commonly used in the lymphocyte proliferation assay comprise peripheral blood (van der Heijden et al. 1995; Kehrer et al. 1998; Rycyzyn et al. 1998; Gauthier et al. 2003; Shelley et al. 2012) and the head kidney (Gauthier et al. 2003; Siwicki et al. 2003; Kowalska et al. 2017). As the main haematopoietic organ in teleosts, the head kidney shows some proliferation activity even in the non-stimulated cells (Gauthier et al. 2003), consequently providing results similar to EDTA (cell activity inhibitor). In our experiment, the values (SI 278.08) obtained with the heparinised blood were significantly higher compared to most other studies (Chilmonczyk 1978; Tillitt et al. 1988).

The lymphocyte proliferation assay is a reliable method of examining the fish immune function. However, there are many variables involved and various authors have used different incubation lengths, temperatures, serum types and types of mitogens. Different results may be obtained due to the various conditions used during this assay. Therefore, unification of this method is essential to ensure comparability. In rainbow trout (*Oncorhynchus mykiss*), a longer incubation period (7 days) is necessary to obtain sufficient proliferation levels. Incubation at 15 °C and the use of FBS instead of trout serum is advisable. Examination of both peripheral blood cells and head kidney cells is appropriate to acquire a complex view of the immune response. Use of heparin instead of EDTA as anticoagulant is highly recommended. Based on our results, 50 µg/ml phytohaemagglutinin is the optimal T cell mitogen (in contrast to concanavalin A), 50–100 µg/ml lipopolysaccharide stimulates B cells and stimula-



<https://doi.org/10.17221/98/2019-VETMED>

tion of both cell types is best provided by 50 µg/ml pokeweed mitogen. Due to the different cell representation in each sample, the peripheral blood lymphocytes are best stimulated by 50 µg/ml pokeweed mitogen while the head kidney lymphocytes show the highest proliferation levels with 50 µg/ml phytohaemagglutinin. Optimisation of this assay improves its utilisation in determining the fish health.

## Acknowledgement

The authors would like to thank Ing. Lenka Leva for technical support, Ing. Ludmila Faldikova for her linguistic assistance and Prof. Jiri Pikula for his critical review of this manuscript.

## REFERENCES

- Alford RH (1970): Metal cation requirements for phytohemagglutinin-induced transformation of human peripheral blood lymphocytes. *Journal of Immunology* (Baltimore, MD: 1950) 104, 698–703.
- Bly JE, Clem LW (1992): Temperature and teleost immune functions. *Fish & Shellfish Immunology* 2, 159–171.
- Chilmonczyk S (1978): In vitro stimulation by mitogens of peripheral blood lymphocytes from rainbow trout (*Salmo gairdneri*). *Annales d'Immunologie* 129, 3–12.
- Daly JG, Moore AR, Olivier G (1995): A colorimetric assay for the quantification of brook trout (*Salvelinus fontinalis*) lymphocyte mitogenesis. *Fish & Shellfish Immunology* 5, 265–273.
- DeKoning J, Kaattari S (1991): Mitogenesis of rainbow trout peripheral blood lymphocytes requires homologous plasma for optimal responsiveness. *In Vitro Cellular & Developmental Biology: Journal of the Tissue Culture Association* 27A, 381–386.
- Divya KB, Sathish MHK, Kapila S, Sabikhi L (2015): Immunosuppressive potential of low fat buffalo milk supplemented with omega-3 fatty acids. *Food and Agricultural Immunology* 26, 558–565.
- Espelid S, Lokken GB, Steiro K, Bogwald J (1996): Effects of cortisol and stress on the immune system in Atlantic Salmon (*Salmo salar* L.). *Fish & Shellfish Immunology* 6, 95–110.
- Etlinger HM, Hodgins HO, Chiller JM (1976): Evolution of the lymphoid system. I. Evidence for lymphocyte heterogeneity in rainbow trout revealed by the organ distribution of mitogenic responses. *Journal of Immunology* (Baltimore, MD: 1950) 116, 1547–1553.
- Finkelstein ME, Grasman KA, Croll DA, Tershy BR, Keitt BS, Jarman WM, Smith DR (2007): Contaminant-associated alteration of immune function in black-footed albatross (*Phoebastria nigripes*), a North Pacific predator. *Environmental Toxicology and Chemistry* 26, 1896–1903.
- Flory CM, Bayne CJ (1991): The influence of adrenergic and cholinergic agents on the chemiluminescent and mitogenic responses of leukocytes from the rainbow trout, *Oncorhynchus mykiss*. *Developmental & Comparative Immunology* 15, 135–142.
- Gauthier DT, Cartwright DD, Densmore CL, Blazer VS, Ottinger CA (2003): Measurement of in vitro leucocyte mitogenesis in fish: ELISA based detection of the thymidine analogue 5'-bromo-2'-deoxyuridine. *Fish & Shellfish Immunology* 14, 279–288.
- Grasman KA, Echols KR, May TM, Peterman PH, Gale RW, Orazio CE (2013): Immunological and reproductive health assessment in herring gulls and black-crowned night herons in the Hudson-Raritan Estuary. *Environmental Toxicology and Chemistry* 32, 548–561.
- Hardie LJ, Fletcher TC, Secombes CJ (1994): Effect of temperature on macrophage activation and the production of macrophage activating factor by rainbow trout (*Oncorhynchus mykiss*) leucocytes. *Developmental & Comparative Immunology* 18, 57–66.
- Harford AJ, O'Halloran K, Wright PFA (2007): Effect of in vitro and in vivo organotin exposures on the immune functions of murray cod (*Maccullochella peelii peelii*). *Environmental Toxicology and Chemistry* 26, 1649–1656.
- Hebert N, Gagne F, Cejka P, Cyr D, Marcogliese DJ, Blaise C, Pellerin J, Fournier M (2008): The effects of a primary-treated municipal effluent on the immune system of rainbow trout (*Oncorhynchus mykiss*): Exposure duration and contribution of suspended particles. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 148, 258–264.
- Janossy G, Greaves MF (1971): Lymphocyte activation. I. Response of T and B lymphocytes to phytomitogens. *Clinical & Experimental Immunology* 9, 483–498.
- Karami Z, Mesdaghi M, Karimzadeh P, Mansouri M, Taghdiri MM, Kayhanidoost Z, Jebelli B, Shekarriz Foumani R, Babaie D, Chavoshzadeh Z (2016): Evaluation of Lymphocyte Transformation Test Results in Patients with Delayed Hypersensitivity Reactions following the Use of Anticonvulsant Drugs. *International Archives of Allergy and Immunology* 170, 158–162.
- Kearns RJ, Hayek MG, Turek JJ, Meydani M, Burr JR, Greene RJ, Marshall CA, Adams SM, Borgert RC, Reinhart GA (1999): Effect of age, breed and dietary omega-6 (n-6): Omega-3 (n-3) fatty acid ratio on immune function, eicosanoid production, and lipid peroxidation in young and

<https://doi.org/10.17221/98/2019-VETMED>

- aged dogs. *Veterinary Immunology and Immunopathology* 69, 165–183.
- Kehrer SR, Hannan CM, Raison RL (1998): Identification of a subpopulation of leucocytes from the rainbow trout (*Oncorhynchus mykiss*) responsive to pokeweed mitogen. *Fish & Shellfish Immunology* 8, 477–487.
- Kollner B, Wasserrab B, Kotterba G, Fischer U (2002): Evaluation of immune functions of rainbow trout (*Oncorhynchus mykiss*)—how can environmental influences be detected? *Toxicology Letters* 131, 83–95.
- Kowalska A, Siwicki AK, Kowalski RK (2017): Dietary resveratrol improves immunity but reduces reproduction of broodstock medaka *Oryzias latipes* (Temminck & Schlegel). *Fish Physiology and Biochemistry* 43, 27–37.
- Kralova-Kovarikova S, Leva L, Knotek Z, Toman M (2016): Changes in lymphocyte function and subset counts in cats with spontaneous chronic kidney disease. *Veterinarni Medicina* 61, 553–559.
- Leonardi MO, Klempau AE (2003): Artificial photoperiod influence on the immune system of juvenile rainbow trout (*Oncorhynchus mykiss*) in the Southern Hemisphere. *Aquaculture* 221, 581–591.
- Lin SH, Ellis AE, Davidson GA, Secombes CJ (1999): Migratory, respiratory burst and mitogenic responses of leucocytes isolated from the gills of rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology* 9, 211–226.
- Lunden T, Bylund G (2000): The influence of in vitro and in vivo exposure to antibiotics on mitogen-induced proliferation of lymphoid cells in rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology* 10, 395–404.
- Mosmann T (1983): Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* 65, 55–63.
- Muller C, Ruby S, Brousseau P, Cyr D, Fournier M, Gagné F (2009): Immunotoxicological effects of an activated-sludge-treated effluent on rainbow trout (*Oncorhynchus mykiss*). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 150, 390–394.
- Nowell PC (1960): Phytohemagglutinin: An Initiator of Mitosis in Cultures of Normal Human Leukocytes. *Cancer Research* 20, 462–466.
- O'Halloran K, Ahokas JT, Wright PFA (1998): Response of fish immune cells to in vitro organotin exposures. *Aquatic Toxicology* 40, 141–156.
- Ottinger CA, Kaattari SL (2000): Long-term immune dysfunction in rainbow trout (*Oncorhynchus mykiss*) exposed as embryos to aflatoxin B<sub>1</sub>. *Fish & Shellfish Immunology* 10, 101–106.
- Reitan L, Thuvander A (1991): In vitro stimulation of salmonid leucocytes with mitogens and with *Aeromonas salmonicida*. *Fish & Shellfish Immunology* 1, 297–307.
- Rycyzyn MA, Wilson MR, Bengten E, Warr GW, Clem LW, Miller NW (1998): Mitogen and growth factor-induced activation of a STAT-like molecule in channel catfish lymphoid cells. *Molecular Immunology* 35, 127–136.
- Sakai DK (1981): Heat inactivation of complements and immune hemolysis reactions in rainbow trout, masu salmon, coho salmon, goldfish and tilapia. *Bulletin of the Japanese Society of Scientific Fisheries* 47, 565–571.
- Scharsack JP, Steinhagen D, Leibold W, Rabe U, Korting W, Schuberth H-J (2001): Flow cytometric analysis of mitogen-induced activation of rainbow trout (*Oncorhynchus mykiss*) peripheral blood leucocytes. *Journal of Veterinary Medicine Series B* 48, 331–339.
- Shelley LK, Ross PS, Kennedy CJ (2012): Immunotoxic and cytotoxic effects of atrazine, permethrin and piperonyl butoxide to rainbow trout following in vitro exposure. *Fish & Shellfish Immunology* 33, 455–458.
- Siwicki AK, Miyazaki T, Komatsu I, Matsusato T (1996): In vitro influence of heat extract from firefly squid *Watasenia scintillans* on the phagocyte and lymphocyte activities in rainbow trout *Oncorhynchus mykiss*. *Fish Pathology* 31, 1–7.
- Siwicki AK, Morand M, Fuller J, Nissen S, Goryczko K, Ostaszewski P, Kazun K, Glombski E (2003): Influence of feeding the leucine metabolite  $\beta$ -hydroxy- $\beta$ -methylbutyrate (HMB) on the non-specific cellular and humoral defence mechanisms of rainbow trout (*Oncorhynchus mykiss*). *Journal of Applied Ichthyology* 19, 44–48.
- Tahir A, Secombes CJ (1995): The effects of diesel oil-based drilling mud extracts on immune responses of rainbow trout. *Archives of Environmental Contamination and Toxicology* 29, 27–32.
- Tillitt DE, Giesy JP, Fromm PO (1988): In vitro mitogenesis of peripheral blood lymphocytes from rainbow trout (*Salmo gairdneri*). *Comparative Biochemistry and Physiology Part A: Comparative Physiology* 89, 25–35.
- van der Heijden MHT, Booms GHR, Tanck MWT, Rombout JHWM, Boon JH (1995): Influence of flumequine on in vivo mitogen responses of European eel (*Anguilla anguilla* L., 1758) lymphoid cells. *Veterinary Immunology and Immunopathology* 47, 143–152.
- Verlhac V, Obach A, Gabaudan J, Schuep W, Hole R (1998): Immunomodulation by dietary vitamin C and glucan in rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology* 8, 409–424.
- Wagner U, Burkhardt E, Failing K (1999): Evaluation of canine lymphocyte proliferation: comparison of three different colorimetric methods with the <sup>3</sup>H-thymidine

<https://doi.org/10.17221/98/2019-VETMED>

- incorporation assay. *Veterinary Immunology and Immunopathology* 70, 151–159.
- Walsh CJ, Luer CA, Noyes DR (2005): Effects of environmental stressors on lymphocyte proliferation in Florida manatees, *Trichechus manatus latirostris*. *Veterinary Immunology and Immunopathology* 103, 247–256.
- Wang C, Noga EJ, Avtalion R, Levy MG (1997): Whole blood assay for examining lymphocyte blastogenesis of percichthyid bass (Morone mrr01). *Veterinary Immunology and Immunopathology* 58, 355–362.
- Wendelaar Bonga SE (1997): The stress response in fish. *Physiological Reviews* 77, 591–625.
- Yaqoob P, Newsholme EA, Calder PC (1994): The effect of dietary lipid manipulation on rat lymphocyte subsets and proliferation. *Immunology* 82, 603–610.
- Yun C-H, Son CG, Jung U, Han SH (2006): Immunosuppressive effect of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) through the inhibition of T-lymphocyte proliferation and IL-2 production. *Toxicology* 217, 31–38.

Received: July 30, 2019

Accepted after corrections: October 25, 2019